Table 9 Reduced Form Effects on Weekly Nutrition for Ages 1-3: Youngest Child

Age reference: 16 December 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. Food outcomes require an age above six calendar months and recorded codes 1-9. Codes 1-3 indicate at least weekly consumption. Ever breastfed uses recorded codes 1 and 2. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.938 P=1,502	-0.008 [0.023] N=1,602	0.010 [0.022] N=1,478	0.003 [0.024] N=1,333	0.001 [0.026] N=1,133	-0.009 [0.031] N=929
Vegetables consumed at least weekly	0.901 P=1,443	0.013 [0.028] N=1,602	0.016 [0.028] N=1,478	0.001 [0.029] N=1,333	0.006 [0.029] N=1,133	-0.022 [0.033] N=929
Meat fish or poultry consumed at least weekly	0.796 P=1,281	**0.076* [0.043] N=1,602**	**0.097** [0.045] N=1,478**	**0.086* [0.049] N=1,333**	0.061 [0.054] N=1,133	0.047 [0.062] N=929
Legumes consumed at least weekly	0.795 P=1,277	-0.048 [0.039] N=1,602	-0.058 [0.040] N=1,478	-0.057 [0.042] N=1,333	-0.049 [0.046] N=1,133	-0.079 [0.051] N=929
Animal products consumed at least weekly	0.914 P=1,466	0.008 [0.031] N=1,602	0.007 [0.032] N=1,478	-0.005 [0.035] N=1,333	-0.005 [0.038] N=1,133	-0.014 [0.043] N=929
Sweets consumed at least weekly	0.666 P=1,063	-0.024 [0.041] N=1,602	-0.009 [0.044] N=1,478	0.005 [0.046] N=1,333	-0.008 [0.049] N=1,133	0.025 [0.056] N=929
Snack foods consumed at least weekly	0.542 P=861	-0.034 [0.042] N=1,602	-0.030 [0.045] N=1,478	-0.007 [0.049] N=1,333	-0.017 [0.053] N=1,133	-0.010 [0.055] N=929
Packaged soft drinks consumed at least weekly	0.322 P=507	0.004 [0.051] N=1,602	0.009 [0.055] N=1,478	0.012 [0.058] N=1,333	0.019 [0.062] N=1,133	0.009 [0.074] N=929
Grains consumed at least weekly	0.912 P=1,460	0.018 [0.030] N=1,602	0.031 [0.032] N=1,478	0.017 [0.033] N=1,333	0.024 [0.035] N=1,133	0.012 [0.044] N=929
Child was ever breastfed	0.964 P=1,541	0.000 [0.016] N=1,604	0.003 [0.017] N=1,480	0.007 [0.018] N=1,334	0.014 [0.020] N=1,134	0.011 [0.020] N=930

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.010 [0.036] N=1,363	0.045 [0.035] N=1,239	0.039 [0.038] N=1,094	0.046 [0.045] N=894	0.047 [0.064] N=690
Vegetables consumed at least weekly	0.042 [0.045] N=1,363	0.053 [0.047] N=1,239	0.032 [0.051] N=1,094	0.052 [0.058] N=894	-0.006 [0.077] N=690
Meat fish or poultry consumed at least weekly	0.075 [0.056] N=1,363	**0.111* [0.058] N=1,239**	0.098 [0.066] N=1,094	0.029 [0.073] N=894	-0.013 [0.090] N=690
Legumes consumed at least weekly	0.001 [0.061] N=1,363	-0.010 [0.064] N=1,239	0.011 [0.070] N=1,094	0.046 [0.086] N=894	0.020 [0.120] N=690
Animal products consumed at least weekly	0.010 [0.042] N=1,363	0.010 [0.045] N=1,239	-0.016 [0.051] N=1,094	-0.008 [0.059] N=894	-0.044 [0.074] N=690
Sweets consumed at least weekly	-0.031 [0.057] N=1,363	-0.011 [0.062] N=1,239	0.010 [0.070] N=1,094	-0.020 [0.080] N=894	0.093 [0.109] N=690
Snack foods consumed at least weekly	-0.028 [0.056] N=1,363	-0.028 [0.061] N=1,239	0.013 [0.073] N=1,094	-0.005 [0.088] N=894	0.019 [0.101] N=690
Packaged soft drinks consumed at least weekly	-0.012 [0.059] N=1,363	-0.010 [0.065] N=1,239	-0.008 [0.068] N=1,094	0.002 [0.075] N=894	-0.065 [0.094] N=690
Grains consumed at least weekly	0.050 [0.048] N=1,363	0.082 [0.053] N=1,239	0.064 [0.057] N=1,094	0.097 [0.069] N=894	0.117 [0.106] N=690
Child was ever breastfed	0.008 [0.026] N=1,365	0.015 [0.030] N=1,241	0.022 [0.037] N=1,095	0.035 [0.044] N=895	0.057 [0.056] N=691

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.938 P=1,502	-0.008 [0.023] N=1,602	0.010 [0.022] N=1,478	0.004 [0.024] N=1,333	0.001 [0.026] N=1,133	-0.010 [0.031] N=929
Vegetables consumed at least weekly	0.901 P=1,443	0.013 [0.028] N=1,602	0.016 [0.028] N=1,478	0.001 [0.029] N=1,333	0.005 [0.029] N=1,133	-0.023 [0.033] N=929
Meat fish or poultry consumed at least weekly	0.796 P=1,281	**0.076* [0.043] N=1,602**	**0.097** [0.045] N=1,478**	**0.087* [0.049] N=1,333**	0.060 [0.054] N=1,133	0.046 [0.062] N=929
Legumes consumed at least weekly	0.795 P=1,277	-0.048 [0.039] N=1,602	-0.058 [0.040] N=1,478	-0.057 [0.042] N=1,333	-0.050 [0.046] N=1,133	-0.080 [0.051] N=929
Animal products consumed at least weekly	0.914 P=1,466	0.007 [0.031] N=1,602	0.007 [0.032] N=1,478	-0.005 [0.035] N=1,333	-0.006 [0.038] N=1,133	-0.015 [0.042] N=929
Sweets consumed at least weekly	0.666 P=1,063	-0.024 [0.041] N=1,602	-0.009 [0.044] N=1,478	0.005 [0.046] N=1,333	-0.008 [0.049] N=1,133	0.025 [0.056] N=929
Snack foods consumed at least weekly	0.542 P=861	-0.035 [0.043] N=1,602	-0.031 [0.045] N=1,478	-0.006 [0.049] N=1,333	-0.017 [0.053] N=1,133	-0.011 [0.056] N=929
Packaged soft drinks consumed at least weekly	0.322 P=507	0.004 [0.051] N=1,602	0.009 [0.055] N=1,478	0.012 [0.058] N=1,333	0.019 [0.062] N=1,133	0.009 [0.074] N=929
Grains consumed at least weekly	0.912 P=1,460	0.018 [0.030] N=1,602	0.031 [0.032] N=1,478	0.018 [0.033] N=1,333	0.024 [0.035] N=1,133	0.012 [0.044] N=929
Child was ever breastfed	0.964 P=1,541	-0.000 [0.016] N=1,604	0.003 [0.017] N=1,480	0.007 [0.018] N=1,334	0.014 [0.020] N=1,134	0.011 [0.020] N=930

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.010 [0.036] N=1,363	0.046 [0.034] N=1,239	0.041 [0.038] N=1,094	0.047 [0.045] N=894	0.049 [0.064] N=690
Vegetables consumed at least weekly	0.042 [0.045] N=1,363	0.054 [0.047] N=1,239	0.032 [0.051] N=1,094	0.052 [0.058] N=894	-0.005 [0.077] N=690
Meat fish or poultry consumed at least weekly	0.075 [0.056] N=1,363	**0.111* [0.058] N=1,239**	0.099 [0.066] N=1,094	0.030 [0.073] N=894	-0.011 [0.089] N=690
Legumes consumed at least weekly	0.001 [0.061] N=1,363	-0.010 [0.064] N=1,239	0.013 [0.070] N=1,094	0.048 [0.086] N=894	0.022 [0.120] N=690
Animal products consumed at least weekly	0.010 [0.041] N=1,363	0.011 [0.044] N=1,239	-0.012 [0.051] N=1,094	-0.006 [0.059] N=894	-0.042 [0.073] N=690
Sweets consumed at least weekly	-0.031 [0.057] N=1,363	-0.011 [0.062] N=1,239	0.010 [0.070] N=1,094	-0.020 [0.080] N=894	0.093 [0.109] N=690
Snack foods consumed at least weekly	-0.028 [0.056] N=1,363	-0.025 [0.062] N=1,239	0.019 [0.073] N=1,094	-0.003 [0.088] N=894	0.023 [0.101] N=690
Packaged soft drinks consumed at least weekly	-0.012 [0.059] N=1,363	-0.010 [0.065] N=1,239	-0.008 [0.068] N=1,094	0.002 [0.075] N=894	-0.067 [0.094] N=690
Grains consumed at least weekly	0.050 [0.048] N=1,363	0.082 [0.053] N=1,239	0.065 [0.057] N=1,094	0.097 [0.069] N=894	0.116 [0.106] N=690
Child was ever breastfed	0.008 [0.026] N=1,365	0.015 [0.031] N=1,241	0.022 [0.037] N=1,095	0.034 [0.044] N=895	0.056 [0.056] N=691

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.938 P=1,502	-0.017 [0.023] N=1,602	-0.000 [0.022] N=1,478	-0.006 [0.024] N=1,333	-0.008 [0.027] N=1,133	-0.018 [0.033] N=929
Vegetables consumed at least weekly	0.901 P=1,443	0.007 [0.028] N=1,602	0.008 [0.028] N=1,478	-0.006 [0.029] N=1,333	-0.001 [0.029] N=1,133	-0.030 [0.033] N=929
Meat fish or poultry consumed at least weekly	0.796 P=1,281	0.054 [0.042] N=1,602	0.070 [0.043] N=1,478	0.063 [0.048] N=1,333	0.040 [0.052] N=1,133	0.025 [0.060] N=929
Legumes consumed at least weekly	0.795 P=1,277	-0.056 [0.040] N=1,602	**-0.067* [0.040] N=1,478**	-0.064 [0.042] N=1,333	-0.056 [0.046] N=1,133	-0.083 [0.051] N=929
Animal products consumed at least weekly	0.914 P=1,466	0.002 [0.031] N=1,602	0.001 [0.033] N=1,478	-0.010 [0.036] N=1,333	-0.011 [0.039] N=1,133	-0.022 [0.043] N=929
Sweets consumed at least weekly	0.666 P=1,063	-0.030 [0.042] N=1,602	-0.016 [0.044] N=1,478	-0.002 [0.046] N=1,333	-0.014 [0.050] N=1,133	0.018 [0.057] N=929
Snack foods consumed at least weekly	0.542 P=861	-0.034 [0.042] N=1,602	-0.030 [0.045] N=1,478	-0.008 [0.048] N=1,333	-0.018 [0.052] N=1,133	-0.012 [0.055] N=929
Packaged soft drinks consumed at least weekly	0.322 P=507	0.016 [0.050] N=1,602	0.025 [0.054] N=1,478	0.025 [0.056] N=1,333	0.031 [0.060] N=1,133	0.021 [0.070] N=929
Grains consumed at least weekly	0.912 P=1,460	0.012 [0.030] N=1,602	0.025 [0.032] N=1,478	0.012 [0.033] N=1,333	0.017 [0.035] N=1,133	0.007 [0.045] N=929
Child was ever breastfed	0.964 P=1,541	-0.004 [0.016] N=1,604	-0.002 [0.017] N=1,480	0.001 [0.018] N=1,334	0.009 [0.020] N=1,134	0.008 [0.020] N=930

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.002 [0.035] N=1,363	0.035 [0.033] N=1,239	0.034 [0.037] N=1,094	0.051 [0.044] N=894	0.057 [0.065] N=690
Vegetables consumed at least weekly	0.037 [0.045] N=1,363	0.047 [0.046] N=1,239	0.028 [0.051] N=1,094	0.057 [0.057] N=894	0.002 [0.077] N=690
Meat fish or poultry consumed at least weekly	0.054 [0.054] N=1,363	0.080 [0.055] N=1,239	0.075 [0.065] N=1,094	0.016 [0.071] N=894	-0.019 [0.090] N=690
Legumes consumed at least weekly	-0.007 [0.061] N=1,363	-0.022 [0.065] N=1,239	0.004 [0.071] N=1,094	0.048 [0.088] N=894	0.027 [0.123] N=690
Animal products consumed at least weekly	0.005 [0.041] N=1,363	0.005 [0.044] N=1,239	-0.018 [0.052] N=1,094	-0.003 [0.060] N=894	-0.033 [0.073] N=690
Sweets consumed at least weekly	-0.035 [0.057] N=1,363	-0.017 [0.063] N=1,239	0.006 [0.071] N=1,094	-0.018 [0.080] N=894	0.097 [0.111] N=690
Snack foods consumed at least weekly	-0.025 [0.056] N=1,363	-0.023 [0.061] N=1,239	0.023 [0.071] N=1,094	0.003 [0.085] N=894	0.029 [0.099] N=690
Packaged soft drinks consumed at least weekly	0.001 [0.057] N=1,363	0.011 [0.064] N=1,239	0.013 [0.066] N=1,094	0.022 [0.074] N=894	-0.051 [0.091] N=690
Grains consumed at least weekly	0.042 [0.048] N=1,363	0.074 [0.052] N=1,239	0.057 [0.057] N=1,094	0.090 [0.068] N=894	0.107 [0.105] N=690
Child was ever breastfed	0.004 [0.027] N=1,365	0.010 [0.031] N=1,241	0.017 [0.037] N=1,095	0.034 [0.045] N=895	0.052 [0.057] N=691

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.938 P=1,502	-0.018 [0.023] N=1,602	-0.001 [0.022] N=1,478	-0.007 [0.024] N=1,333	-0.009 [0.026] N=1,133	-0.017 [0.032] N=929
Vegetables consumed at least weekly	0.901 P=1,443	0.006 [0.028] N=1,602	0.009 [0.028] N=1,478	-0.005 [0.029] N=1,333	-0.001 [0.029] N=1,133	-0.027 [0.033] N=929
Meat fish or poultry consumed at least weekly	0.796 P=1,281	0.050 [0.042] N=1,602	0.066 [0.043] N=1,478	0.056 [0.047] N=1,333	0.034 [0.052] N=1,133	0.018 [0.059] N=929
Legumes consumed at least weekly	0.795 P=1,277	-0.057 [0.040] N=1,602	**-0.070* [0.041] N=1,478**	-0.070 [0.043] N=1,333	-0.060 [0.047] N=1,133	-0.087 [0.052] N=929
Animal products consumed at least weekly	0.914 P=1,466	0.001 [0.031] N=1,602	0.002 [0.032] N=1,478	-0.009 [0.035] N=1,333	-0.012 [0.038] N=1,133	-0.022 [0.042] N=929
Sweets consumed at least weekly	0.666 P=1,063	-0.027 [0.041] N=1,602	-0.015 [0.044] N=1,478	-0.003 [0.046] N=1,333	-0.013 [0.049] N=1,133	0.019 [0.057] N=929
Snack foods consumed at least weekly	0.542 P=861	-0.032 [0.043] N=1,602	-0.028 [0.045] N=1,478	-0.009 [0.049] N=1,333	-0.016 [0.053] N=1,133	-0.011 [0.056] N=929
Packaged soft drinks consumed at least weekly	0.322 P=507	0.020 [0.050] N=1,602	0.027 [0.054] N=1,478	0.026 [0.057] N=1,333	0.035 [0.060] N=1,133	0.026 [0.069] N=929
Grains consumed at least weekly	0.912 P=1,460	0.013 [0.030] N=1,602	0.026 [0.032] N=1,478	0.014 [0.033] N=1,333	0.019 [0.035] N=1,133	0.010 [0.046] N=929
Child was ever breastfed	0.964 P=1,541	-0.005 [0.017] N=1,604	-0.002 [0.018] N=1,480	0.000 [0.019] N=1,334	0.008 [0.021] N=1,134	0.006 [0.021] N=930

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.001 [0.035] N=1,363	0.035 [0.034] N=1,239	0.031 [0.038] N=1,094	0.045 [0.045] N=894	0.043 [0.066] N=690
Vegetables consumed at least weekly	0.037 [0.044] N=1,363	0.046 [0.046] N=1,239	0.026 [0.051] N=1,094	0.050 [0.057] N=894	-0.017 [0.078] N=690
Meat fish or poultry consumed at least weekly	0.051 [0.054] N=1,363	0.075 [0.056] N=1,239	0.068 [0.065] N=1,094	0.015 [0.072] N=894	-0.029 [0.093] N=690
Legumes consumed at least weekly	-0.004 [0.061] N=1,363	-0.020 [0.064] N=1,239	0.004 [0.070] N=1,094	0.058 [0.085] N=894	0.028 [0.126] N=690
Animal products consumed at least weekly	0.004 [0.041] N=1,363	0.007 [0.044] N=1,239	-0.018 [0.051] N=1,094	-0.014 [0.059] N=894	-0.054 [0.076] N=690
Sweets consumed at least weekly	-0.029 [0.057] N=1,363	-0.013 [0.062] N=1,239	0.007 [0.068] N=1,094	-0.008 [0.080] N=894	0.104 [0.111] N=690
Snack foods consumed at least weekly	-0.019 [0.054] N=1,363	-0.019 [0.059] N=1,239	0.022 [0.069] N=1,094	0.016 [0.083] N=894	0.046 [0.096] N=690
Packaged soft drinks consumed at least weekly	0.008 [0.056] N=1,363	0.013 [0.063] N=1,239	0.014 [0.066] N=1,094	0.028 [0.075] N=894	-0.033 [0.089] N=690
Grains consumed at least weekly	0.042 [0.047] N=1,363	0.073 [0.051] N=1,239	0.056 [0.055] N=1,094	0.088 [0.064] N=894	0.097 [0.099] N=690
Child was ever breastfed	0.005 [0.027] N=1,365	0.011 [0.031] N=1,241	0.017 [0.037] N=1,095	0.036 [0.046] N=895	0.059 [0.058] N=691

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.939 P=1,394	-0.017 [0.023] N=1,487	-0.004 [0.023] N=1,378	-0.010 [0.025] N=1,241	-0.012 [0.027] N=1,059	-0.027 [0.033] N=868
Vegetables consumed at least weekly	0.901 P=1,340	-0.003 [0.029] N=1,487	-0.002 [0.029] N=1,378	-0.021 [0.030] N=1,241	-0.010 [0.030] N=1,059	-0.037 [0.033] N=868
Meat fish or poultry consumed at least weekly	0.795 P=1,186	0.063 [0.041] N=1,487	**0.073* [0.042] N=1,378**	0.055 [0.045] N=1,241	0.026 [0.050] N=1,059	-0.001 [0.057] N=868
Legumes consumed at least weekly	0.790 P=1,179	**-0.067* [0.039] N=1,487**	**-0.082** [0.040] N=1,378**	**-0.085** [0.041] N=1,241**	-0.072 [0.046] N=1,059	**-0.105** [0.050] N=868**
Animal products consumed at least weekly	0.915 P=1,363	0.003 [0.031] N=1,487	0.000 [0.032] N=1,378	-0.013 [0.035] N=1,241	-0.019 [0.037] N=1,059	-0.029 [0.042] N=868
Sweets consumed at least weekly	0.669 P=990	-0.009 [0.043] N=1,487	-0.001 [0.045] N=1,378	0.010 [0.047] N=1,241	-0.001 [0.051] N=1,059	0.033 [0.059] N=868
Snack foods consumed at least weekly	0.544 P=801	-0.036 [0.045] N=1,487	-0.033 [0.048] N=1,378	-0.010 [0.052] N=1,241	-0.021 [0.058] N=1,059	-0.000 [0.063] N=868
Packaged soft drinks consumed at least weekly	0.328 P=480	0.035 [0.051] N=1,487	0.041 [0.055] N=1,378	0.034 [0.058] N=1,241	0.034 [0.061] N=1,059	0.036 [0.072] N=868
Grains consumed at least weekly	0.911 P=1,354	0.010 [0.031] N=1,487	0.020 [0.032] N=1,378	0.002 [0.033] N=1,241	0.012 [0.035] N=1,059	-0.001 [0.043] N=868
Child was ever breastfed	0.964 P=1,431	-0.004 [0.017] N=1,489	0.001 [0.019] N=1,380	0.004 [0.020] N=1,242	0.016 [0.022] N=1,060	0.013 [0.022] N=869

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.007 [0.037] N=1,263	0.033 [0.037] N=1,154	0.025 [0.040] N=1,017	0.033 [0.047] N=835	0.026 [0.066] N=644
Vegetables consumed at least weekly	0.025 [0.046] N=1,263	0.029 [0.049] N=1,154	-0.003 [0.053] N=1,017	0.030 [0.060] N=835	-0.025 [0.076] N=644
Meat fish or poultry consumed at least weekly	0.092 [0.056] N=1,263	**0.109* [0.059] N=1,154**	0.089 [0.066] N=1,017	0.022 [0.075] N=835	-0.026 [0.090] N=644
Legumes consumed at least weekly	-0.010 [0.062] N=1,263	-0.033 [0.064] N=1,154	-0.022 [0.068] N=1,017	0.022 [0.086] N=835	-0.017 [0.118] N=644
Animal products consumed at least weekly	0.009 [0.043] N=1,263	0.005 [0.047] N=1,154	-0.023 [0.053] N=1,017	-0.021 [0.061] N=835	-0.053 [0.077] N=644
Sweets consumed at least weekly	-0.009 [0.059] N=1,263	-0.001 [0.066] N=1,154	0.005 [0.071] N=1,017	-0.024 [0.082] N=835	0.066 [0.113] N=644
Snack foods consumed at least weekly	-0.030 [0.057] N=1,263	-0.031 [0.064] N=1,154	0.006 [0.074] N=1,017	-0.028 [0.088] N=835	-0.008 [0.102] N=644
Packaged soft drinks consumed at least weekly	0.024 [0.061] N=1,263	0.025 [0.066] N=1,154	0.007 [0.068] N=1,017	0.002 [0.075] N=835	-0.047 [0.098] N=644
Grains consumed at least weekly	0.046 [0.049] N=1,263	0.071 [0.054] N=1,154	0.040 [0.057] N=1,017	0.081 [0.068] N=835	0.100 [0.105] N=644
Child was ever breastfed	-0.000 [0.028] N=1,265	0.010 [0.032] N=1,156	0.015 [0.037] N=1,018	0.033 [0.044] N=836	0.051 [0.057] N=645

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.939 P=1,394	-0.019 [0.023] N=1,487	-0.005 [0.023] N=1,378	-0.011 [0.024] N=1,241	-0.013 [0.027] N=1,059	-0.027 [0.031] N=868
Vegetables consumed at least weekly	0.901 P=1,340	-0.004 [0.029] N=1,487	-0.003 [0.029] N=1,378	-0.022 [0.030] N=1,241	-0.010 [0.030] N=1,059	-0.037 [0.032] N=868
Meat fish or poultry consumed at least weekly	0.795 P=1,186	0.065 [0.042] N=1,487	**0.076* [0.043] N=1,378**	0.060 [0.046] N=1,241	0.033 [0.051] N=1,059	0.014 [0.059] N=868
Legumes consumed at least weekly	0.790 P=1,179	**-0.069* [0.039] N=1,487**	**-0.084** [0.040] N=1,378**	**-0.088** [0.041] N=1,241**	**-0.078* [0.046] N=1,059**	**-0.117** [0.051] N=868**
Animal products consumed at least weekly	0.915 P=1,363	0.002 [0.031] N=1,487	-0.001 [0.032] N=1,378	-0.014 [0.035] N=1,241	-0.019 [0.037] N=1,059	-0.029 [0.041] N=868
Sweets consumed at least weekly	0.669 P=990	-0.009 [0.042] N=1,487	-0.002 [0.045] N=1,378	0.012 [0.046] N=1,241	0.001 [0.050] N=1,059	0.034 [0.058] N=868
Snack foods consumed at least weekly	0.544 P=801	-0.037 [0.045] N=1,487	-0.035 [0.048] N=1,378	-0.010 [0.052] N=1,241	-0.023 [0.057] N=1,059	-0.008 [0.061] N=868
Packaged soft drinks consumed at least weekly	0.328 P=480	0.032 [0.052] N=1,487	0.037 [0.055] N=1,378	0.032 [0.059] N=1,241	0.031 [0.062] N=1,059	0.027 [0.073] N=868
Grains consumed at least weekly	0.911 P=1,354	0.009 [0.031] N=1,487	0.019 [0.032] N=1,378	0.000 [0.033] N=1,241	0.009 [0.035] N=1,059	-0.006 [0.043] N=868
Child was ever breastfed	0.964 P=1,431	-0.005 [0.017] N=1,489	0.001 [0.018] N=1,380	0.003 [0.019] N=1,242	0.015 [0.021] N=1,060	0.012 [0.021] N=869

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.004 [0.036] N=1,263	0.032 [0.036] N=1,154	0.027 [0.039] N=1,017	0.034 [0.047] N=835	0.024 [0.064] N=644
Vegetables consumed at least weekly	0.022 [0.046] N=1,263	0.028 [0.048] N=1,154	-0.002 [0.053] N=1,017	0.029 [0.060] N=835	-0.024 [0.074] N=644
Meat fish or poultry consumed at least weekly	0.087 [0.055] N=1,263	**0.105* [0.058] N=1,154**	0.088 [0.065] N=1,017	0.023 [0.075] N=835	-0.017 [0.089] N=644
Legumes consumed at least weekly	-0.011 [0.062] N=1,263	-0.032 [0.065] N=1,154	-0.017 [0.070] N=1,017	0.027 [0.088] N=835	-0.020 [0.120] N=644
Animal products consumed at least weekly	0.007 [0.043] N=1,263	0.003 [0.046] N=1,154	-0.022 [0.052] N=1,017	-0.023 [0.061] N=835	-0.056 [0.076] N=644
Sweets consumed at least weekly	-0.020 [0.059] N=1,263	-0.017 [0.065] N=1,154	-0.003 [0.072] N=1,017	-0.027 [0.083] N=835	0.069 [0.113] N=644
Snack foods consumed at least weekly	-0.040 [0.057] N=1,263	-0.043 [0.063] N=1,154	0.000 [0.074] N=1,017	-0.030 [0.089] N=835	-0.006 [0.102] N=644
Packaged soft drinks consumed at least weekly	0.015 [0.061] N=1,263	0.016 [0.066] N=1,154	0.004 [0.069] N=1,017	-0.000 [0.076] N=835	-0.053 [0.097] N=644
Grains consumed at least weekly	0.045 [0.049] N=1,263	0.072 [0.055] N=1,154	0.046 [0.059] N=1,017	0.084 [0.072] N=835	0.097 [0.107] N=644
Child was ever breastfed	0.001 [0.028] N=1,265	0.010 [0.033] N=1,156	0.015 [0.039] N=1,018	0.036 [0.047] N=836	0.052 [0.061] N=645

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.943 P=1,343	0.001 [0.028] N=1,427	0.002 [0.029] N=1,293	-0.000 [0.033] N=1,132	0.009 [0.033] N=950	0.021 [0.040] N=768
Vegetables consumed at least weekly	0.902 P=1,286	0.028 [0.033] N=1,427	0.042 [0.035] N=1,293	0.041 [0.038] N=1,132	0.045 [0.041] N=950	0.055 [0.047] N=768
Meat fish or poultry consumed at least weekly	0.784 P=1,129	**0.083** [0.040] N=1,427**	**0.087** [0.042] N=1,293**	**0.079* [0.045] N=1,132**	**0.124*** [0.046] N=950**	**0.129** [0.054] N=768**
Legumes consumed at least weekly	0.794 P=1,140	-0.005 [0.046] N=1,427	0.019 [0.049] N=1,293	0.008 [0.052] N=1,132	-0.019 [0.056] N=950	0.008 [0.069] N=768
Animal products consumed at least weekly	0.913 P=1,304	-0.002 [0.036] N=1,427	0.002 [0.038] N=1,293	0.007 [0.040] N=1,132	-0.013 [0.042] N=950	0.014 [0.049] N=768
Sweets consumed at least weekly	0.673 P=962	0.017 [0.051] N=1,427	0.013 [0.053] N=1,293	0.008 [0.057] N=1,132	0.003 [0.059] N=950	-0.049 [0.068] N=768
Snack foods consumed at least weekly	0.546 P=773	-0.019 [0.050] N=1,427	-0.017 [0.053] N=1,293	-0.009 [0.056] N=1,132	-0.005 [0.060] N=950	-0.053 [0.070] N=768
Packaged soft drinks consumed at least weekly	0.314 P=441	-0.015 [0.047] N=1,427	-0.013 [0.049] N=1,293	0.025 [0.051] N=1,132	0.025 [0.056] N=950	0.026 [0.062] N=768
Grains consumed at least weekly	0.911 P=1,300	0.050 [0.041] N=1,427	0.057 [0.043] N=1,293	0.062 [0.046] N=1,132	0.066 [0.051] N=950	0.070 [0.059] N=768
Child was ever breastfed	0.963 P=1,373	-0.005 [0.021] N=1,428	-0.003 [0.021] N=1,294	-0.001 [0.021] N=1,133	-0.007 [0.022] N=951	-0.016 [0.027] N=769

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.041 [0.037] N=1,236	-0.047 [0.039] N=1,102	-0.061 [0.048] N=941	-0.037 [0.048] N=759	-0.026 [0.074] N=577
Vegetables consumed at least weekly	-0.019 [0.038] N=1,236	-0.006 [0.041] N=1,102	-0.019 [0.049] N=941	-0.008 [0.052] N=759	0.007 [0.068] N=577
Meat fish or poultry consumed at least weekly	0.015 [0.060] N=1,236	0.016 [0.063] N=1,102	-0.028 [0.071] N=941	0.045 [0.074] N=759	0.017 [0.099] N=577
Legumes consumed at least weekly	-0.014 [0.054] N=1,236	0.023 [0.059] N=1,102	0.006 [0.066] N=941	-0.029 [0.070] N=759	0.044 [0.085] N=577
Animal products consumed at least weekly	-0.051 [0.044] N=1,236	-0.044 [0.048] N=1,102	-0.044 [0.052] N=941	**-0.091* [0.053] N=759**	-0.070 [0.066] N=577
Sweets consumed at least weekly	0.013 [0.070] N=1,236	0.008 [0.075] N=1,102	-0.001 [0.084] N=941	-0.010 [0.097] N=759	-0.161 [0.124] N=577
Snack foods consumed at least weekly	-0.011 [0.073] N=1,236	-0.008 [0.080] N=1,102	0.013 [0.088] N=941	0.039 [0.102] N=759	-0.078 [0.135] N=577
Packaged soft drinks consumed at least weekly	-0.056 [0.075] N=1,236	-0.054 [0.081] N=1,102	0.006 [0.093] N=941	0.002 [0.112] N=759	-0.031 [0.141] N=577
Grains consumed at least weekly	-0.008 [0.042] N=1,236	-0.008 [0.045] N=1,102	-0.014 [0.051] N=941	-0.032 [0.061] N=759	-0.103 [0.080] N=577
Child was ever breastfed	0.022 [0.040] N=1,237	0.030 [0.041] N=1,103	0.040 [0.043] N=942	0.033 [0.048] N=760	0.023 [0.066] N=578

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.943 P=1,343	0.001 [0.028] N=1,427	0.003 [0.029] N=1,293	-0.000 [0.033] N=1,132	0.009 [0.033] N=950	0.021 [0.040] N=768
Vegetables consumed at least weekly	0.902 P=1,286	0.029 [0.033] N=1,427	0.043 [0.035] N=1,293	0.042 [0.038] N=1,132	0.045 [0.041] N=950	0.055 [0.047] N=768
Meat fish or poultry consumed at least weekly	0.784 P=1,129	**0.084** [0.040] N=1,427**	**0.089** [0.041] N=1,293**	**0.080* [0.045] N=1,132**	**0.124*** [0.046] N=950**	**0.129** [0.054] N=768**
Legumes consumed at least weekly	0.794 P=1,140	-0.005 [0.046] N=1,427	0.019 [0.049] N=1,293	0.008 [0.052] N=1,132	-0.019 [0.056] N=950	0.008 [0.069] N=768
Animal products consumed at least weekly	0.913 P=1,304	0.001 [0.035] N=1,427	0.005 [0.037] N=1,293	0.009 [0.040] N=1,132	-0.012 [0.042] N=950	0.014 [0.048] N=768
Sweets consumed at least weekly	0.673 P=962	0.019 [0.051] N=1,427	0.015 [0.053] N=1,293	0.010 [0.057] N=1,132	0.003 [0.059] N=950	-0.049 [0.067] N=768
Snack foods consumed at least weekly	0.546 P=773	-0.016 [0.049] N=1,427	-0.015 [0.053] N=1,293	-0.007 [0.056] N=1,132	-0.004 [0.060] N=950	-0.054 [0.069] N=768
Packaged soft drinks consumed at least weekly	0.314 P=441	-0.014 [0.047] N=1,427	-0.012 [0.048] N=1,293	0.027 [0.050] N=1,132	0.025 [0.056] N=950	0.025 [0.061] N=768
Grains consumed at least weekly	0.911 P=1,300	0.050 [0.041] N=1,427	0.057 [0.043] N=1,293	0.062 [0.046] N=1,132	0.066 [0.052] N=950	0.070 [0.059] N=768
Child was ever breastfed	0.963 P=1,373	-0.004 [0.021] N=1,428	-0.002 [0.021] N=1,294	-0.001 [0.021] N=1,133	-0.007 [0.022] N=951	-0.016 [0.027] N=769

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.040 [0.037] N=1,236	-0.045 [0.039] N=1,102	-0.061 [0.048] N=941	-0.038 [0.049] N=759	-0.027 [0.075] N=577
Vegetables consumed at least weekly	-0.018 [0.038] N=1,236	-0.004 [0.041] N=1,102	-0.018 [0.049] N=941	-0.006 [0.051] N=759	0.010 [0.068] N=577
Meat fish or poultry consumed at least weekly	0.018 [0.060] N=1,236	0.021 [0.062] N=1,102	-0.023 [0.071] N=941	0.050 [0.074] N=759	0.026 [0.100] N=577
Legumes consumed at least weekly	-0.013 [0.054] N=1,236	0.024 [0.059] N=1,102	0.007 [0.066] N=941	-0.027 [0.069] N=759	0.040 [0.084] N=577
Animal products consumed at least weekly	-0.046 [0.042] N=1,236	-0.037 [0.047] N=1,102	-0.038 [0.051] N=941	-0.085 [0.053] N=759	-0.060 [0.064] N=577
Sweets consumed at least weekly	0.017 [0.070] N=1,236	0.013 [0.075] N=1,102	0.003 [0.084] N=941	-0.009 [0.097] N=759	-0.158 [0.123] N=577
Snack foods consumed at least weekly	-0.004 [0.073] N=1,236	-0.000 [0.080] N=1,102	0.020 [0.088] N=941	0.042 [0.103] N=759	-0.070 [0.136] N=577
Packaged soft drinks consumed at least weekly	-0.053 [0.074] N=1,236	-0.052 [0.080] N=1,102	0.010 [0.092] N=941	0.004 [0.112] N=759	-0.021 [0.140] N=577
Grains consumed at least weekly	-0.008 [0.043] N=1,236	-0.008 [0.045] N=1,102	-0.016 [0.051] N=941	-0.035 [0.061] N=759	-0.110 [0.081] N=577
Child was ever breastfed	0.023 [0.040] N=1,237	0.032 [0.041] N=1,103	0.041 [0.043] N=942	0.033 [0.048] N=760	0.024 [0.066] N=578

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.943 P=1,343	-0.001 [0.028] N=1,427	0.001 [0.029] N=1,293	0.000 [0.033] N=1,132	0.008 [0.033] N=950	0.020 [0.040] N=768
Vegetables consumed at least weekly	0.902 P=1,286	0.027 [0.033] N=1,427	0.041 [0.034] N=1,293	0.042 [0.037] N=1,132	0.046 [0.041] N=950	0.055 [0.047] N=768
Meat fish or poultry consumed at least weekly	0.784 P=1,129	**0.076* [0.039] N=1,427**	**0.083** [0.040] N=1,293**	**0.081* [0.043] N=1,132**	**0.124*** [0.043] N=950**	**0.128** [0.051] N=768**
Legumes consumed at least weekly	0.794 P=1,140	-0.007 [0.046] N=1,427	0.018 [0.049] N=1,293	0.009 [0.053] N=1,132	-0.019 [0.056] N=950	0.009 [0.069] N=768
Animal products consumed at least weekly	0.913 P=1,304	-0.001 [0.035] N=1,427	0.004 [0.037] N=1,293	0.010 [0.040] N=1,132	-0.013 [0.042] N=950	0.014 [0.048] N=768
Sweets consumed at least weekly	0.673 P=962	0.019 [0.051] N=1,427	0.014 [0.053] N=1,293	0.010 [0.057] N=1,132	0.002 [0.059] N=950	-0.050 [0.067] N=768
Snack foods consumed at least weekly	0.546 P=773	-0.013 [0.050] N=1,427	-0.013 [0.053] N=1,293	-0.007 [0.056] N=1,132	-0.007 [0.060] N=950	-0.055 [0.070] N=768
Packaged soft drinks consumed at least weekly	0.314 P=441	-0.008 [0.048] N=1,427	-0.009 [0.050] N=1,293	0.026 [0.052] N=1,132	0.022 [0.058] N=950	0.025 [0.064] N=768
Grains consumed at least weekly	0.911 P=1,300	0.047 [0.041] N=1,427	0.055 [0.043] N=1,293	0.062 [0.046] N=1,132	0.066 [0.051] N=950	0.069 [0.059] N=768
Child was ever breastfed	0.963 P=1,373	-0.005 [0.021] N=1,428	-0.003 [0.021] N=1,294	-0.001 [0.021] N=1,133	-0.007 [0.022] N=951	-0.017 [0.027] N=769

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.041 [0.036] N=1,236	-0.045 [0.038] N=1,102	-0.056 [0.047] N=941	-0.037 [0.049] N=759	-0.025 [0.076] N=577
Vegetables consumed at least weekly	-0.019 [0.038] N=1,236	-0.004 [0.040] N=1,102	-0.014 [0.047] N=941	-0.006 [0.050] N=759	0.012 [0.067] N=577
Meat fish or poultry consumed at least weekly	0.013 [0.057] N=1,236	0.019 [0.059] N=1,102	-0.013 [0.066] N=941	0.057 [0.071] N=759	0.033 [0.091] N=577
Legumes consumed at least weekly	-0.011 [0.053] N=1,236	0.026 [0.059] N=1,102	0.008 [0.067] N=941	-0.026 [0.070] N=759	0.037 [0.088] N=577
Animal products consumed at least weekly	-0.047 [0.042] N=1,236	-0.039 [0.047] N=1,102	-0.037 [0.051] N=941	**-0.091* [0.053] N=759**	-0.064 [0.063] N=577
Sweets consumed at least weekly	0.017 [0.070] N=1,236	0.011 [0.075] N=1,102	0.002 [0.083] N=941	-0.018 [0.098] N=759	-0.167 [0.123] N=577
Snack foods consumed at least weekly	0.003 [0.073] N=1,236	0.003 [0.081] N=1,102	0.016 [0.089] N=941	0.031 [0.103] N=759	-0.077 [0.139] N=577
Packaged soft drinks consumed at least weekly	-0.053 [0.075] N=1,236	-0.056 [0.081] N=1,102	-0.006 [0.094] N=941	-0.022 [0.114] N=759	-0.037 [0.141] N=577
Grains consumed at least weekly	-0.011 [0.042] N=1,236	-0.010 [0.044] N=1,102	-0.014 [0.050] N=941	-0.038 [0.060] N=759	-0.113 [0.080] N=577
Child was ever breastfed	0.022 [0.040] N=1,237	0.030 [0.041] N=1,103	0.042 [0.043] N=942	0.037 [0.048] N=760	0.026 [0.065] N=578

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.943 P=1,343	-0.001 [0.027] N=1,427	0.000 [0.029] N=1,293	-0.001 [0.032] N=1,132	0.002 [0.033] N=950	0.012 [0.040] N=768
Vegetables consumed at least weekly	0.902 P=1,286	0.027 [0.034] N=1,427	0.041 [0.035] N=1,293	0.041 [0.039] N=1,132	0.042 [0.042] N=950	0.051 [0.048] N=768
Meat fish or poultry consumed at least weekly	0.784 P=1,129	**0.078* [0.040] N=1,427**	**0.084** [0.041] N=1,293**	**0.078* [0.044] N=1,132**	**0.113** [0.046] N=950**	**0.110** [0.053] N=768**
Legumes consumed at least weekly	0.794 P=1,140	-0.001 [0.046] N=1,427	0.023 [0.049] N=1,293	0.013 [0.052] N=1,132	-0.019 [0.056] N=950	0.009 [0.069] N=768
Animal products consumed at least weekly	0.913 P=1,304	0.004 [0.035] N=1,427	0.010 [0.037] N=1,293	0.017 [0.040] N=1,132	-0.009 [0.042] N=950	0.015 [0.048] N=768
Sweets consumed at least weekly	0.673 P=962	0.027 [0.052] N=1,427	0.022 [0.054] N=1,293	0.015 [0.058] N=1,132	0.004 [0.061] N=950	-0.047 [0.070] N=768
Snack foods consumed at least weekly	0.546 P=773	-0.003 [0.050] N=1,427	-0.004 [0.053] N=1,293	0.001 [0.056] N=1,132	-0.001 [0.060] N=950	-0.047 [0.069] N=768
Packaged soft drinks consumed at least weekly	0.314 P=441	0.004 [0.048] N=1,427	0.001 [0.050] N=1,293	0.036 [0.052] N=1,132	0.034 [0.058] N=950	0.040 [0.064] N=768
Grains consumed at least weekly	0.911 P=1,300	0.047 [0.041] N=1,427	0.054 [0.043] N=1,293	0.059 [0.046] N=1,132	0.063 [0.051] N=950	0.066 [0.059] N=768
Child was ever breastfed	0.963 P=1,373	-0.005 [0.021] N=1,428	-0.003 [0.021] N=1,294	-0.001 [0.021] N=1,133	-0.008 [0.022] N=951	-0.019 [0.028] N=769

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.043 [0.036] N=1,236	-0.048 [0.038] N=1,102	-0.059 [0.047] N=941	-0.046 [0.050] N=759	-0.047 [0.075] N=577
Vegetables consumed at least weekly	-0.020 [0.038] N=1,236	-0.004 [0.041] N=1,102	-0.016 [0.049] N=941	-0.007 [0.051] N=759	0.006 [0.068] N=577
Meat fish or poultry consumed at least weekly	0.015 [0.056] N=1,236	0.019 [0.058] N=1,102	-0.019 [0.064] N=941	0.037 [0.072] N=759	-0.013 [0.090] N=577
Legumes consumed at least weekly	-0.006 [0.053] N=1,236	0.030 [0.059] N=1,102	0.016 [0.066] N=941	-0.023 [0.070] N=759	0.047 [0.084] N=577
Animal products consumed at least weekly	-0.043 [0.040] N=1,236	-0.032 [0.044] N=1,102	-0.029 [0.048] N=941	-0.084 [0.051] N=759	-0.062 [0.060] N=577
Sweets consumed at least weekly	0.026 [0.071] N=1,236	0.021 [0.076] N=1,102	0.008 [0.084] N=941	-0.013 [0.099] N=759	-0.160 [0.121] N=577
Snack foods consumed at least weekly	0.010 [0.073] N=1,236	0.009 [0.081] N=1,102	0.025 [0.088] N=941	0.039 [0.102] N=759	-0.064 [0.134] N=577
Packaged soft drinks consumed at least weekly	-0.035 [0.075] N=1,236	-0.040 [0.081] N=1,102	0.014 [0.093] N=941	0.010 [0.113] N=759	0.014 [0.142] N=577
Grains consumed at least weekly	-0.013 [0.042] N=1,236	-0.015 [0.045] N=1,102	-0.021 [0.050] N=941	-0.046 [0.061] N=759	-0.127 [0.080] N=577
Child was ever breastfed	0.023 [0.039] N=1,237	0.033 [0.040] N=1,103	0.043 [0.042] N=942	0.033 [0.048] N=760	0.022 [0.067] N=578

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.944 P=1,247	0.006 [0.030] N=1,324	0.007 [0.031] N=1,198	0.013 [0.035] N=1,054	0.012 [0.035] N=885	0.022 [0.042] N=718
Vegetables consumed at least weekly	0.904 P=1,196	0.030 [0.034] N=1,324	0.043 [0.036] N=1,198	0.055 [0.039] N=1,054	0.049 [0.042] N=885	0.042 [0.049] N=718
Meat fish or poultry consumed at least weekly	0.783 P=1,046	**0.082* [0.043] N=1,324**	**0.083* [0.044] N=1,198**	**0.082* [0.048] N=1,054**	**0.100** [0.050] N=885**	**0.103* [0.057] N=718**
Legumes consumed at least weekly	0.792 P=1,056	0.001 [0.048] N=1,324	0.028 [0.052] N=1,198	0.019 [0.055] N=1,054	-0.021 [0.060] N=885	0.004 [0.074] N=718
Animal products consumed at least weekly	0.914 P=1,213	0.005 [0.037] N=1,324	0.014 [0.039] N=1,198	0.024 [0.042] N=1,054	-0.005 [0.043] N=885	0.019 [0.049] N=718
Sweets consumed at least weekly	0.684 P=905	0.023 [0.052] N=1,324	0.021 [0.054] N=1,198	0.009 [0.059] N=1,054	-0.006 [0.063] N=885	-0.052 [0.074] N=718
Snack foods consumed at least weekly	0.553 P=726	-0.007 [0.050] N=1,324	-0.009 [0.054] N=1,198	-0.011 [0.056] N=1,054	-0.009 [0.061] N=885	-0.042 [0.071] N=718
Packaged soft drinks consumed at least weekly	0.324 P=424	-0.018 [0.048] N=1,324	-0.022 [0.050] N=1,198	0.016 [0.052] N=1,054	0.017 [0.058] N=885	0.017 [0.065] N=718
Grains consumed at least weekly	0.911 P=1,206	0.060 [0.043] N=1,324	0.067 [0.045] N=1,198	0.075 [0.048] N=1,054	0.075 [0.053] N=885	0.073 [0.060] N=718
Child was ever breastfed	0.965 P=1,278	-0.010 [0.019] N=1,325	-0.011 [0.020] N=1,199	-0.010 [0.019] N=1,055	-0.021 [0.020] N=886	-0.027 [0.027] N=719

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.029 [0.039] N=1,143	-0.032 [0.042] N=1,017	-0.027 [0.052] N=873	-0.022 [0.053] N=704	-0.017 [0.076] N=537
Vegetables consumed at least weekly	-0.010 [0.040] N=1,143	0.005 [0.043] N=1,017	0.018 [0.050] N=873	0.017 [0.055] N=704	-0.005 [0.068] N=537
Meat fish or poultry consumed at least weekly	0.038 [0.068] N=1,143	0.042 [0.071] N=1,017	0.016 [0.080] N=873	0.040 [0.087] N=704	0.008 [0.114] N=537
Legumes consumed at least weekly	0.011 [0.056] N=1,143	0.053 [0.061] N=1,017	0.048 [0.069] N=873	-0.002 [0.074] N=704	0.067 [0.091] N=537
Animal products consumed at least weekly	-0.038 [0.047] N=1,143	-0.020 [0.052] N=1,017	-0.009 [0.055] N=873	-0.067 [0.057] N=704	-0.041 [0.067] N=537
Sweets consumed at least weekly	0.024 [0.073] N=1,143	0.021 [0.079] N=1,017	-0.007 [0.088] N=873	-0.040 [0.099] N=704	-0.205 [0.127] N=537
Snack foods consumed at least weekly	0.013 [0.075] N=1,143	0.012 [0.083] N=1,017	0.010 [0.092] N=873	0.027 [0.105] N=704	-0.050 [0.141] N=537
Packaged soft drinks consumed at least weekly	-0.067 [0.075] N=1,143	-0.075 [0.081] N=1,017	-0.021 [0.093] N=873	-0.028 [0.114] N=704	-0.071 [0.145] N=537
Grains consumed at least weekly	0.013 [0.046] N=1,143	0.015 [0.049] N=1,017	0.019 [0.056] N=873	-0.008 [0.067] N=704	-0.097 [0.087] N=537
Child was ever breastfed	0.017 [0.037] N=1,144	0.020 [0.038] N=1,018	0.028 [0.038] N=874	0.010 [0.041] N=705	0.001 [0.061] N=538

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	0.944 P=1,247	0.006 [0.029] N=1,324	0.007 [0.031] N=1,198	0.013 [0.035] N=1,054	0.011 [0.034] N=885	0.021 [0.041] N=718
Vegetables consumed at least weekly	0.904 P=1,196	0.031 [0.034] N=1,324	0.043 [0.036] N=1,198	0.055 [0.039] N=1,054	0.047 [0.042] N=885	0.039 [0.049] N=718
Meat fish or poultry consumed at least weekly	0.783 P=1,046	**0.085** [0.043] N=1,324**	**0.088** [0.044] N=1,198**	**0.086* [0.047] N=1,054**	**0.104** [0.049] N=885**	**0.106* [0.056] N=718**
Legumes consumed at least weekly	0.792 P=1,056	-0.000 [0.048] N=1,324	0.026 [0.052] N=1,198	0.018 [0.055] N=1,054	-0.026 [0.060] N=885	0.000 [0.074] N=718
Animal products consumed at least weekly	0.914 P=1,213	0.005 [0.037] N=1,324	0.013 [0.039] N=1,198	0.023 [0.042] N=1,054	-0.006 [0.043] N=885	0.017 [0.049] N=718
Sweets consumed at least weekly	0.684 P=905	0.023 [0.052] N=1,324	0.021 [0.055] N=1,198	0.009 [0.060] N=1,054	-0.006 [0.064] N=885	-0.054 [0.074] N=718
Snack foods consumed at least weekly	0.553 P=726	-0.005 [0.049] N=1,324	-0.007 [0.053] N=1,198	-0.008 [0.056] N=1,054	-0.008 [0.060] N=885	-0.041 [0.071] N=718
Packaged soft drinks consumed at least weekly	0.324 P=424	-0.018 [0.048] N=1,324	-0.022 [0.050] N=1,198	0.017 [0.051] N=1,054	0.016 [0.058] N=885	0.014 [0.065] N=718
Grains consumed at least weekly	0.911 P=1,206	0.059 [0.043] N=1,324	0.066 [0.045] N=1,198	0.073 [0.048] N=1,054	0.073 [0.053] N=885	0.071 [0.060] N=718
Child was ever breastfed	0.965 P=1,278	-0.009 [0.020] N=1,325	-0.010 [0.020] N=1,199	-0.006 [0.020] N=1,055	-0.015 [0.021] N=886	-0.021 [0.027] N=719

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
Fruit consumed at least weekly	-0.028 [0.040] N=1,143	-0.030 [0.043] N=1,017	-0.026 [0.053] N=873	-0.023 [0.054] N=704	-0.015 [0.081] N=537
Vegetables consumed at least weekly	-0.010 [0.040] N=1,143	0.004 [0.043] N=1,017	0.018 [0.050] N=873	0.012 [0.055] N=704	-0.004 [0.071] N=537
Meat fish or poultry consumed at least weekly	0.041 [0.067] N=1,143	0.046 [0.070] N=1,017	0.019 [0.080] N=873	0.043 [0.087] N=704	0.009 [0.115] N=537
Legumes consumed at least weekly	0.010 [0.056] N=1,143	0.054 [0.061] N=1,017	0.047 [0.070] N=873	-0.010 [0.075] N=704	0.066 [0.090] N=537
Animal products consumed at least weekly	-0.039 [0.047] N=1,143	-0.021 [0.052] N=1,017	-0.011 [0.055] N=873	-0.072 [0.058] N=704	-0.050 [0.068] N=537
Sweets consumed at least weekly	0.027 [0.073] N=1,143	0.025 [0.080] N=1,017	0.002 [0.089] N=873	-0.027 [0.101] N=704	-0.182 [0.131] N=537
Snack foods consumed at least weekly	0.014 [0.075] N=1,143	0.013 [0.083] N=1,017	0.015 [0.092] N=873	0.037 [0.105] N=704	-0.044 [0.141] N=537
Packaged soft drinks consumed at least weekly	-0.066 [0.075] N=1,143	-0.074 [0.081] N=1,017	-0.015 [0.093] N=873	-0.020 [0.112] N=704	-0.058 [0.144] N=537
Grains consumed at least weekly	0.012 [0.045] N=1,143	0.014 [0.048] N=1,017	0.017 [0.055] N=873	-0.012 [0.066] N=704	-0.102 [0.086] N=537
Child was ever breastfed	0.018 [0.038] N=1,144	0.020 [0.039] N=1,018	0.032 [0.040] N=874	0.020 [0.045] N=705	0.007 [0.065] N=538

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.