Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for Ages 1-3: Youngest Child

Age reference: 16 December 2022. Any-child household eligibility; one youngest child within the age interval is retained before outcome exclusions. WHO 2006 standards apply throughout these cohorts. WHO outcomes defined only at ages 5 and above are outside this analysis. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.560	0.275 [0.264] N=1,392	0.231 [0.287] N=1,290	0.230 [0.314] N=1,165	0.382 [0.327] N=992	0.205 [0.360] N=818
WHO weight-for-age z score	-0.050	-0.056 [0.142] N=1,594	-0.031 [0.155] N=1,471	-0.011 [0.161] N=1,326	-0.016 [0.176] N=1,128	-0.198 [0.196] N=925
WHO weight-for-length/height z score, 0-60 months	0.365	0.053 [0.217] N=1,368	0.014 [0.232] N=1,271	-0.066 [0.261] N=1,149	-0.141 [0.288] N=979	-0.089 [0.319] N=804
WHO BMI-for-age z score	0.383	0.197 [0.254] N=1,337	0.239 [0.273] N=1,243	0.144 [0.308] N=1,122	-0.058 [0.340] N=956	-0.010 [0.383] N=788
Stunted: HAZ below -2	0.271 P=368	**-0.077* [0.044] N=1,392**	**-0.082* [0.047] N=1,290**	-0.071 [0.051] N=1,165	-0.078 [0.054] N=992	-0.039 [0.059] N=818
Underweight: WAZ below -2	0.075 P=121	**0.038* [0.023] N=1,594**	0.035 [0.025] N=1,471	0.034 [0.026] N=1,326	0.042 [0.030] N=1,128	**0.071** [0.036] N=925**
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=151	-0.029 [0.035] N=1,368	-0.028 [0.036] N=1,271	-0.028 [0.038] N=1,149	-0.016 [0.042] N=979	-0.033 [0.044] N=804
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=262	0.024 [0.047] N=1,368	0.018 [0.051] N=1,271	0.001 [0.056] N=1,149	0.002 [0.062] N=979	-0.004 [0.069] N=804
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=121	-0.005 [0.028] N=1,368	-0.013 [0.030] N=1,271	-0.004 [0.032] N=1,149	-0.005 [0.033] N=979	-0.001 [0.037] N=804
CIAF: any stunting, underweight, or wasting, 0-60 months	0.353 P=463	**-0.104** [0.045] N=1,325**	**-0.111** [0.047] N=1,232**	**-0.108** [0.050] N=1,114**	**-0.108** [0.053] N=949**	-0.079 [0.058] N=781
BMI-for-age z score below -2	0.118 P=160	-0.028 [0.039] N=1,337	-0.033 [0.040] N=1,243	-0.024 [0.043] N=1,122	0.013 [0.047] N=956	0.005 [0.051] N=788

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.173 [0.360] N=1,176	0.049 [0.413] N=1,074	0.065 [0.494] N=949	0.435 [0.531] N=776	-0.007 [0.694] N=602
WHO weight-for-age z score	-0.180 [0.184] N=1,356	-0.174 [0.208] N=1,233	-0.151 [0.220] N=1,088	-0.187 [0.276] N=890	**-0.799** [0.329] N=687**
WHO weight-for-length/height z score, 0-60 months	0.011 [0.263] N=1,155	-0.062 [0.292] N=1,058	-0.265 [0.360] N=936	-0.517 [0.425] N=766	-0.634 [0.492] N=591
WHO BMI-for-age z score	0.310 [0.289] N=1,131	0.408 [0.319] N=1,037	0.219 [0.396] N=916	-0.233 [0.452] N=750	-0.268 [0.560] N=582
Stunted: HAZ below -2	-0.087 [0.061] N=1,176	-0.095 [0.068] N=1,074	-0.087 [0.078] N=949	-0.120 [0.084] N=776	-0.051 [0.115] N=602
Underweight: WAZ below -2	0.025 [0.031] N=1,356	0.021 [0.035] N=1,233	0.019 [0.039] N=1,088	0.033 [0.048] N=890	**0.123* [0.063] N=687**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.017 [0.051] N=1,155	-0.012 [0.054] N=1,058	-0.008 [0.062] N=936	0.027 [0.078] N=766	-0.008 [0.097] N=591
Overweight: WLZ/WHZ above +2, 0-60 months	0.011 [0.052] N=1,155	-0.001 [0.057] N=1,058	-0.042 [0.066] N=936	-0.055 [0.079] N=766	-0.127 [0.088] N=591
Obese: WLZ/WHZ above +3, 0-60 months	-0.020 [0.038] N=1,155	-0.036 [0.043] N=1,058	-0.020 [0.049] N=936	-0.022 [0.055] N=766	-0.032 [0.068] N=591
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.087 [0.070] N=1,119	-0.095 [0.076] N=1,026	-0.091 [0.085] N=908	-0.094 [0.098] N=743	-0.031 [0.136] N=575
BMI-for-age z score below -2	-0.048 [0.051] N=1,131	-0.061 [0.053] N=1,037	-0.050 [0.062] N=916	0.019 [0.078] N=750	-0.017 [0.096] N=582

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.560	0.275 [0.264] N=1,392	0.231 [0.287] N=1,290	0.230 [0.314] N=1,165	0.384 [0.326] N=992	0.209 [0.357] N=818
WHO weight-for-age z score	-0.050	-0.060 [0.141] N=1,594	-0.035 [0.154] N=1,471	-0.011 [0.161] N=1,326	-0.018 [0.176] N=1,128	-0.205 [0.194] N=925
WHO weight-for-length/height z score, 0-60 months	0.365	0.051 [0.216] N=1,368	0.010 [0.232] N=1,271	-0.066 [0.261] N=1,149	-0.146 [0.288] N=979	-0.100 [0.317] N=804
WHO BMI-for-age z score	0.383	0.195 [0.254] N=1,337	0.236 [0.273] N=1,243	0.145 [0.308] N=1,122	-0.061 [0.340] N=956	-0.022 [0.382] N=788
Stunted: HAZ below -2	0.271 P=368	**-0.076* [0.044] N=1,392**	**-0.082* [0.047] N=1,290**	-0.072 [0.050] N=1,165	-0.078 [0.054] N=992	-0.038 [0.059] N=818
Underweight: WAZ below -2	0.075 P=121	**0.039* [0.023] N=1,594**	0.035 [0.025] N=1,471	0.034 [0.026] N=1,326	0.042 [0.030] N=1,128	**0.072** [0.036] N=925**
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=151	-0.028 [0.036] N=1,368	-0.026 [0.036] N=1,271	-0.028 [0.038] N=1,149	-0.014 [0.043] N=979	-0.029 [0.044] N=804
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=262	0.023 [0.047] N=1,368	0.018 [0.051] N=1,271	0.001 [0.056] N=1,149	0.001 [0.062] N=979	-0.004 [0.069] N=804
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=121	-0.005 [0.028] N=1,368	-0.012 [0.030] N=1,271	-0.004 [0.032] N=1,149	-0.005 [0.033] N=979	-0.000 [0.037] N=804
CIAF: any stunting, underweight, or wasting, 0-60 months	0.353 P=463	**-0.102** [0.045] N=1,325**	**-0.108** [0.047] N=1,232**	**-0.108** [0.050] N=1,114**	**-0.105** [0.052] N=949**	-0.071 [0.057] N=781
BMI-for-age z score below -2	0.118 P=160	-0.028 [0.039] N=1,337	-0.032 [0.040] N=1,243	-0.024 [0.043] N=1,122	0.014 [0.047] N=956	0.009 [0.051] N=788

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.170 [0.359] N=1,176	0.046 [0.412] N=1,074	0.058 [0.492] N=949	0.421 [0.529] N=776	-0.022 [0.691] N=602
WHO weight-for-age z score	-0.179 [0.182] N=1,356	-0.169 [0.207] N=1,233	-0.138 [0.219] N=1,088	-0.183 [0.275] N=890	**-0.784** [0.323] N=687**
WHO weight-for-length/height z score, 0-60 months	0.010 [0.262] N=1,155	-0.064 [0.291] N=1,058	-0.257 [0.358] N=936	-0.520 [0.424] N=766	-0.640 [0.495] N=591
WHO BMI-for-age z score	0.311 [0.289] N=1,131	0.409 [0.318] N=1,037	0.232 [0.394] N=916	-0.231 [0.452] N=750	-0.279 [0.564] N=582
Stunted: HAZ below -2	-0.088 [0.061] N=1,176	-0.097 [0.068] N=1,074	-0.089 [0.077] N=949	-0.121 [0.084] N=776	-0.051 [0.115] N=602
Underweight: WAZ below -2	0.025 [0.030] N=1,356	0.020 [0.035] N=1,233	0.017 [0.038] N=1,088	0.032 [0.048] N=890	**0.120* [0.062] N=687**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.017 [0.051] N=1,155	-0.012 [0.054] N=1,058	-0.010 [0.062] N=936	0.028 [0.079] N=766	-0.006 [0.099] N=591
Overweight: WLZ/WHZ above +2, 0-60 months	0.011 [0.052] N=1,155	-0.001 [0.057] N=1,058	-0.042 [0.066] N=936	-0.055 [0.079] N=766	-0.127 [0.089] N=591
Obese: WLZ/WHZ above +3, 0-60 months	-0.020 [0.038] N=1,155	-0.035 [0.044] N=1,058	-0.021 [0.049] N=936	-0.022 [0.055] N=766	-0.032 [0.069] N=591
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.088 [0.070] N=1,119	-0.097 [0.077] N=1,026	-0.098 [0.086] N=908	-0.097 [0.100] N=743	-0.030 [0.138] N=575
BMI-for-age z score below -2	-0.048 [0.052] N=1,131	-0.061 [0.054] N=1,037	-0.053 [0.062] N=916	0.019 [0.079] N=750	-0.015 [0.098] N=582

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.560	0.247 [0.265] N=1,392	0.189 [0.289] N=1,290	0.198 [0.316] N=1,165	0.363 [0.323] N=992	0.168 [0.352] N=818
WHO weight-for-age z score	-0.050	-0.102 [0.135] N=1,594	-0.079 [0.149] N=1,471	-0.052 [0.155] N=1,326	-0.048 [0.169] N=1,128	-0.240 [0.184] N=925
WHO weight-for-length/height z score, 0-60 months	0.365	0.032 [0.220] N=1,368	-0.006 [0.236] N=1,271	-0.071 [0.262] N=1,149	-0.145 [0.286] N=979	-0.112 [0.316] N=804
WHO BMI-for-age z score	0.383	0.182 [0.257] N=1,337	0.222 [0.277] N=1,243	0.138 [0.310] N=1,122	-0.061 [0.339] N=956	-0.032 [0.383] N=788
Stunted: HAZ below -2	0.271 P=368	-0.067 [0.043] N=1,392	-0.067 [0.046] N=1,290	-0.062 [0.049] N=1,165	-0.073 [0.052] N=992	-0.028 [0.056] N=818
Underweight: WAZ below -2	0.075 P=121	**0.047** [0.022] N=1,594**	**0.045* [0.023] N=1,471**	**0.044* [0.025] N=1,326**	**0.050* [0.028] N=1,128**	**0.082** [0.033] N=925**
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=151	-0.022 [0.035] N=1,368	-0.018 [0.036] N=1,271	-0.021 [0.038] N=1,149	-0.011 [0.043] N=979	-0.025 [0.044] N=804
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=262	0.027 [0.047] N=1,368	0.023 [0.051] N=1,271	0.006 [0.056] N=1,149	0.006 [0.062] N=979	-0.001 [0.069] N=804
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=121	-0.004 [0.029] N=1,368	-0.010 [0.031] N=1,271	-0.002 [0.032] N=1,149	-0.003 [0.033] N=979	-0.001 [0.037] N=804
CIAF: any stunting, underweight, or wasting, 0-60 months	0.353 P=463	**-0.083* [0.044] N=1,325**	**-0.082* [0.046] N=1,232**	**-0.089* [0.049] N=1,114**	**-0.093* [0.051] N=949**	-0.052 [0.056] N=781
BMI-for-age z score below -2	0.118 P=160	-0.021 [0.039] N=1,337	-0.022 [0.041] N=1,243	-0.016 [0.044] N=1,122	0.019 [0.047] N=956	0.012 [0.051] N=788

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.131 [0.362] N=1,176	-0.028 [0.417] N=1,074	-0.008 [0.495] N=949	0.387 [0.524] N=776	-0.090 [0.675] N=602
WHO weight-for-age z score	-0.219 [0.178] N=1,356	-0.215 [0.203] N=1,233	-0.175 [0.218] N=1,088	-0.194 [0.276] N=890	**-0.790** [0.321] N=687**
WHO weight-for-length/height z score, 0-60 months	0.005 [0.267] N=1,155	-0.061 [0.299] N=1,058	-0.234 [0.362] N=936	-0.515 [0.424] N=766	-0.647 [0.497] N=591
WHO BMI-for-age z score	0.308 [0.292] N=1,131	0.411 [0.325] N=1,037	0.247 [0.397] N=916	-0.241 [0.450] N=750	-0.287 [0.563] N=582
Stunted: HAZ below -2	-0.079 [0.061] N=1,176	-0.077 [0.068] N=1,074	-0.075 [0.077] N=949	-0.121 [0.082] N=776	-0.044 [0.109] N=602
Underweight: WAZ below -2	0.032 [0.029] N=1,356	0.031 [0.034] N=1,233	0.027 [0.037] N=1,088	0.035 [0.048] N=890	**0.117* [0.062] N=687**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.012 [0.051] N=1,155	-0.003 [0.055] N=1,058	-0.007 [0.063] N=936	0.028 [0.081] N=766	-0.003 [0.102] N=591
Overweight: WLZ/WHZ above +2, 0-60 months	0.016 [0.052] N=1,155	0.008 [0.058] N=1,058	-0.035 [0.066] N=936	-0.053 [0.077] N=766	-0.117 [0.085] N=591
Obese: WLZ/WHZ above +3, 0-60 months	-0.019 [0.039] N=1,155	-0.031 [0.045] N=1,058	-0.017 [0.050] N=936	-0.020 [0.055] N=766	-0.031 [0.068] N=591
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.071 [0.070] N=1,119	-0.066 [0.076] N=1,026	-0.081 [0.085] N=908	-0.098 [0.099] N=743	-0.020 [0.135] N=575
BMI-for-age z score below -2	-0.042 [0.051] N=1,131	-0.049 [0.054] N=1,037	-0.046 [0.064] N=916	0.022 [0.080] N=750	-0.012 [0.101] N=582

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.560	0.198 [0.260] N=1,392	0.147 [0.285] N=1,290	0.158 [0.312] N=1,165	0.322 [0.321] N=992	0.122 [0.349] N=818
WHO weight-for-age z score	-0.050	-0.121 [0.134] N=1,594	-0.105 [0.147] N=1,471	-0.079 [0.154] N=1,326	-0.085 [0.170] N=1,128	-0.289 [0.186] N=925
WHO weight-for-length/height z score, 0-60 months	0.365	0.049 [0.215] N=1,368	0.006 [0.231] N=1,271	-0.067 [0.258] N=1,149	-0.151 [0.289] N=979	-0.120 [0.319] N=804
WHO BMI-for-age z score	0.383	0.206 [0.253] N=1,337	0.245 [0.273] N=1,243	0.159 [0.307] N=1,122	-0.050 [0.342] N=956	-0.016 [0.385] N=788
Stunted: HAZ below -2	0.271 P=368	-0.058 [0.043] N=1,392	-0.058 [0.046] N=1,290	-0.051 [0.049] N=1,165	-0.060 [0.051] N=992	-0.016 [0.055] N=818
Underweight: WAZ below -2	0.075 P=121	**0.050** [0.022] N=1,594**	**0.049** [0.024] N=1,471**	**0.049* [0.025] N=1,326**	**0.055* [0.029] N=1,128**	**0.088** [0.034] N=925**
Wasted: WLZ/WHZ below -2, 0-60 months	0.107 P=151	-0.023 [0.035] N=1,368	-0.020 [0.037] N=1,271	-0.021 [0.039] N=1,149	-0.010 [0.043] N=979	-0.027 [0.045] N=804
Overweight: WLZ/WHZ above +2, 0-60 months	0.197 P=262	0.030 [0.047] N=1,368	0.027 [0.051] N=1,271	0.009 [0.056] N=1,149	0.006 [0.062] N=979	-0.004 [0.070] N=804
Obese: WLZ/WHZ above +3, 0-60 months	0.089 P=121	-0.002 [0.028] N=1,368	-0.010 [0.030] N=1,271	-0.003 [0.032] N=1,149	-0.005 [0.033] N=979	-0.004 [0.037] N=804
CIAF: any stunting, underweight, or wasting, 0-60 months	0.353 P=463	-0.074 [0.045] N=1,325	-0.074 [0.047] N=1,232	-0.075 [0.049] N=1,114	-0.075 [0.051] N=949	-0.036 [0.057] N=781
BMI-for-age z score below -2	0.118 P=160	-0.022 [0.039] N=1,337	-0.025 [0.041] N=1,243	-0.016 [0.044] N=1,122	0.020 [0.048] N=956	0.011 [0.052] N=788

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.102 [0.363] N=1,176	-0.043 [0.415] N=1,074	-0.001 [0.495] N=949	0.404 [0.520] N=776	-0.117 [0.660] N=602
WHO weight-for-age z score	-0.239 [0.179] N=1,356	-0.242 [0.200] N=1,233	-0.207 [0.216] N=1,088	-0.216 [0.276] N=890	**-0.797** [0.319] N=687**
WHO weight-for-length/height z score, 0-60 months	-0.002 [0.265] N=1,155	-0.069 [0.293] N=1,058	-0.249 [0.359] N=936	-0.491 [0.428] N=766	-0.551 [0.501] N=591
WHO BMI-for-age z score	0.310 [0.292] N=1,131	0.416 [0.321] N=1,037	0.256 [0.394] N=916	-0.180 [0.451] N=750	-0.152 [0.567] N=582
Stunted: HAZ below -2	-0.074 [0.061] N=1,176	-0.077 [0.068] N=1,074	-0.078 [0.077] N=949	-0.123 [0.081] N=776	-0.036 [0.109] N=602
Underweight: WAZ below -2	0.035 [0.030] N=1,356	0.033 [0.034] N=1,233	0.032 [0.037] N=1,088	0.037 [0.048] N=890	**0.121* [0.065] N=687**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.010 [0.051] N=1,155	-0.006 [0.055] N=1,058	-0.008 [0.063] N=936	0.027 [0.081] N=766	-0.009 [0.100] N=591
Overweight: WLZ/WHZ above +2, 0-60 months	0.017 [0.051] N=1,155	0.009 [0.057] N=1,058	-0.034 [0.064] N=936	-0.045 [0.076] N=766	-0.098 [0.084] N=591
Obese: WLZ/WHZ above +3, 0-60 months	-0.015 [0.039] N=1,155	-0.028 [0.044] N=1,058	-0.014 [0.050] N=936	-0.014 [0.055] N=766	-0.021 [0.068] N=591
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.063 [0.071] N=1,119	-0.068 [0.076] N=1,026	-0.079 [0.084] N=908	-0.094 [0.098] N=743	-0.009 [0.135] N=575
BMI-for-age z score below -2	-0.043 [0.051] N=1,131	-0.057 [0.054] N=1,037	-0.054 [0.063] N=916	0.012 [0.079] N=750	-0.025 [0.099] N=582

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.574	0.227 [0.279] N=1,297	0.174 [0.301] N=1,208	0.178 [0.326] N=1,089	0.328 [0.329] N=933	0.196 [0.349] N=768
WHO weight-for-age z score	-0.047	-0.009 [0.145] N=1,480	0.010 [0.157] N=1,372	0.044 [0.164] N=1,235	0.041 [0.176] N=1,055	-0.113 [0.202] N=865
WHO weight-for-length/height z score, 0-60 months	0.371	0.179 [0.215] N=1,271	0.149 [0.229] N=1,188	0.059 [0.258] N=1,073	-0.016 [0.281] N=920	0.064 [0.308] N=754
WHO BMI-for-age z score	0.390	0.344 [0.252] N=1,243	0.357 [0.268] N=1,161	0.265 [0.302] N=1,048	0.085 [0.325] N=899	0.168 [0.360] N=739
Stunted: HAZ below -2	0.272 P=342	-0.064 [0.048] N=1,297	-0.070 [0.051] N=1,208	-0.059 [0.054] N=1,089	-0.071 [0.057] N=933	-0.034 [0.061] N=768
Underweight: WAZ below -2	0.075 P=113	0.032 [0.024] N=1,480	0.030 [0.026] N=1,372	0.027 [0.028] N=1,235	0.035 [0.030] N=1,055	0.061 [0.038] N=865
Wasted: WLZ/WHZ below -2, 0-60 months	0.106 P=139	-0.039 [0.036] N=1,271	-0.038 [0.036] N=1,188	-0.035 [0.038] N=1,073	-0.023 [0.043] N=920	-0.035 [0.044] N=754
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=243	0.041 [0.049] N=1,271	0.037 [0.053] N=1,188	0.020 [0.059] N=1,073	0.017 [0.065] N=920	0.018 [0.073] N=754
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=116	-0.005 [0.031] N=1,271	-0.008 [0.033] N=1,188	0.002 [0.035] N=1,073	0.001 [0.036] N=920	0.011 [0.039] N=754
CIAF: any stunting, underweight, or wasting, 0-60 months	0.353 P=429	**-0.095* [0.049] N=1,233**	**-0.103** [0.051] N=1,153**	**-0.104* [0.054] N=1,041**	**-0.114* [0.058] N=893**	-0.079 [0.065] N=733
BMI-for-age z score below -2	0.116 P=147	-0.039 [0.038] N=1,243	-0.041 [0.039] N=1,161	-0.033 [0.042] N=1,048	0.002 [0.047] N=899	-0.003 [0.051] N=739

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.157 [0.383] N=1,094	0.034 [0.440] N=1,005	0.047 [0.524] N=886	0.442 [0.560] N=730	0.088 [0.727] N=565
WHO weight-for-age z score	-0.188 [0.191] N=1,257	-0.196 [0.212] N=1,149	-0.164 [0.228] N=1,012	-0.228 [0.274] N=832	**-0.788** [0.361] N=642**
WHO weight-for-length/height z score, 0-60 months	0.028 [0.264] N=1,071	-0.054 [0.290] N=988	-0.284 [0.354] N=873	-0.531 [0.427] N=720	-0.534 [0.499] N=554
WHO BMI-for-age z score	0.348 [0.303] N=1,050	0.373 [0.328] N=968	0.178 [0.397] N=855	-0.245 [0.460] N=706	-0.160 [0.571] N=546
Stunted: HAZ below -2	-0.085 [0.066] N=1,094	-0.100 [0.072] N=1,005	-0.090 [0.082] N=886	-0.139 [0.089] N=730	-0.077 [0.121] N=565
Underweight: WAZ below -2	0.023 [0.032] N=1,257	0.021 [0.036] N=1,149	0.019 [0.040] N=1,012	0.041 [0.045] N=832	**0.127* [0.066] N=642**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.028 [0.052] N=1,071	-0.024 [0.053] N=988	-0.019 [0.061] N=873	0.003 [0.079] N=720	-0.054 [0.099] N=554
Overweight: WLZ/WHZ above +2, 0-60 months	0.013 [0.054] N=1,071	-0.005 [0.059] N=988	-0.050 [0.068] N=873	-0.078 [0.084] N=720	-0.124 [0.097] N=554
Obese: WLZ/WHZ above +3, 0-60 months	-0.033 [0.042] N=1,071	-0.042 [0.049] N=988	-0.028 [0.055] N=873	-0.031 [0.060] N=720	-0.033 [0.073] N=554
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.087 [0.074] N=1,040	-0.103 [0.078] N=960	-0.112 [0.087] N=848	-0.147 [0.101] N=700	-0.101 [0.143] N=540
BMI-for-age z score below -2	-0.055 [0.051] N=1,050	-0.063 [0.051] N=968	-0.056 [0.061] N=855	0.002 [0.078] N=706	-0.048 [0.099] N=546

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.574	0.216 [0.279] N=1,297	0.164 [0.300] N=1,208	0.157 [0.328] N=1,089	0.297 [0.337] N=933	0.128 [0.365] N=768
WHO weight-for-age z score	-0.047	-0.017 [0.144] N=1,480	0.003 [0.156] N=1,372	0.029 [0.163] N=1,235	0.019 [0.176] N=1,055	-0.151 [0.201] N=865
WHO weight-for-length/height z score, 0-60 months	0.371	0.165 [0.214] N=1,271	0.134 [0.228] N=1,188	0.046 [0.258] N=1,073	-0.024 [0.285] N=920	0.062 [0.317] N=754
WHO BMI-for-age z score	0.390	0.330 [0.252] N=1,243	0.343 [0.268] N=1,161	0.254 [0.303] N=1,048	0.079 [0.333] N=899	0.168 [0.376] N=739
Stunted: HAZ below -2	0.272 P=342	-0.062 [0.048] N=1,297	-0.068 [0.051] N=1,208	-0.056 [0.054] N=1,089	-0.067 [0.057] N=933	-0.025 [0.061] N=768
Underweight: WAZ below -2	0.075 P=113	0.034 [0.024] N=1,480	0.031 [0.026] N=1,372	0.030 [0.027] N=1,235	0.041 [0.030] N=1,055	**0.072* [0.037] N=865**
Wasted: WLZ/WHZ below -2, 0-60 months	0.106 P=139	-0.038 [0.036] N=1,271	-0.037 [0.036] N=1,188	-0.036 [0.039] N=1,073	-0.026 [0.045] N=920	-0.043 [0.048] N=754
Overweight: WLZ/WHZ above +2, 0-60 months	0.198 P=243	0.040 [0.049] N=1,271	0.035 [0.053] N=1,188	0.017 [0.059] N=1,073	0.013 [0.065] N=920	0.017 [0.072] N=754
Obese: WLZ/WHZ above +3, 0-60 months	0.092 P=116	-0.005 [0.030] N=1,271	-0.008 [0.033] N=1,188	0.002 [0.035] N=1,073	0.002 [0.035] N=920	0.012 [0.039] N=754
CIAF: any stunting, underweight, or wasting, 0-60 months	0.353 P=429	**-0.093* [0.049] N=1,233**	**-0.101** [0.051] N=1,153**	**-0.103* [0.054] N=1,041**	**-0.112* [0.058] N=893**	-0.076 [0.063] N=733
BMI-for-age z score below -2	0.116 P=147	-0.037 [0.039] N=1,243	-0.040 [0.040] N=1,161	-0.033 [0.043] N=1,048	0.000 [0.048] N=899	-0.009 [0.054] N=739

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.172 [0.383] N=1,094	0.049 [0.438] N=1,005	0.067 [0.519] N=886	0.442 [0.567] N=730	0.052 [0.752] N=565
WHO weight-for-age z score	-0.157 [0.188] N=1,257	-0.160 [0.209] N=1,149	-0.137 [0.224] N=1,012	-0.210 [0.273] N=832	**-0.812** [0.362] N=642**
WHO weight-for-length/height z score, 0-60 months	0.057 [0.259] N=1,071	-0.012 [0.285] N=988	-0.245 [0.352] N=873	-0.508 [0.427] N=720	-0.553 [0.497] N=554
WHO BMI-for-age z score	0.368 [0.295] N=1,050	0.407 [0.319] N=968	0.206 [0.391] N=855	-0.228 [0.452] N=706	-0.192 [0.559] N=546
Stunted: HAZ below -2	-0.083 [0.066] N=1,094	-0.096 [0.072] N=1,005	-0.090 [0.081] N=886	-0.138 [0.089] N=730	-0.075 [0.122] N=565
Underweight: WAZ below -2	0.019 [0.031] N=1,257	0.015 [0.036] N=1,149	0.013 [0.039] N=1,012	0.035 [0.045] N=832	**0.130** [0.064] N=642**
Wasted: WLZ/WHZ below -2, 0-60 months	-0.029 [0.051] N=1,071	-0.026 [0.053] N=988	-0.021 [0.061] N=873	0.002 [0.078] N=720	-0.056 [0.097] N=554
Overweight: WLZ/WHZ above +2, 0-60 months	0.019 [0.054] N=1,071	0.004 [0.058] N=988	-0.042 [0.068] N=873	-0.070 [0.083] N=720	-0.125 [0.096] N=554
Obese: WLZ/WHZ above +3, 0-60 months	-0.033 [0.042] N=1,071	-0.042 [0.049] N=988	-0.026 [0.055] N=873	-0.030 [0.061] N=720	-0.033 [0.073] N=554
CIAF: any stunting, underweight, or wasting, 0-60 months	-0.084 [0.073] N=1,040	-0.100 [0.077] N=960	-0.110 [0.086] N=848	-0.143 [0.100] N=700	-0.100 [0.141] N=540
BMI-for-age z score below -2	-0.053 [0.051] N=1,050	-0.064 [0.051] N=968	-0.056 [0.060] N=855	0.002 [0.078] N=706	-0.049 [0.098] N=546

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.440	-0.148 [0.294] N=1,256	-0.225 [0.306] N=1,127	-0.258 [0.316] N=986	-0.506 [0.348] N=835	-0.620 [0.391] N=683
WHO weight-for-age z score	-0.033	0.085 [0.143] N=1,419	0.106 [0.150] N=1,285	-0.018 [0.147] N=1,124	0.057 [0.155] N=944	0.021 [0.177] N=764
WHO weight-for-length/height z score, 0-60 months	0.361	0.125 [0.205] N=1,241	0.063 [0.217] N=1,117	0.000 [0.236] N=979	0.240 [0.249] N=827	0.099 [0.278] N=676
WHO BMI-for-age z score	0.340	0.162 [0.236] N=1,205	0.102 [0.246] N=1,085	-0.006 [0.265] N=952	0.154 [0.287] N=803	0.070 [0.313] N=660
Stunted: HAZ below -2	0.251 P=319	0.012 [0.060] N=1,256	0.008 [0.062] N=1,127	0.016 [0.063] N=986	0.050 [0.068] N=835	0.068 [0.073] N=683
Underweight: WAZ below -2	0.072 P=103	-0.004 [0.032] N=1,419	-0.011 [0.034] N=1,285	0.007 [0.036] N=1,124	-0.001 [0.037] N=944	0.001 [0.041] N=764
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=137	0.026 [0.037] N=1,241	0.029 [0.040] N=1,117	0.054 [0.044] N=979	-0.017 [0.042] N=827	-0.005 [0.048] N=676
Overweight: WLZ/WHZ above +2, 0-60 months	0.189 P=239	**0.091** [0.045] N=1,241**	**0.085* [0.048] N=1,117**	0.083 [0.051] N=979	**0.104* [0.058] N=827**	0.073 [0.059] N=676
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=115	0.043 [0.038] N=1,241	0.030 [0.039] N=1,117	0.024 [0.042] N=979	0.029 [0.049] N=827	-0.013 [0.048] N=676
CIAF: any stunting, underweight, or wasting, 0-60 months	0.341 P=412	0.039 [0.064] N=1,200	0.028 [0.065] N=1,076	0.065 [0.067] N=943	0.034 [0.070] N=798	0.074 [0.074] N=654
BMI-for-age z score below -2	0.121 P=145	0.029 [0.043] N=1,205	0.034 [0.047] N=1,085	0.041 [0.051] N=952	-0.013 [0.052] N=803	-0.006 [0.057] N=660

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.018 [0.414] N=1,083	-0.066 [0.449] N=954	-0.088 [0.507] N=813	-0.676 [0.621] N=662	-1.047 [0.774] N=510
WHO weight-for-age z score	0.144 [0.196] N=1,229	0.166 [0.213] N=1,095	-0.056 [0.211] N=934	0.108 [0.216] N=754	0.142 [0.296] N=574
WHO weight-for-length/height z score, 0-60 months	-0.006 [0.306] N=1,068	-0.143 [0.334] N=944	-0.282 [0.386] N=806	0.246 [0.433] N=654	-0.123 [0.537] N=503
WHO BMI-for-age z score	-0.142 [0.338] N=1,043	-0.276 [0.364] N=923	-0.510 [0.418] N=790	-0.198 [0.482] N=641	-0.447 [0.555] N=498
Stunted: HAZ below -2	0.008 [0.098] N=1,083	-0.006 [0.103] N=954	0.010 [0.116] N=813	0.102 [0.140] N=662	0.189 [0.166] N=510
Underweight: WAZ below -2	-0.026 [0.045] N=1,229	-0.037 [0.049] N=1,095	-0.007 [0.054] N=934	-0.039 [0.054] N=754	-0.055 [0.063] N=574
Wasted: WLZ/WHZ below -2, 0-60 months	0.078 [0.049] N=1,068	0.091 [0.057] N=944	**0.148** [0.064] N=806**	0.003 [0.061] N=654	0.030 [0.074] N=503
Overweight: WLZ/WHZ above +2, 0-60 months	**0.128** [0.058] N=1,068**	**0.116* [0.062] N=944**	**0.127* [0.068] N=806**	**0.193** [0.082] N=654**	0.125 [0.095] N=503
Obese: WLZ/WHZ above +3, 0-60 months	**0.100** [0.041] N=1,068**	**0.080* [0.041] N=944**	**0.081* [0.048] N=806**	**0.109* [0.065] N=654**	-0.003 [0.069] N=503
CIAF: any stunting, underweight, or wasting, 0-60 months	0.061 [0.101] N=1,033	0.042 [0.107] N=909	0.115 [0.121] N=776	0.065 [0.142] N=631	0.169 [0.163] N=487
BMI-for-age z score below -2	**0.119** [0.052] N=1,043**	**0.137** [0.062] N=923**	**0.165** [0.073] N=790**	0.051 [0.072] N=641	0.074 [0.088] N=498

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.440	-0.152 [0.293] N=1,256	-0.228 [0.305] N=1,127	-0.261 [0.314] N=986	-0.509 [0.347] N=835	-0.624 [0.390] N=683
WHO weight-for-age z score	-0.033	0.091 [0.143] N=1,419	0.111 [0.149] N=1,285	-0.012 [0.146] N=1,124	0.057 [0.155] N=944	0.021 [0.176] N=764
WHO weight-for-length/height z score, 0-60 months	0.361	0.131 [0.205] N=1,241	0.069 [0.217] N=1,117	0.005 [0.235] N=979	0.242 [0.249] N=827	0.105 [0.278] N=676
WHO BMI-for-age z score	0.340	0.163 [0.236] N=1,205	0.103 [0.246] N=1,085	-0.009 [0.264] N=952	0.154 [0.287] N=803	0.070 [0.314] N=660
Stunted: HAZ below -2	0.251 P=319	0.010 [0.060] N=1,256	0.005 [0.062] N=1,127	0.012 [0.063] N=986	0.049 [0.068] N=835	0.068 [0.072] N=683
Underweight: WAZ below -2	0.072 P=103	-0.006 [0.032] N=1,419	-0.013 [0.033] N=1,285	0.006 [0.035] N=1,124	-0.002 [0.037] N=944	0.001 [0.041] N=764
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=137	0.023 [0.038] N=1,241	0.026 [0.041] N=1,117	0.050 [0.044] N=979	-0.018 [0.042] N=827	-0.006 [0.048] N=676
Overweight: WLZ/WHZ above +2, 0-60 months	0.189 P=239	**0.092** [0.045] N=1,241**	**0.085* [0.048] N=1,117**	0.082 [0.051] N=979	**0.103* [0.058] N=827**	0.073 [0.059] N=676
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=115	0.041 [0.038] N=1,241	0.029 [0.038] N=1,117	0.022 [0.042] N=979	0.028 [0.049] N=827	-0.013 [0.048] N=676
CIAF: any stunting, underweight, or wasting, 0-60 months	0.341 P=412	0.034 [0.064] N=1,200	0.020 [0.066] N=1,076	0.056 [0.067] N=943	0.032 [0.071] N=798	0.073 [0.075] N=654
BMI-for-age z score below -2	0.121 P=145	0.028 [0.043] N=1,205	0.033 [0.047] N=1,085	0.040 [0.051] N=952	-0.014 [0.052] N=803	-0.006 [0.057] N=660

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.008 [0.411] N=1,083	-0.077 [0.445] N=954	-0.102 [0.500] N=813	-0.709 [0.608] N=662	-1.107 [0.763] N=510
WHO weight-for-age z score	0.158 [0.193] N=1,229	0.182 [0.209] N=1,095	-0.041 [0.206] N=934	0.122 [0.211] N=754	0.174 [0.288] N=574
WHO weight-for-length/height z score, 0-60 months	0.008 [0.307] N=1,068	-0.128 [0.335] N=944	-0.271 [0.386] N=806	0.262 [0.436] N=654	-0.050 [0.552] N=503
WHO BMI-for-age z score	-0.130 [0.339] N=1,043	-0.261 [0.365] N=923	-0.501 [0.419] N=790	-0.188 [0.486] N=641	-0.392 [0.573] N=498
Stunted: HAZ below -2	0.005 [0.098] N=1,083	-0.013 [0.102] N=954	0.001 [0.115] N=813	0.098 [0.139] N=662	0.188 [0.164] N=510
Underweight: WAZ below -2	-0.028 [0.045] N=1,229	-0.041 [0.048] N=1,095	-0.009 [0.053] N=934	-0.042 [0.054] N=754	-0.060 [0.062] N=574
Wasted: WLZ/WHZ below -2, 0-60 months	0.072 [0.050] N=1,068	0.082 [0.058] N=944	**0.137** [0.065] N=806**	-0.004 [0.061] N=654	0.015 [0.076] N=503
Overweight: WLZ/WHZ above +2, 0-60 months	**0.130** [0.058] N=1,068**	**0.118* [0.062] N=944**	**0.126* [0.068] N=806**	**0.190** [0.082] N=654**	0.126 [0.095] N=503
Obese: WLZ/WHZ above +3, 0-60 months	**0.098** [0.040] N=1,068**	**0.076* [0.041] N=944**	0.076 [0.047] N=806	0.102 [0.065] N=654	-0.010 [0.067] N=503
CIAF: any stunting, underweight, or wasting, 0-60 months	0.049 [0.102] N=1,033	0.022 [0.108] N=909	0.089 [0.122] N=776	0.048 [0.142] N=631	0.146 [0.163] N=487
BMI-for-age z score below -2	**0.115** [0.052] N=1,043**	**0.130** [0.062] N=923**	**0.157** [0.072] N=790**	0.047 [0.072] N=641	0.065 [0.089] N=498

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.440	-0.165 [0.289] N=1,256	-0.228 [0.301] N=1,127	-0.259 [0.311] N=986	-0.504 [0.340] N=835	**-0.643* [0.380] N=683**
WHO weight-for-age z score	-0.033	0.079 [0.143] N=1,419	0.106 [0.148] N=1,285	-0.009 [0.147] N=1,124	0.062 [0.154] N=944	0.023 [0.174] N=764
WHO weight-for-length/height z score, 0-60 months	0.361	0.124 [0.205] N=1,241	0.062 [0.218] N=1,117	0.003 [0.236] N=979	0.236 [0.248] N=827	0.110 [0.278] N=676
WHO BMI-for-age z score	0.340	0.151 [0.238] N=1,205	0.091 [0.248] N=1,085	-0.014 [0.264] N=952	0.150 [0.285] N=803	0.081 [0.314] N=660
Stunted: HAZ below -2	0.251 P=319	0.013 [0.060] N=1,256	0.004 [0.061] N=1,127	0.008 [0.062] N=986	0.045 [0.067] N=835	0.068 [0.071] N=683
Underweight: WAZ below -2	0.072 P=103	-0.004 [0.032] N=1,419	-0.012 [0.034] N=1,285	0.005 [0.035] N=1,124	-0.003 [0.036] N=944	0.001 [0.040] N=764
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=137	0.026 [0.038] N=1,241	0.028 [0.041] N=1,117	0.050 [0.045] N=979	-0.019 [0.043] N=827	-0.007 [0.049] N=676
Overweight: WLZ/WHZ above +2, 0-60 months	0.189 P=239	**0.093** [0.045] N=1,241**	**0.084* [0.048] N=1,117**	0.081 [0.051] N=979	**0.101* [0.058] N=827**	0.073 [0.060] N=676
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=115	0.042 [0.038] N=1,241	0.028 [0.038] N=1,117	0.021 [0.042] N=979	0.026 [0.049] N=827	-0.014 [0.049] N=676
CIAF: any stunting, underweight, or wasting, 0-60 months	0.341 P=412	0.040 [0.064] N=1,200	0.023 [0.066] N=1,076	0.054 [0.068] N=943	0.028 [0.072] N=798	0.072 [0.075] N=654
BMI-for-age z score below -2	0.121 P=145	0.031 [0.043] N=1,205	0.036 [0.047] N=1,085	0.041 [0.051] N=952	-0.013 [0.052] N=803	-0.004 [0.058] N=660

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.002 [0.405] N=1,083	-0.077 [0.441] N=954	-0.089 [0.492] N=813	-0.689 [0.599] N=662	-1.102 [0.747] N=510
WHO weight-for-age z score	0.135 [0.195] N=1,229	0.165 [0.211] N=1,095	-0.041 [0.212] N=934	0.121 [0.221] N=754	0.168 [0.290] N=574
WHO weight-for-length/height z score, 0-60 months	-0.015 [0.306] N=1,068	-0.147 [0.334] N=944	-0.289 [0.384] N=806	0.242 [0.433] N=654	-0.066 [0.544] N=503
WHO BMI-for-age z score	-0.149 [0.338] N=1,043	-0.279 [0.364] N=923	-0.526 [0.412] N=790	-0.223 [0.475] N=641	-0.406 [0.558] N=498
Stunted: HAZ below -2	0.004 [0.097] N=1,083	-0.015 [0.102] N=954	-0.007 [0.114] N=813	0.086 [0.138] N=662	0.175 [0.160] N=510
Underweight: WAZ below -2	-0.027 [0.046] N=1,229	-0.040 [0.049] N=1,095	-0.012 [0.055] N=934	-0.046 [0.056] N=754	-0.064 [0.064] N=574
Wasted: WLZ/WHZ below -2, 0-60 months	0.073 [0.050] N=1,068	0.082 [0.058] N=944	**0.136** [0.066] N=806**	-0.007 [0.062] N=654	0.010 [0.078] N=503
Overweight: WLZ/WHZ above +2, 0-60 months	**0.131** [0.059] N=1,068**	**0.119* [0.062] N=944**	**0.124* [0.069] N=806**	**0.188** [0.083] N=654**	0.125 [0.097] N=503
Obese: WLZ/WHZ above +3, 0-60 months	**0.095** [0.041] N=1,068**	**0.073* [0.041] N=944**	0.072 [0.048] N=806	0.097 [0.066] N=654	-0.016 [0.067] N=503
CIAF: any stunting, underweight, or wasting, 0-60 months	0.051 [0.102] N=1,033	0.021 [0.108] N=909	0.082 [0.122] N=776	0.037 [0.142] N=631	0.132 [0.160] N=487
BMI-for-age z score below -2	**0.114** [0.053] N=1,043**	**0.129** [0.063] N=923**	**0.154** [0.072] N=790**	0.040 [0.072] N=641	0.057 [0.091] N=498

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.440	-0.133 [0.285] N=1,256	-0.194 [0.297] N=1,127	-0.222 [0.308] N=986	-0.487 [0.334] N=835	-0.630 [0.381] N=683
WHO weight-for-age z score	-0.033	0.088 [0.145] N=1,419	0.113 [0.152] N=1,285	-0.000 [0.150] N=1,124	0.063 [0.160] N=944	0.020 [0.180] N=764
WHO weight-for-length/height z score, 0-60 months	0.361	0.113 [0.205] N=1,241	0.055 [0.217] N=1,117	-0.003 [0.234] N=979	0.245 [0.247] N=827	0.123 [0.278] N=676
WHO BMI-for-age z score	0.340	0.144 [0.236] N=1,205	0.085 [0.247] N=1,085	-0.021 [0.262] N=952	0.153 [0.282] N=803	0.097 [0.313] N=660
Stunted: HAZ below -2	0.251 P=319	0.007 [0.058] N=1,256	-0.001 [0.060] N=1,127	0.006 [0.062] N=986	0.046 [0.066] N=835	0.070 [0.071] N=683
Underweight: WAZ below -2	0.072 P=103	-0.005 [0.032] N=1,419	-0.013 [0.034] N=1,285	0.004 [0.036] N=1,124	-0.003 [0.037] N=944	-0.001 [0.042] N=764
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=137	0.026 [0.038] N=1,241	0.028 [0.041] N=1,117	0.053 [0.044] N=979	-0.017 [0.042] N=827	-0.008 [0.048] N=676
Overweight: WLZ/WHZ above +2, 0-60 months	0.189 P=239	**0.093** [0.045] N=1,241**	**0.086* [0.047] N=1,117**	**0.085* [0.050] N=979**	**0.107* [0.056] N=827**	0.079 [0.057] N=676
Obese: WLZ/WHZ above +3, 0-60 months	0.088 P=115	0.044 [0.037] N=1,241	0.032 [0.037] N=1,117	0.025 [0.041] N=979	0.032 [0.048] N=827	-0.009 [0.047] N=676
CIAF: any stunting, underweight, or wasting, 0-60 months	0.341 P=412	0.037 [0.064] N=1,200	0.022 [0.066] N=1,076	0.056 [0.068] N=943	0.034 [0.071] N=798	0.079 [0.075] N=654
BMI-for-age z score below -2	0.121 P=145	0.028 [0.043] N=1,205	0.033 [0.047] N=1,085	0.040 [0.051] N=952	-0.015 [0.052] N=803	-0.011 [0.056] N=660

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.041 [0.390] N=1,083	-0.023 [0.424] N=954	-0.013 [0.475] N=813	-0.662 [0.569] N=662	-1.146 [0.700] N=510
WHO weight-for-age z score	0.170 [0.189] N=1,229	0.198 [0.208] N=1,095	0.000 [0.205] N=934	0.139 [0.213] N=754	0.174 [0.275] N=574
WHO weight-for-length/height z score, 0-60 months	-0.020 [0.306] N=1,068	-0.165 [0.333] N=944	-0.313 [0.382] N=806	0.227 [0.440] N=654	-0.028 [0.576] N=503
WHO BMI-for-age z score	-0.167 [0.337] N=1,043	-0.309 [0.363] N=923	-0.560 [0.412] N=790	-0.234 [0.481] N=641	-0.341 [0.588] N=498
Stunted: HAZ below -2	-0.004 [0.093] N=1,083	-0.025 [0.099] N=954	-0.015 [0.111] N=813	0.092 [0.134] N=662	0.203 [0.155] N=510
Underweight: WAZ below -2	-0.030 [0.045] N=1,229	-0.041 [0.048] N=1,095	-0.015 [0.054] N=934	-0.043 [0.054] N=754	-0.064 [0.063] N=574
Wasted: WLZ/WHZ below -2, 0-60 months	0.077 [0.050] N=1,068	0.090 [0.057] N=944	**0.146** [0.064] N=806**	0.002 [0.062] N=654	0.016 [0.077] N=503
Overweight: WLZ/WHZ above +2, 0-60 months	**0.128** [0.058] N=1,068**	**0.116* [0.062] N=944**	**0.123* [0.069] N=806**	**0.187** [0.083] N=654**	0.137 [0.099] N=503
Obese: WLZ/WHZ above +3, 0-60 months	**0.098** [0.041] N=1,068**	**0.077* [0.041] N=944**	0.074 [0.047] N=806	0.098 [0.065] N=654	-0.011 [0.069] N=503
CIAF: any stunting, underweight, or wasting, 0-60 months	0.047 [0.099] N=1,033	0.019 [0.107] N=909	0.082 [0.120] N=776	0.048 [0.140] N=631	0.168 [0.156] N=487
BMI-for-age z score below -2	**0.117** [0.053] N=1,043**	**0.134** [0.063] N=923**	**0.162** [0.072] N=790**	0.051 [0.072] N=641	0.066 [0.090] N=498

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.450	-0.174 [0.304] N=1,170	-0.257 [0.325] N=1,049	-0.252 [0.334] N=923	-0.522 [0.365] N=780	**-0.683* [0.406] N=639**
WHO weight-for-age z score	-0.033	0.122 [0.151] N=1,317	0.159 [0.160] N=1,191	0.013 [0.155] N=1,047	0.060 [0.166] N=880	-0.028 [0.181] N=715
WHO weight-for-length/height z score, 0-60 months	0.364	0.213 [0.223] N=1,153	0.160 [0.240] N=1,037	0.043 [0.256] N=912	0.256 [0.272] N=770	0.090 [0.297] N=631
WHO BMI-for-age z score	0.341	0.190 [0.258] N=1,120	0.142 [0.274] N=1,008	-0.027 [0.289] N=888	0.117 [0.312] N=749	-0.013 [0.334] N=617
Stunted: HAZ below -2	0.252 P=296	0.002 [0.063] N=1,170	0.000 [0.065] N=1,049	-0.006 [0.066] N=923	0.027 [0.070] N=780	0.054 [0.075] N=639
Underweight: WAZ below -2	0.072 P=96	-0.013 [0.033] N=1,317	-0.023 [0.035] N=1,191	-0.005 [0.037] N=1,047	-0.009 [0.038] N=880	0.001 [0.042] N=715
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=127	0.025 [0.040] N=1,153	0.026 [0.043] N=1,037	0.056 [0.046] N=912	-0.004 [0.045] N=770	0.014 [0.050] N=631
Overweight: WLZ/WHZ above +2, 0-60 months	0.190 P=222	**0.100** [0.046] N=1,153**	**0.093* [0.048] N=1,037**	0.082 [0.052] N=912	**0.097* [0.058] N=770**	0.065 [0.059] N=631
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=110	0.050 [0.037] N=1,153	0.036 [0.038] N=1,037	0.027 [0.041] N=912	0.038 [0.049] N=770	-0.002 [0.047] N=631
CIAF: any stunting, underweight, or wasting, 0-60 months	0.342 P=383	0.031 [0.068] N=1,117	0.017 [0.069] N=1,001	0.046 [0.071] N=881	0.019 [0.075] N=744	0.077 [0.079] N=611
BMI-for-age z score below -2	0.120 P=134	0.036 [0.046] N=1,120	0.038 [0.051] N=1,008	0.048 [0.054] N=888	0.002 [0.056] N=749	0.017 [0.060] N=617

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.051 [0.426] N=1,005	-0.037 [0.471] N=884	-0.030 [0.534] N=758	-0.702 [0.638] N=615	-1.192 [0.800] N=474
WHO weight-for-age z score	0.260 [0.195] N=1,137	0.324 [0.210] N=1,011	0.066 [0.204] N=867	0.232 [0.204] N=700	0.227 [0.264] N=535
WHO weight-for-length/height z score, 0-60 months	0.098 [0.337] N=989	-0.009 [0.376] N=873	-0.232 [0.427] N=748	0.265 [0.475] N=606	-0.045 [0.613] N=467
WHO BMI-for-age z score	-0.085 [0.375] N=965	-0.187 [0.416] N=853	-0.517 [0.470] N=733	-0.203 [0.526] N=594	-0.476 [0.623] N=462
Stunted: HAZ below -2	-0.012 [0.103] N=1,005	-0.025 [0.109] N=884	-0.029 [0.121] N=758	0.050 [0.143] N=615	0.153 [0.167] N=474
Underweight: WAZ below -2	-0.047 [0.046] N=1,137	-0.066 [0.049] N=1,011	-0.040 [0.054] N=867	-0.075 [0.054] N=700	-0.087 [0.060] N=535
Wasted: WLZ/WHZ below -2, 0-60 months	0.070 [0.054] N=989	0.074 [0.062] N=873	**0.137* [0.069] N=748**	0.008 [0.068] N=606	0.035 [0.084] N=467
Overweight: WLZ/WHZ above +2, 0-60 months	**0.142** [0.063] N=989**	**0.131* [0.067] N=873**	**0.128* [0.074] N=748**	**0.182** [0.089] N=606**	0.127 [0.108] N=467
Obese: WLZ/WHZ above +3, 0-60 months	**0.099** [0.045] N=989**	0.076 [0.046] N=873	0.072 [0.052] N=748	0.102 [0.071] N=606	-0.016 [0.076] N=467
CIAF: any stunting, underweight, or wasting, 0-60 months	0.035 [0.109] N=958	0.003 [0.116] N=842	0.064 [0.131] N=722	0.005 [0.153] N=585	0.138 [0.174] N=452
BMI-for-age z score below -2	**0.125** [0.058] N=965**	**0.134* [0.070] N=853**	**0.165** [0.080] N=733**	0.065 [0.081] N=594	0.106 [0.098] N=462

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	-0.450	-0.198 [0.310] N=1,170	-0.283 [0.329] N=1,049	-0.278 [0.340] N=923	-0.568 [0.373] N=780	**-0.736* [0.419] N=639**
WHO weight-for-age z score	-0.033	0.120 [0.151] N=1,317	0.150 [0.161] N=1,191	0.005 [0.155] N=1,047	0.048 [0.166] N=880	-0.041 [0.183] N=715
WHO weight-for-length/height z score, 0-60 months	0.364	0.232 [0.222] N=1,153	0.175 [0.239] N=1,037	0.061 [0.256] N=912	0.279 [0.270] N=770	0.120 [0.297] N=631
WHO BMI-for-age z score	0.341	0.213 [0.260] N=1,120	0.160 [0.276] N=1,008	-0.004 [0.292] N=888	0.144 [0.313] N=749	0.021 [0.338] N=617
Stunted: HAZ below -2	0.252 P=296	0.005 [0.062] N=1,170	0.002 [0.064] N=1,049	-0.003 [0.065] N=923	0.031 [0.070] N=780	0.060 [0.075] N=639
Underweight: WAZ below -2	0.072 P=96	-0.012 [0.033] N=1,317	-0.022 [0.035] N=1,191	-0.004 [0.036] N=1,047	-0.008 [0.038] N=880	0.003 [0.042] N=715
Wasted: WLZ/WHZ below -2, 0-60 months	0.109 P=127	0.022 [0.041] N=1,153	0.022 [0.044] N=1,037	0.051 [0.047] N=912	-0.011 [0.046] N=770	0.007 [0.051] N=631
Overweight: WLZ/WHZ above +2, 0-60 months	0.190 P=222	**0.101** [0.045] N=1,153**	**0.094** [0.047] N=1,037**	0.081 [0.051] N=912	**0.098* [0.057] N=770**	0.068 [0.058] N=631
Obese: WLZ/WHZ above +3, 0-60 months	0.091 P=110	0.051 [0.037] N=1,153	0.036 [0.038] N=1,037	0.026 [0.041] N=912	0.037 [0.048] N=770	-0.002 [0.047] N=631
CIAF: any stunting, underweight, or wasting, 0-60 months	0.342 P=383	0.031 [0.066] N=1,117	0.015 [0.068] N=1,001	0.042 [0.069] N=881	0.017 [0.072] N=744	0.076 [0.076] N=611
BMI-for-age z score below -2	0.120 P=134	0.032 [0.047] N=1,120	0.034 [0.052] N=1,008	0.042 [0.055] N=888	-0.004 [0.057] N=749	0.010 [0.061] N=617

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO length/height-for-age z score	0.020 [0.432] N=1,005	-0.064 [0.473] N=884	-0.057 [0.535] N=758	-0.802 [0.649] N=615	-1.348 [0.811] N=474
WHO weight-for-age z score	0.251 [0.198] N=1,137	0.303 [0.215] N=1,011	0.043 [0.207] N=867	0.195 [0.210] N=700	0.174 [0.272] N=535
WHO weight-for-length/height z score, 0-60 months	0.102 [0.335] N=989	-0.016 [0.370] N=873	-0.238 [0.423] N=748	0.292 [0.470] N=606	-0.017 [0.604] N=467
WHO BMI-for-age z score	-0.072 [0.375] N=965	-0.186 [0.412] N=853	-0.509 [0.470] N=733	-0.169 [0.527] N=594	-0.395 [0.630] N=462
Stunted: HAZ below -2	-0.008 [0.103] N=1,005	-0.024 [0.108] N=884	-0.028 [0.120] N=758	0.061 [0.143] N=615	0.167 [0.166] N=474
Underweight: WAZ below -2	-0.045 [0.047] N=1,137	-0.064 [0.050] N=1,011	-0.036 [0.054] N=867	-0.067 [0.054] N=700	-0.074 [0.061] N=535
Wasted: WLZ/WHZ below -2, 0-60 months	0.069 [0.054] N=989	0.075 [0.062] N=873	**0.136* [0.069] N=748**	0.002 [0.067] N=606	0.026 [0.082] N=467
Overweight: WLZ/WHZ above +2, 0-60 months	**0.144** [0.062] N=989**	**0.131** [0.065] N=873**	**0.126* [0.072] N=748**	**0.185** [0.087] N=606**	0.128 [0.106] N=467
Obese: WLZ/WHZ above +3, 0-60 months	**0.099** [0.044] N=989**	**0.075* [0.045] N=873**	0.069 [0.050] N=748	0.101 [0.069] N=606	-0.015 [0.075] N=467
CIAF: any stunting, underweight, or wasting, 0-60 months	0.038 [0.108] N=958	0.005 [0.115] N=842	0.062 [0.129] N=722	0.005 [0.151] N=585	0.133 [0.171] N=452
BMI-for-age z score below -2	**0.123** [0.058] N=965**	**0.134* [0.070] N=853**	**0.163** [0.079] N=733**	0.059 [0.080] N=594	0.093 [0.098] N=462

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 12 to 47 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Aid and income controls are not combined. Aid specifications exclude households with a blank or non-yes/no answer on any of the six main aid questions. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.