Table 15 Reduced Form Effects on WHO Anthropometric Outcomes With UNICEF Age Controls

Continuous z scores and prespecified adverse clinical classifications exclude WHO-flagged measurements. Child age is based on 16 December 2022. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 UNICEF age-interval fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.509	0.206 [0.221] N=1,517	0.200 [0.235] N=1,404	0.225 [0.244] N=1,262	0.244 [0.256] N=1,078	0.224 [0.292] N=886
WHO weight-for-age z score	0.024	0.103 [0.130] N=1,679	0.141 [0.137] N=1,550	0.079 [0.137] N=1,395	0.096 [0.160] N=1,189	0.037 [0.162] N=977
WHO weight-for-height z score	0.367	-0.146 [0.194] N=1,440	-0.138 [0.200] N=1,342	-0.260 [0.216] N=1,206	-0.288 [0.240] N=1,030	-0.163 [0.255] N=845
WHO BMI-for-age z score	0.368	-0.173 [0.218] N=1,442	-0.152 [0.230] N=1,341	-0.310 [0.248] N=1,200	-0.386 [0.281] N=1,024	-0.273 [0.304] N=844
Stunted HAZ below -2	0.238 P=353	-0.050 [0.040] N=1,517	-0.054 [0.042] N=1,404	-0.050 [0.044] N=1,262	-0.060 [0.047] N=1,078	-0.018 [0.052] N=886
Underweight WAZ below -2	0.064 P=108	**-0.037* [0.022] N=1,679**	**-0.041* [0.023] N=1,550**	**-0.041* [0.024] N=1,395**	-0.041 [0.026] N=1,189	-0.031 [0.029] N=977
Wasted WHZ below -2	0.106 P=159	-0.019 [0.034] N=1,440	-0.009 [0.035] N=1,342	0.009 [0.038] N=1,206	0.003 [0.043] N=1,030	-0.012 [0.048] N=845
Overweight WHZ above +2	0.196 P=273	-0.021 [0.042] N=1,440	-0.014 [0.044] N=1,342	-0.017 [0.046] N=1,206	-0.024 [0.051] N=1,030	-0.003 [0.056] N=845
Obese WHZ above +3	0.090 P=128	-0.019 [0.029] N=1,440	-0.020 [0.029] N=1,342	-0.017 [0.031] N=1,206	-0.023 [0.034] N=1,030	-0.015 [0.038] N=845
CIAF any stunting underweight or wasting	0.319 P=450	**-0.099** [0.043] N=1,413**	**-0.089** [0.044] N=1,315**	-0.062 [0.045] N=1,181	-0.078 [0.049] N=1,008	-0.039 [0.053] N=825
Below-normal BMI-for-age BMIZ below -2	0.109 P=164	0.006 [0.037] N=1,442	0.013 [0.038] N=1,341	0.036 [0.042] N=1,200	0.043 [0.047] N=1,024	0.046 [0.052] N=844

Specification 1 UNICEF age-interval fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.083 [0.327] N=1,284	0.046 [0.366] N=1,171	0.065 [0.398] N=1,029	0.094 [0.445] N=845	-0.052 [0.602] N=653
WHO weight-for-age z score	0.093 [0.178] N=1,423	0.156 [0.188] N=1,294	0.053 [0.192] N=1,139	0.124 [0.268] N=933	0.005 [0.311] N=721
WHO weight-for-height z score	-0.199 [0.275] N=1,216	-0.193 [0.294] N=1,118	-0.447 [0.345] N=982	-0.570 [0.429] N=806	-0.276 [0.514] N=621
WHO BMI-for-age z score	-0.099 [0.288] N=1,220	-0.047 [0.313] N=1,119	-0.331 [0.362] N=978	-0.507 [0.451] N=802	-0.157 [0.530] N=622
Stunted HAZ below -2	-0.065 [0.060] N=1,284	-0.074 [0.067] N=1,171	-0.074 [0.072] N=1,029	-0.112 [0.083] N=845	-0.017 [0.109] N=653
Underweight WAZ below -2	**-0.060* [0.032] N=1,423**	**-0.072** [0.034] N=1,294**	**-0.079** [0.037] N=1,139**	**-0.096** [0.044] N=933**	-0.095 [0.058] N=721
Wasted WHZ below -2	-0.000 [0.039] N=1,216	0.020 [0.039] N=1,118	0.062 [0.045] N=982	0.064 [0.056] N=806	0.041 [0.061] N=621
Overweight WHZ above +2	-0.028 [0.056] N=1,216	-0.018 [0.060] N=1,118	-0.025 [0.065] N=982	-0.047 [0.080] N=806	0.016 [0.098] N=621
Obese WHZ above +3	-0.013 [0.040] N=1,216	-0.016 [0.043] N=1,118	-0.004 [0.049] N=982	-0.013 [0.061] N=806	0.014 [0.083] N=621
CIAF any stunting underweight or wasting	**-0.102* [0.061] N=1,199**	-0.086 [0.063] N=1,101	-0.035 [0.070] N=967	-0.070 [0.082] N=794	0.034 [0.109] N=611
Below-normal BMI-for-age BMIZ below -2	-0.006 [0.041] N=1,220	0.003 [0.042] N=1,119	0.046 [0.048] N=978	0.059 [0.057] N=802	0.066 [0.067] N=622

Specification 2 UNICEF age intervals and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.509	0.209 [0.221] N=1,517	0.205 [0.235] N=1,404	0.232 [0.244] N=1,262	0.252 [0.255] N=1,078	0.230 [0.291] N=886
WHO weight-for-age z score	0.024	0.103 [0.130] N=1,679	0.140 [0.137] N=1,550	0.079 [0.137] N=1,395	0.096 [0.160] N=1,189	0.037 [0.162] N=977
WHO weight-for-height z score	0.367	-0.149 [0.195] N=1,440	-0.143 [0.201] N=1,342	-0.266 [0.218] N=1,206	-0.289 [0.241] N=1,030	-0.164 [0.256] N=845
WHO BMI-for-age z score	0.368	-0.176 [0.218] N=1,442	-0.157 [0.231] N=1,341	-0.315 [0.248] N=1,200	-0.386 [0.280] N=1,024	-0.273 [0.301] N=844
Stunted HAZ below -2	0.238 P=353	-0.050 [0.040] N=1,517	-0.053 [0.042] N=1,404	-0.050 [0.044] N=1,262	-0.060 [0.047] N=1,078	-0.018 [0.052] N=886
Underweight WAZ below -2	0.064 P=108	-0.036 [0.022] N=1,679	**-0.041* [0.023] N=1,550**	**-0.040* [0.024] N=1,395**	-0.041 [0.026] N=1,189	-0.030 [0.028] N=977
Wasted WHZ below -2	0.106 P=159	-0.017 [0.034] N=1,440	-0.006 [0.035] N=1,342	0.012 [0.038] N=1,206	0.004 [0.042] N=1,030	-0.012 [0.046] N=845
Overweight WHZ above +2	0.196 P=273	-0.021 [0.042] N=1,440	-0.013 [0.043] N=1,342	-0.015 [0.046] N=1,206	-0.023 [0.050] N=1,030	-0.003 [0.056] N=845
Obese WHZ above +3	0.090 P=128	-0.018 [0.029] N=1,440	-0.020 [0.029] N=1,342	-0.016 [0.031] N=1,206	-0.023 [0.034] N=1,030	-0.015 [0.038] N=845
CIAF any stunting underweight or wasting	0.319 P=450	**-0.097** [0.043] N=1,413**	**-0.085** [0.043] N=1,315**	-0.058 [0.044] N=1,181	-0.075 [0.048] N=1,008	-0.036 [0.052] N=825
Below-normal BMI-for-age BMIZ below -2	0.109 P=164	0.007 [0.037] N=1,442	0.015 [0.038] N=1,341	0.038 [0.041] N=1,200	0.043 [0.046] N=1,024	0.046 [0.051] N=844

Specification 2 UNICEF age intervals and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.089 [0.326] N=1,284	0.056 [0.364] N=1,171	0.082 [0.396] N=1,029	0.123 [0.441] N=845	-0.015 [0.598] N=653
WHO weight-for-age z score	0.092 [0.179] N=1,423	0.154 [0.189] N=1,294	0.052 [0.193] N=1,139	0.124 [0.269] N=933	0.003 [0.313] N=721
WHO weight-for-height z score	-0.206 [0.277] N=1,216	-0.207 [0.296] N=1,118	-0.467 [0.349] N=982	-0.584 [0.432] N=806	-0.309 [0.523] N=621
WHO BMI-for-age z score	-0.107 [0.289] N=1,220	-0.062 [0.315] N=1,119	-0.348 [0.364] N=978	-0.515 [0.454] N=802	-0.184 [0.538] N=622
Stunted HAZ below -2	-0.064 [0.060] N=1,284	-0.073 [0.066] N=1,171	-0.075 [0.072] N=1,029	-0.112 [0.083] N=845	-0.020 [0.110] N=653
Underweight WAZ below -2	**-0.060* [0.032] N=1,423**	**-0.072** [0.034] N=1,294**	**-0.078** [0.037] N=1,139**	**-0.096** [0.044] N=933**	-0.094 [0.057] N=721
Wasted WHZ below -2	0.003 [0.040] N=1,216	0.026 [0.040] N=1,118	0.070 [0.047] N=982	0.069 [0.058] N=806	0.051 [0.065] N=621
Overweight WHZ above +2	-0.027 [0.056] N=1,216	-0.016 [0.059] N=1,118	-0.022 [0.065] N=982	-0.045 [0.080] N=806	0.020 [0.097] N=621
Obese WHZ above +3	-0.012 [0.040] N=1,216	-0.015 [0.042] N=1,118	-0.003 [0.048] N=982	-0.013 [0.061] N=806	0.016 [0.082] N=621
CIAF any stunting underweight or wasting	-0.097 [0.061] N=1,199	-0.078 [0.063] N=1,101	-0.026 [0.070] N=967	-0.064 [0.081] N=794	0.045 [0.108] N=611
Below-normal BMI-for-age BMIZ below -2	-0.004 [0.042] N=1,220	0.007 [0.043] N=1,119	0.051 [0.049] N=978	0.062 [0.058] N=802	0.073 [0.071] N=622

Specification 3 UNICEF age intervals female and household income groups No donut hole

Outcome	**Mean**	**8 years**	**7 years**	**6 years**	**5 years**	**4 years**
WHO height-for-age z score	-0.509	0.161 [0.217] N=1,517	0.140 [0.230] N=1,404	0.183 [0.240] N=1,262	0.186 [0.251] N=1,078	0.155 [0.282] N=886
WHO weight-for-age z score	0.024	0.047 [0.128] N=1,679	0.086 [0.136] N=1,550	0.026 [0.139] N=1,395	0.032 [0.161] N=1,189	-0.034 [0.166] N=977
WHO weight-for-height z score	0.367	-0.184 [0.193] N=1,440	-0.171 [0.200] N=1,342	-0.298 [0.216] N=1,206	-0.324 [0.238] N=1,030	-0.207 [0.254] N=845
WHO BMI-for-age z score	0.368	-0.193 [0.217] N=1,442	-0.168 [0.229] N=1,341	-0.330 [0.245] N=1,200	-0.404 [0.275] N=1,024	-0.293 [0.298] N=844
Stunted HAZ below -2	0.238 P=353	-0.039 [0.039] N=1,517	-0.039 [0.041] N=1,404	-0.039 [0.043] N=1,262	-0.047 [0.046] N=1,078	-0.001 [0.050] N=886
Underweight WAZ below -2	0.064 P=108	-0.027 [0.022] N=1,679	-0.029 [0.022] N=1,550	-0.029 [0.023] N=1,395	-0.027 [0.025] N=1,189	-0.013 [0.027] N=977
Wasted WHZ below -2	0.106 P=159	-0.011 [0.034] N=1,440	0.002 [0.034] N=1,342	0.020 [0.037] N=1,206	0.011 [0.041] N=1,030	-0.005 [0.046] N=845
Overweight WHZ above +2	0.196 P=273	-0.022 [0.043] N=1,440	-0.012 [0.044] N=1,342	-0.016 [0.046] N=1,206	-0.026 [0.051] N=1,030	-0.008 [0.057] N=845
Obese WHZ above +3	0.090 P=128	-0.018 [0.028] N=1,440	-0.018 [0.029] N=1,342	-0.016 [0.031] N=1,206	-0.025 [0.034] N=1,030	-0.020 [0.038] N=845
CIAF any stunting underweight or wasting	0.319 P=450	**-0.080* [0.041] N=1,413**	-0.065 [0.040] N=1,315	-0.041 [0.041] N=1,181	-0.057 [0.046] N=1,008	-0.014 [0.050] N=825
Below-normal BMI-for-age BMIZ below -2	0.109 P=164	0.011 [0.037] N=1,442	0.020 [0.038] N=1,341	0.044 [0.041] N=1,200	0.049 [0.045] N=1,024	0.053 [0.050] N=844

Specification 3 UNICEF age intervals female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.057 [0.320] N=1,284	-0.002 [0.356] N=1,171	0.038 [0.392] N=1,029	0.043 [0.438] N=845	-0.127 [0.581] N=653
WHO weight-for-age z score	0.040 [0.174] N=1,423	0.100 [0.185] N=1,294	-0.001 [0.190] N=1,139	0.042 [0.266] N=933	-0.104 [0.318] N=721
WHO weight-for-height z score	-0.229 [0.277] N=1,216	-0.220 [0.299] N=1,118	-0.479 [0.350] N=982	-0.612 [0.435] N=806	-0.357 [0.530] N=621
WHO BMI-for-age z score	-0.112 [0.290] N=1,220	-0.060 [0.316] N=1,119	-0.346 [0.365] N=978	-0.547 [0.459] N=802	-0.208 [0.551] N=622
Stunted HAZ below -2	-0.058 [0.058] N=1,284	-0.061 [0.064] N=1,171	-0.065 [0.070] N=1,029	-0.099 [0.081] N=845	-0.001 [0.105] N=653
Underweight WAZ below -2	**-0.052* [0.031] N=1,423**	**-0.059* [0.033] N=1,294**	**-0.066* [0.036] N=1,139**	**-0.080* [0.042] N=933**	-0.069 [0.055] N=721
Wasted WHZ below -2	0.008 [0.040] N=1,216	0.034 [0.041] N=1,118	0.077 [0.047] N=982	0.080 [0.058] N=806	0.062 [0.067] N=621
Overweight WHZ above +2	-0.024 [0.057] N=1,216	-0.010 [0.060] N=1,118	-0.016 [0.066] N=982	-0.043 [0.082] N=806	0.020 [0.099] N=621
Obese WHZ above +3	-0.011 [0.040] N=1,216	-0.010 [0.042] N=1,118	0.000 [0.048] N=982	-0.014 [0.060] N=806	0.006 [0.081] N=621
CIAF any stunting underweight or wasting	-0.086 [0.059] N=1,199	-0.060 [0.060] N=1,101	-0.012 [0.065] N=967	-0.043 [0.079] N=794	0.074 [0.104] N=611
Below-normal BMI-for-age BMIZ below -2	-0.003 [0.042] N=1,220	0.011 [0.043] N=1,119	0.055 [0.049] N=978	0.073 [0.059] N=802	0.081 [0.073] N=622

Specification 4 UNICEF age intervals female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.509	0.109 [0.216] N=1,517	0.083 [0.230] N=1,404	0.125 [0.243] N=1,262	0.125 [0.253] N=1,078	0.063 [0.284] N=886
WHO weight-for-age z score	0.024	0.049 [0.127] N=1,679	0.081 [0.135] N=1,550	0.023 [0.139] N=1,395	0.021 [0.163] N=1,189	-0.064 [0.164] N=977
WHO weight-for-height z score	0.367	-0.113 [0.193] N=1,440	-0.101 [0.198] N=1,342	-0.237 [0.216] N=1,206	-0.270 [0.242] N=1,030	-0.135 [0.254] N=845
WHO BMI-for-age z score	0.368	-0.132 [0.219] N=1,442	-0.105 [0.231] N=1,341	-0.272 [0.250] N=1,200	-0.353 [0.285] N=1,024	-0.232 [0.309] N=844
Stunted HAZ below -2	0.238 P=353	-0.032 [0.039] N=1,517	-0.032 [0.041] N=1,404	-0.031 [0.043] N=1,262	-0.035 [0.046] N=1,078	0.012 [0.050] N=886
Underweight WAZ below -2	0.064 P=108	-0.026 [0.022] N=1,679	-0.029 [0.023] N=1,550	-0.027 [0.024] N=1,395	-0.025 [0.026] N=1,189	-0.011 [0.029] N=977
Wasted WHZ below -2	0.106 P=159	-0.015 [0.034] N=1,440	-0.003 [0.035] N=1,342	0.017 [0.038] N=1,206	0.009 [0.042] N=1,030	-0.009 [0.047] N=845
Overweight WHZ above +2	0.196 P=273	-0.012 [0.043] N=1,440	-0.004 [0.044] N=1,342	-0.009 [0.046] N=1,206	-0.018 [0.052] N=1,030	0.001 [0.057] N=845
Obese WHZ above +3	0.090 P=128	-0.011 [0.029] N=1,440	-0.012 [0.029] N=1,342	-0.009 [0.031] N=1,206	-0.017 [0.035] N=1,030	-0.009 [0.039] N=845
CIAF any stunting underweight or wasting	0.319 P=450	**-0.076* [0.042] N=1,413**	-0.061 [0.042] N=1,315	-0.036 [0.044] N=1,181	-0.047 [0.047] N=1,008	-0.005 [0.051] N=825
Below-normal BMI-for-age BMIZ below -2	0.109 P=164	0.008 [0.037] N=1,442	0.017 [0.039] N=1,341	0.042 [0.042] N=1,200	0.046 [0.046] N=1,024	0.048 [0.052] N=844

Specification 4 UNICEF age intervals female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.035 [0.314] N=1,284	-0.030 [0.348] N=1,171	0.022 [0.387] N=1,029	0.037 [0.424] N=845	-0.224 [0.561] N=653
WHO weight-for-age z score	0.041 [0.173] N=1,423	0.097 [0.185] N=1,294	-0.005 [0.194] N=1,139	0.045 [0.270] N=933	-0.103 [0.320] N=721
WHO weight-for-height z score	-0.204 [0.277] N=1,216	-0.181 [0.296] N=1,118	-0.455 [0.351] N=982	-0.588 [0.440] N=806	-0.256 [0.530] N=621
WHO BMI-for-age z score	-0.103 [0.290] N=1,220	-0.033 [0.314] N=1,119	-0.332 [0.368] N=978	-0.523 [0.461] N=802	-0.132 [0.552] N=622
Stunted HAZ below -2	-0.053 [0.059] N=1,284	-0.057 [0.065] N=1,171	-0.064 [0.071] N=1,029	-0.097 [0.082] N=845	0.019 [0.107] N=653
Underweight WAZ below -2	-0.049 [0.031] N=1,423	**-0.057* [0.034] N=1,294**	**-0.063* [0.038] N=1,139**	**-0.074* [0.044] N=933**	-0.062 [0.058] N=721
Wasted WHZ below -2	0.010 [0.040] N=1,216	0.032 [0.041] N=1,118	0.076 [0.047] N=982	0.080 [0.058] N=806	0.056 [0.067] N=621
Overweight WHZ above +2	-0.019 [0.057] N=1,216	-0.006 [0.060] N=1,118	-0.014 [0.066] N=982	-0.037 [0.083] N=806	0.038 [0.099] N=621
Obese WHZ above +3	-0.007 [0.040] N=1,216	-0.005 [0.042] N=1,118	0.005 [0.049] N=982	-0.007 [0.060] N=806	0.023 [0.081] N=621
CIAF any stunting underweight or wasting	-0.077 [0.060] N=1,199	-0.056 [0.062] N=1,101	-0.010 [0.068] N=967	-0.038 [0.079] N=794	0.092 [0.105] N=611
Below-normal BMI-for-age BMIZ below -2	0.001 [0.041] N=1,220	0.010 [0.043] N=1,119	0.054 [0.050] N=978	0.070 [0.058] N=802	0.075 [0.071] N=622

Specification 5 UNICEF age intervals female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.515	0.165 [0.228] N=1,430	0.134 [0.242] N=1,329	0.171 [0.245] N=1,192	0.233 [0.249] N=1,021	0.223 [0.287] N=835
WHO weight-for-age z score	0.017	0.100 [0.135] N=1,581	0.133 [0.141] N=1,465	0.070 [0.145] N=1,317	0.085 [0.168] N=1,126	0.049 [0.178] N=922
WHO weight-for-height z score	0.369	-0.072 [0.200] N=1,359	-0.065 [0.204] N=1,273	-0.188 [0.223] N=1,143	-0.221 [0.246] N=979	-0.082 [0.268] N=800
WHO BMI-for-age z score	0.376	-0.086 [0.220] N=1,358	-0.066 [0.229] N=1,269	-0.211 [0.247] N=1,135	-0.287 [0.278] N=971	-0.165 [0.303] N=797
Stunted HAZ below -2	0.239 P=332	-0.035 [0.043] N=1,430	-0.036 [0.045] N=1,329	-0.034 [0.046] N=1,192	-0.056 [0.049] N=1,021	-0.012 [0.054] N=835
Underweight WAZ below -2	0.065 P=104	-0.031 [0.023] N=1,581	-0.033 [0.024] N=1,465	-0.032 [0.025] N=1,317	-0.034 [0.027] N=1,126	-0.027 [0.032] N=922
Wasted WHZ below -2	0.107 P=152	-0.015 [0.036] N=1,359	-0.007 [0.036] N=1,273	0.014 [0.039] N=1,143	0.004 [0.044] N=979	-0.009 [0.049] N=800
Overweight WHZ above +2	0.197 P=257	-0.011 [0.044] N=1,359	-0.006 [0.045] N=1,273	-0.009 [0.048] N=1,143	-0.021 [0.053] N=979	0.000 [0.060] N=800
Obese WHZ above +3	0.093 P=125	-0.017 [0.030] N=1,359	-0.018 [0.031] N=1,273	-0.013 [0.033] N=1,143	-0.022 [0.035] N=979	-0.013 [0.040] N=800
CIAF any stunting underweight or wasting	0.322 P=427	-0.077 [0.047] N=1,334	-0.068 [0.047] N=1,248	-0.041 [0.049] N=1,119	-0.071 [0.054] N=958	-0.025 [0.060] N=781
Below-normal BMI-for-age BMIZ below -2	0.109 P=154	0.007 [0.037] N=1,358	0.011 [0.037] N=1,269	0.034 [0.040] N=1,135	0.036 [0.046] N=971	0.040 [0.050] N=797

Specification 5 UNICEF age intervals female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.028 [0.343] N=1,212	-0.058 [0.381] N=1,111	-0.031 [0.411] N=974	0.120 [0.447] N=803	0.004 [0.626] N=617
WHO weight-for-age z score	0.067 [0.183] N=1,341	0.116 [0.187] N=1,225	-0.010 [0.198] N=1,077	0.055 [0.273] N=886	-0.039 [0.330] N=682
WHO weight-for-height z score	-0.129 [0.283] N=1,147	-0.133 [0.297] N=1,061	-0.385 [0.350] N=931	-0.499 [0.429] N=767	-0.149 [0.521] N=588
WHO BMI-for-age z score	-0.051 [0.295] N=1,150	-0.012 [0.316] N=1,061	-0.269 [0.368] N=927	-0.434 [0.449] N=763	-0.055 [0.539] N=589
Stunted HAZ below -2	-0.042 [0.063] N=1,212	-0.046 [0.069] N=1,111	-0.049 [0.076] N=974	-0.110 [0.084] N=803	-0.016 [0.114] N=617
Underweight WAZ below -2	-0.053 [0.032] N=1,341	**-0.061* [0.034] N=1,225**	-0.060 [0.038] N=1,077	**-0.076* [0.044] N=886**	-0.076 [0.062] N=682
Wasted WHZ below -2	0.002 [0.040] N=1,147	0.019 [0.040] N=1,061	0.067 [0.046] N=931	0.059 [0.058] N=767	0.033 [0.065] N=588
Overweight WHZ above +2	-0.016 [0.058] N=1,147	-0.013 [0.059] N=1,061	-0.011 [0.066] N=931	-0.043 [0.081] N=767	0.038 [0.102] N=588
Obese WHZ above +3	-0.009 [0.043] N=1,147	-0.010 [0.045] N=1,061	0.003 [0.051] N=931	-0.010 [0.062] N=767	0.036 [0.086] N=588
CIAF any stunting underweight or wasting	-0.070 [0.064] N=1,132	-0.056 [0.065] N=1,046	0.000 [0.073] N=917	-0.063 [0.083] N=756	0.044 [0.116] N=579
Below-normal BMI-for-age BMIZ below -2	0.001 [0.042] N=1,150	0.006 [0.042] N=1,061	0.051 [0.049] N=927	0.055 [0.058] N=763	0.061 [0.071] N=589

Specification 6 UNICEF age intervals female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.515	0.162 [0.225] N=1,430	0.134 [0.237] N=1,329	0.134 [0.244] N=1,192	0.180 [0.252] N=1,021	0.141 [0.290] N=835
WHO weight-for-age z score	0.017	0.102 [0.136] N=1,581	0.135 [0.142] N=1,465	0.065 [0.144] N=1,317	0.086 [0.166] N=1,126	0.042 [0.175] N=922
WHO weight-for-height z score	0.369	-0.072 [0.200] N=1,359	-0.064 [0.204] N=1,273	-0.184 [0.221] N=1,143	-0.205 [0.246] N=979	-0.062 [0.265] N=800
WHO BMI-for-age z score	0.376	-0.089 [0.219] N=1,358	-0.069 [0.229] N=1,269	-0.210 [0.246] N=1,135	-0.270 [0.279] N=971	-0.143 [0.304] N=797
Stunted HAZ below -2	0.239 P=332	-0.036 [0.043] N=1,430	-0.039 [0.045] N=1,329	-0.033 [0.047] N=1,192	-0.052 [0.050] N=1,021	-0.009 [0.055] N=835
Underweight WAZ below -2	0.065 P=104	-0.034 [0.023] N=1,581	-0.037 [0.024] N=1,465	-0.035 [0.025] N=1,317	-0.037 [0.027] N=1,126	-0.029 [0.031] N=922
Wasted WHZ below -2	0.107 P=152	-0.015 [0.036] N=1,359	-0.007 [0.036] N=1,273	0.013 [0.039] N=1,143	0.002 [0.045] N=979	-0.016 [0.050] N=800
Overweight WHZ above +2	0.197 P=257	-0.011 [0.044] N=1,359	-0.006 [0.045] N=1,273	-0.006 [0.048] N=1,143	-0.017 [0.053] N=979	0.006 [0.059] N=800
Obese WHZ above +3	0.093 P=125	-0.016 [0.030] N=1,359	-0.017 [0.030] N=1,273	-0.012 [0.032] N=1,143	-0.020 [0.035] N=979	-0.008 [0.039] N=800
CIAF any stunting underweight or wasting	0.322 P=427	**-0.078*** **[0.047]** **N=1,334**	-0.070 [0.047] N=1,248	-0.041 [0.049] N=1,119	-0.071 [0.054] N=958	-0.034 [0.060] N=781
Below-normal BMI-for-age BMIZ below -2	0.109 P=154	0.006 [0.038] N=1,358	0.011 [0.038] N=1,269	0.033 [0.041] N=1,135	0.034 [0.046] N=971	0.034 [0.052] N=797

Specification 6 UNICEF age intervals female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.095 [0.343] N=1,212	0.027 [0.379] N=1,111	0.008 [0.411] N=974	0.134 [0.462] N=803	-0.027 [0.648] N=617
WHO weight-for-age z score	0.088 [0.184] N=1,341	0.141 [0.191] N=1,225	0.022 [0.197] N=1,077	0.102 [0.268] N=886	0.018 [0.323] N=682
WHO weight-for-height z score	-0.123 [0.282] N=1,147	-0.119 [0.296] N=1,061	-0.371 [0.349] N=931	-0.469 [0.434] N=767	-0.111 [0.529] N=588
WHO BMI-for-age z score	-0.049 [0.295] N=1,150	-0.005 [0.316] N=1,061	-0.266 [0.366] N=927	-0.410 [0.456] N=763	-0.023 [0.551] N=589
Stunted HAZ below -2	-0.052 [0.063] N=1,212	-0.058 [0.069] N=1,111	-0.056 [0.075] N=974	-0.118 [0.085] N=803	-0.024 [0.114] N=617
Underweight WAZ below -2	**-0.057* [0.033] N=1,341**	**-0.066* [0.035] N=1,225**	**-0.070* [0.038] N=1,077**	**-0.087** [0.044] N=886**	-0.092 [0.060] N=682
Wasted WHZ below -2	0.004 [0.040] N=1,147	0.021 [0.039] N=1,061	0.068 [0.046] N=931	0.058 [0.058] N=767	0.027 [0.065] N=588
Overweight WHZ above +2	-0.013 [0.058] N=1,147	-0.007 [0.060] N=1,061	-0.006 [0.067] N=931	-0.034 [0.082] N=767	0.047 [0.103] N=588
Obese WHZ above +3	-0.009 [0.043] N=1,147	-0.011 [0.045] N=1,061	0.005 [0.052] N=931	-0.007 [0.063] N=767	0.040 [0.087] N=588
CIAF any stunting underweight or wasting	-0.071 [0.063] N=1,132	-0.058 [0.065] N=1,046	-0.003 [0.073] N=917	-0.070 [0.084] N=756	0.028 [0.114] N=579
Below-normal BMI-for-age BMIZ below -2	0.001 [0.042] N=1,150	0.007 [0.042] N=1,061	0.050 [0.049] N=927	0.051 [0.058] N=763	0.051 [0.072] N=589

Mother RDD secondary analysis

Specification 1 UNICEF age-interval fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.464	0.017 [0.241] N=1,410	0.012 [0.257] N=1,274	0.054 [0.283] N=1,115	0.001 [0.309] N=943	-0.026 [0.326] N=766
WHO weight-for-age z score	0.029	0.060 [0.137] N=1,550	0.146 [0.145] N=1,410	-0.009 [0.144] N=1,228	-0.025 [0.155] N=1,034	0.018 [0.166] N=830
WHO weight-for-height z score	0.385	0.063 [0.194] N=1,342	0.106 [0.197] N=1,214	0.048 [0.208] N=1,066	0.015 [0.227] N=899	-0.042 [0.254] N=728
WHO BMI-for-age z score	0.352	-0.053 [0.216] N=1,339	-0.027 [0.231] N=1,213	-0.165 [0.249] N=1,065	-0.177 [0.268] N=897	-0.151 [0.294] N=731
Stunted HAZ below -2	0.236 P=334	0.031 [0.046] N=1,410	0.035 [0.048] N=1,274	0.034 [0.050] N=1,115	0.055 [0.055] N=943	0.060 [0.058] N=766
Underweight WAZ below -2	0.062 P=98	-0.020 [0.026] N=1,550	-0.035 [0.027] N=1,410	-0.014 [0.029] N=1,228	-0.018 [0.031] N=1,034	-0.022 [0.034] N=830
Wasted WHZ below -2	0.105 P=142	**0.055* [0.031] N=1,342**	0.042 [0.033] N=1,214	**0.073** [0.035] N=1,066**	0.050 [0.035] N=899	0.060 [0.038] N=728
Overweight WHZ above +2	0.193 P=262	0.043 [0.041] N=1,342	0.050 [0.042] N=1,214	0.056 [0.046] N=1,066	0.049 [0.050] N=899	0.047 [0.054] N=728
Obese WHZ above +3	0.096 P=133	0.020 [0.030] N=1,342	0.028 [0.030] N=1,214	0.017 [0.033] N=1,066	0.022 [0.038] N=899	0.004 [0.040] N=728
CIAF any stunting underweight or wasting	0.318 P=421	**0.099* [0.053] N=1,316**	0.076 [0.057] N=1,188	**0.118* [0.059] N=1,041**	**0.113* [0.065] N=877**	**0.149** [0.067] N=711**
Below-normal BMI-for-age BMIZ below -2	0.107 P=145	0.050 [0.037] N=1,339	0.041 [0.041] N=1,213	0.070 [0.044] N=1,065	0.059 [0.046] N=897	0.063 [0.050] N=731

Specification 1 UNICEF age-interval fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.281 [0.303] N=1,219	0.334 [0.330] N=1,083	0.511 [0.386] N=924	0.485 [0.486] N=752	0.631 [0.552] N=575
WHO weight-for-age z score	0.130 [0.205] N=1,345	0.301 [0.215] N=1,205	0.043 [0.231] N=1,023	0.022 [0.272] N=829	0.206 [0.328] N=625
WHO weight-for-height z score	-0.111 [0.287] N=1,167	-0.061 [0.295] N=1,039	-0.177 [0.328] N=891	-0.280 [0.402] N=724	-0.585 [0.532] N=553
WHO BMI-for-age z score	-0.319 [0.281] N=1,165	-0.318 [0.309] N=1,039	**-0.635* [0.345] N=891**	**-0.764* [0.422] N=723**	**-0.913* [0.532] N=557**
Stunted HAZ below -2	-0.005 [0.067] N=1,219	-0.007 [0.074] N=1,083	-0.024 [0.081] N=924	0.008 [0.103] N=752	-0.009 [0.121] N=575
Underweight WAZ below -2	-0.035 [0.036] N=1,345	-0.062 [0.039] N=1,205	-0.030 [0.045] N=1,023	-0.049 [0.052] N=829	-0.086 [0.058] N=625
Wasted WHZ below -2	**0.095** [0.040] N=1,167**	**0.077* [0.046] N=1,039**	**0.144*** [0.050] N=891**	**0.108** [0.050] N=724**	**0.152*** [0.057] N=553**
Overweight WHZ above +2	0.029 [0.062] N=1,167	0.034 [0.065] N=1,039	0.048 [0.076] N=891	0.022 [0.093] N=724	-0.009 [0.118] N=553
Obese WHZ above +3	0.006 [0.046] N=1,167	0.016 [0.048] N=1,039	-0.003 [0.056] N=891	0.005 [0.073] N=724	-0.074 [0.087] N=553
CIAF any stunting underweight or wasting	0.078 [0.072] N=1,145	0.029 [0.080] N=1,017	0.095 [0.090] N=870	0.071 [0.107] N=706	0.124 [0.121] N=540
Below-normal BMI-for-age BMIZ below -2	**0.100*** [0.036] N=1,165**	**0.094** [0.045] N=1,039**	**0.161*** [0.050] N=891**	**0.154*** [0.050] N=723**	**0.190*** [0.050] N=557**

Specification 2 UNICEF age intervals and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.464	0.009 [0.240] N=1,410	0.002 [0.256] N=1,274	0.045 [0.282] N=1,115	-0.011 [0.307] N=943	-0.017 [0.327] N=766
WHO weight-for-age z score	0.029	0.061 [0.137] N=1,550	0.146 [0.144] N=1,410	-0.008 [0.143] N=1,228	-0.024 [0.155] N=1,034	0.016 [0.167] N=830
WHO weight-for-height z score	0.385	0.065 [0.194] N=1,342	0.109 [0.197] N=1,214	0.052 [0.208] N=1,066	0.016 [0.228] N=899	-0.060 [0.257] N=728
WHO BMI-for-age z score	0.352	-0.051 [0.216] N=1,339	-0.025 [0.231] N=1,213	-0.162 [0.249] N=1,065	-0.176 [0.268] N=897	-0.170 [0.297] N=731
Stunted HAZ below -2	0.236 P=334	0.031 [0.046] N=1,410	0.035 [0.048] N=1,274	0.033 [0.050] N=1,115	0.055 [0.055] N=943	0.060 [0.058] N=766
Underweight WAZ below -2	0.062 P=98	-0.021 [0.026] N=1,550	-0.035 [0.027] N=1,410	-0.015 [0.029] N=1,228	-0.018 [0.031] N=1,034	-0.020 [0.034] N=830
Wasted WHZ below -2	0.105 P=142	**0.054* [0.031] N=1,342**	0.041 [0.033] N=1,214	**0.071** [0.035] N=1,066**	0.050 [0.035] N=899	**0.066* [0.038] N=728**
Overweight WHZ above +2	0.193 P=262	0.043 [0.041] N=1,342	0.049 [0.042] N=1,214	0.056 [0.046] N=1,066	0.049 [0.050] N=899	0.048 [0.055] N=728
Obese WHZ above +3	0.096 P=133	0.019 [0.030] N=1,342	0.027 [0.030] N=1,214	0.016 [0.033] N=1,066	0.022 [0.038] N=899	0.007 [0.040] N=728
CIAF any stunting underweight or wasting	0.318 P=421	**0.098* [0.054] N=1,316**	0.074 [0.058] N=1,188	**0.115* [0.061] N=1,041**	**0.112* [0.066] N=877**	**0.155** [0.068] N=711**
Below-normal BMI-for-age BMIZ below -2	0.107 P=145	0.049 [0.036] N=1,339	0.041 [0.040] N=1,213	0.069 [0.043] N=1,065	0.059 [0.045] N=897	0.068 [0.050] N=731

Specification 2 UNICEF age intervals and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.265 [0.299] N=1,219	0.315 [0.326] N=1,083	0.501 [0.382] N=924	0.445 [0.475] N=752	0.608 [0.544] N=575
WHO weight-for-age z score	0.132 [0.204] N=1,345	0.303 [0.214] N=1,205	0.045 [0.228] N=1,023	0.026 [0.270] N=829	0.207 [0.332] N=625
WHO weight-for-height z score	-0.097 [0.288] N=1,167	-0.044 [0.295] N=1,039	-0.162 [0.327] N=891	-0.253 [0.401] N=724	-0.565 [0.540] N=553
WHO BMI-for-age z score	-0.303 [0.283] N=1,165	-0.299 [0.311] N=1,039	**-0.620* [0.345] N=891**	**-0.736* [0.424] N=723**	-0.902 [0.545] N=557
Stunted HAZ below -2	-0.005 [0.067] N=1,219	-0.007 [0.074] N=1,083	-0.028 [0.082] N=924	0.007 [0.103] N=752	-0.009 [0.122] N=575
Underweight WAZ below -2	-0.037 [0.036] N=1,345	-0.063 [0.039] N=1,205	-0.031 [0.044] N=1,023	-0.051 [0.052] N=829	-0.086 [0.059] N=625
Wasted WHZ below -2	**0.089** [0.040] N=1,167**	0.068 [0.046] N=1,039	**0.132** [0.051] N=891**	**0.096* [0.051] N=724**	**0.144** [0.057] N=553**
Overweight WHZ above +2	0.028 [0.062] N=1,167	0.033 [0.065] N=1,039	0.044 [0.075] N=891	0.020 [0.093] N=724	-0.010 [0.118] N=553
Obese WHZ above +3	0.003 [0.046] N=1,167	0.011 [0.048] N=1,039	-0.011 [0.056] N=891	-0.002 [0.072] N=724	-0.078 [0.086] N=553
CIAF any stunting underweight or wasting	0.072 [0.073] N=1,145	0.017 [0.083] N=1,017	0.078 [0.093] N=870	0.054 [0.110] N=706	0.115 [0.126] N=540
Below-normal BMI-for-age BMIZ below -2	**0.095** [0.037] N=1,165**	**0.087* [0.045] N=1,039**	**0.153*** [0.050] N=891**	**0.145*** [0.050] N=723**	**0.187*** [0.051] N=557**

Specification 3 UNICEF age intervals female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.464	-0.016 [0.234] N=1,410	-0.012 [0.250] N=1,274	0.028 [0.278] N=1,115	-0.014 [0.301] N=943	-0.045 [0.311] N=766
WHO weight-for-age z score	0.029	0.047 [0.137] N=1,550	0.139 [0.145] N=1,410	-0.015 [0.144] N=1,228	-0.035 [0.155] N=1,034	0.006 [0.165] N=830
WHO weight-for-height z score	0.385	0.062 [0.198] N=1,342	0.108 [0.202] N=1,214	0.065 [0.216] N=1,066	0.027 [0.235] N=899	-0.036 [0.265] N=728
WHO BMI-for-age z score	0.352	-0.048 [0.217] N=1,339	-0.020 [0.232] N=1,213	-0.143 [0.253] N=1,065	-0.166 [0.273] N=897	-0.150 [0.301] N=731
Stunted HAZ below -2	0.236 P=334	0.037 [0.045] N=1,410	0.038 [0.047] N=1,274	0.038 [0.049] N=1,115	0.056 [0.053] N=943	0.064 [0.055] N=766
Underweight WAZ below -2	0.062 P=98	-0.018 [0.026] N=1,550	-0.033 [0.028] N=1,410	-0.013 [0.029] N=1,228	-0.017 [0.031] N=1,034	-0.018 [0.033] N=830
Wasted WHZ below -2	0.105 P=142	**0.053* [0.031] N=1,342**	0.039 [0.033] N=1,214	**0.068* [0.035] N=1,066**	0.046 [0.035] N=899	0.058 [0.036] N=728
Overweight WHZ above +2	0.193 P=262	0.044 [0.042] N=1,342	0.049 [0.043] N=1,214	0.058 [0.046] N=1,066	0.048 [0.050] N=899	0.047 [0.054] N=728
Obese WHZ above +3	0.096 P=133	0.021 [0.030] N=1,342	0.028 [0.031] N=1,214	0.019 [0.034] N=1,066	0.024 [0.038] N=899	0.009 [0.041] N=728
CIAF any stunting underweight or wasting	0.318 P=421	**0.102* [0.054] N=1,316**	0.075 [0.058] N=1,188	**0.116* [0.061] N=1,041**	0.108 [0.066] N=877	**0.148** [0.067] N=711**
Below-normal BMI-for-age BMIZ below -2	0.107 P=145	0.049 [0.036] N=1,339	0.039 [0.040] N=1,213	0.067 [0.043] N=1,065	0.057 [0.045] N=897	0.065 [0.049] N=731

Specification 3 UNICEF age intervals female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.295 [0.298] N=1,219	0.361 [0.325] N=1,083	0.569 [0.383] N=924	0.561 [0.478] N=752	0.795 [0.528] N=575
WHO weight-for-age z score	0.129 [0.202] N=1,345	0.315 [0.214] N=1,205	0.061 [0.231] N=1,023	0.027 [0.277] N=829	0.184 [0.348] N=625
WHO weight-for-height z score	-0.133 [0.292] N=1,167	-0.070 [0.303] N=1,039	-0.192 [0.338] N=891	-0.327 [0.421] N=724	-0.704 [0.555] N=553
WHO BMI-for-age z score	-0.324 [0.285] N=1,165	-0.313 [0.315] N=1,039	**-0.647* [0.353] N=891**	**-0.821* [0.440] N=723**	**-1.054* [0.552] N=557**
Stunted HAZ below -2	-0.011 [0.068] N=1,219	-0.016 [0.075] N=1,083	-0.038 [0.084] N=924	-0.015 [0.105] N=752	-0.047 [0.122] N=575
Underweight WAZ below -2	-0.040 [0.036] N=1,345	**-0.070* [0.040] N=1,205**	-0.041 [0.045] N=1,023	-0.061 [0.053] N=829	-0.093 [0.060] N=625
Wasted WHZ below -2	**0.087** [0.041] N=1,167**	0.065 [0.047] N=1,039	**0.130** [0.052] N=891**	**0.091* [0.052] N=724**	**0.139** [0.058] N=553**
Overweight WHZ above +2	0.024 [0.064] N=1,167	0.030 [0.067] N=1,039	0.040 [0.079] N=891	0.004 [0.098] N=724	-0.040 [0.123] N=553
Obese WHZ above +3	-0.002 [0.046] N=1,167	0.006 [0.048] N=1,039	-0.014 [0.057] N=891	-0.010 [0.073] N=724	-0.094 [0.085] N=553
CIAF any stunting underweight or wasting	0.067 [0.075] N=1,145	0.010 [0.085] N=1,017	0.071 [0.097] N=870	0.037 [0.115] N=706	0.085 [0.133] N=540
Below-normal BMI-for-age BMIZ below -2	**0.092** [0.037] N=1,165**	**0.083* [0.046] N=1,039**	**0.150*** [0.050] N=891**	**0.140*** [0.051] N=723**	**0.183*** [0.050] N=557**

Specification 4 UNICEF age intervals female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.464	-0.002 [0.231] N=1,410	0.000 [0.247] N=1,274	0.042 [0.276] N=1,115	0.003 [0.298] N=943	-0.016 [0.310] N=766
WHO weight-for-age z score	0.029	0.060 [0.136] N=1,550	0.152 [0.144] N=1,410	-0.002 [0.143] N=1,228	-0.011 [0.154] N=1,034	0.015 [0.163] N=830
WHO weight-for-height z score	0.385	0.068 [0.193] N=1,342	0.118 [0.194] N=1,214	0.068 [0.205] N=1,066	0.025 [0.223] N=899	-0.059 [0.255] N=728
WHO BMI-for-age z score	0.352	-0.041 [0.211] N=1,339	-0.010 [0.225] N=1,213	-0.144 [0.243] N=1,065	-0.164 [0.259] N=897	-0.156 [0.287] N=731
Stunted HAZ below -2	0.236 P=334	0.036 [0.045] N=1,410	0.037 [0.047] N=1,274	0.036 [0.050] N=1,115	0.054 [0.054] N=943	0.059 [0.056] N=766
Underweight WAZ below -2	0.062 P=98	-0.020 [0.026] N=1,550	-0.035 [0.027] N=1,410	-0.015 [0.029] N=1,228	-0.020 [0.031] N=1,034	-0.021 [0.034] N=830
Wasted WHZ below -2	0.105 P=142	**0.054* [0.030] N=1,342**	0.040 [0.033] N=1,214	**0.071** [0.034] N=1,066**	0.051 [0.034] N=899	**0.063* [0.037] N=728**
Overweight WHZ above +2	0.193 P=262	0.046 [0.041] N=1,342	0.052 [0.042] N=1,214	0.060 [0.046] N=1,066	0.053 [0.050] N=899	0.047 [0.054] N=728
Obese WHZ above +3	0.096 P=133	0.022 [0.030] N=1,342	0.030 [0.030] N=1,214	0.020 [0.033] N=1,066	0.026 [0.037] N=899	0.009 [0.040] N=728
CIAF any stunting underweight or wasting	0.318 P=421	**0.103* [0.054] N=1,316**	0.076 [0.059] N=1,188	**0.117* [0.062] N=1,041**	**0.112* [0.067] N=877**	**0.150** [0.068] N=711**
Below-normal BMI-for-age BMIZ below -2	0.107 P=145	0.048 [0.036] N=1,339	0.039 [0.040] N=1,213	0.068 [0.043] N=1,065	0.059 [0.045] N=897	0.062 [0.048] N=731

Specification 4 UNICEF age intervals female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.277 [0.290] N=1,219	0.341 [0.317] N=1,083	0.531 [0.380] N=924	0.532 [0.470] N=752	0.713 [0.511] N=575
WHO weight-for-age z score	0.138 [0.196] N=1,345	0.317 [0.208] N=1,205	0.068 [0.220] N=1,023	0.069 [0.259] N=829	0.200 [0.305] N=625
WHO weight-for-height z score	-0.112 [0.285] N=1,167	-0.053 [0.289] N=1,039	-0.170 [0.316] N=891	-0.285 [0.391] N=724	-0.645 [0.525] N=553
WHO BMI-for-age z score	-0.314 [0.277] N=1,165	-0.304 [0.304] N=1,039	**-0.629* [0.334] N=891**	**-0.777* [0.412] N=723**	**-0.955* [0.521] N=557**
Stunted HAZ below -2	-0.005 [0.067] N=1,219	-0.012 [0.074] N=1,083	-0.034 [0.084] N=924	-0.012 [0.103] N=752	-0.032 [0.125] N=575
Underweight WAZ below -2	-0.036 [0.035] N=1,345	**-0.065* [0.039] N=1,205**	-0.036 [0.044] N=1,023	-0.059 [0.051] N=829	-0.089 [0.058] N=625
Wasted WHZ below -2	**0.090** [0.040] N=1,167**	0.067 [0.047] N=1,039	**0.132** [0.051] N=891**	**0.095* [0.051] N=724**	**0.139** [0.056] N=553**
Overweight WHZ above +2	0.027 [0.062] N=1,167	0.031 [0.064] N=1,039	0.043 [0.075] N=891	0.014 [0.092] N=724	-0.028 [0.117] N=553
Obese WHZ above +3	0.002 [0.046] N=1,167	0.010 [0.048] N=1,039	-0.012 [0.056] N=891	-0.006 [0.072] N=724	-0.087 [0.085] N=553
CIAF any stunting underweight or wasting	0.076 [0.076] N=1,145	0.016 [0.086] N=1,017	0.074 [0.098] N=870	0.043 [0.114] N=706	0.094 [0.132] N=540
Below-normal BMI-for-age BMIZ below -2	**0.095** [0.037] N=1,165**	**0.086* [0.046] N=1,039**	**0.153*** [0.050] N=891**	**0.145*** [0.051] N=723**	**0.179*** [0.050] N=557**

Specification 5 UNICEF age intervals female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.478	-0.026 [0.241] N=1,332	-0.034 [0.259] N=1,200	-0.001 [0.287] N=1,050	-0.045 [0.301] N=887	-0.074 [0.322] N=722
WHO weight-for-age z score	0.018	0.105 [0.142] N=1,464	0.203 [0.148] N=1,328	0.025 [0.148] N=1,156	-0.001 [0.160] N=973	0.001 [0.170] N=783
WHO weight-for-height z score	0.384	0.107 [0.207] N=1,269	0.165 [0.211] N=1,145	0.089 [0.225] N=1,005	0.022 [0.249] N=848	-0.094 [0.282] N=688
WHO BMI-for-age z score	0.353	-0.002 [0.224] N=1,263	0.038 [0.241] N=1,141	-0.119 [0.259] N=1,001	-0.167 [0.277] N=843	-0.186 [0.311] N=689
Stunted HAZ below -2	0.238 P=317	0.036 [0.046] N=1,332	0.043 [0.048] N=1,200	0.038 [0.050] N=1,050	0.060 [0.053] N=887	0.075 [0.058] N=722
Underweight WAZ below -2	0.063 P=95	-0.025 [0.027] N=1,464	-0.042 [0.028] N=1,328	-0.022 [0.030] N=1,156	-0.021 [0.033] N=973	-0.015 [0.035] N=783
Wasted WHZ below -2	0.107 P=137	**0.059* [0.032] N=1,269**	0.042 [0.035] N=1,145	**0.072** [0.036] N=1,005**	0.060 [0.037] N=848	**0.082** [0.039] N=688**
Overweight WHZ above +2	0.194 P=247	0.049 [0.044] N=1,269	0.057 [0.045] N=1,145	0.059 [0.049] N=1,005	0.050 [0.054] N=848	0.051 [0.060] N=688
Obese WHZ above +3	0.100 P=130	0.020 [0.032] N=1,269	0.028 [0.032] N=1,145	0.016 [0.035] N=1,005	0.024 [0.041] N=848	0.013 [0.044] N=688
CIAF any stunting underweight or wasting	0.324 P=403	**0.104* [0.056] N=1,245**	0.079 [0.061] N=1,121	**0.117* [0.064] N=982**	**0.118* [0.068] N=827**	**0.178** [0.071] N=672**
Below-normal BMI-for-age BMIZ below -2	0.107 P=137	0.055 [0.038] N=1,263	0.045 [0.042] N=1,141	0.073 [0.045] N=1,001	0.076 [0.047] N=843	**0.090* [0.052] N=689**

Specification 5 UNICEF age intervals female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.282 [0.318] N=1,152	0.348 [0.353] N=1,020	0.535 [0.426] N=870	0.530 [0.490] N=707	0.600 [0.572] N=542
WHO weight-for-age z score	0.204 [0.210] N=1,271	**0.403* [0.218] N=1,135**	0.108 [0.233] N=963	0.088 [0.277] N=780	0.204 [0.345] N=590
WHO weight-for-height z score	0.020 [0.299] N=1,102	0.098 [0.308] N=978	-0.043 [0.345] N=838	-0.183 [0.417] N=681	-0.509 [0.588] N=521
WHO BMI-for-age z score	-0.200 [0.293] N=1,098	-0.172 [0.327] N=976	-0.527 [0.365] N=836	-0.695 [0.434] N=678	-0.848 [0.588] N=524
Stunted HAZ below -2	-0.015 [0.070] N=1,152	-0.017 [0.078] N=1,020	-0.051 [0.087] N=870	-0.027 [0.104] N=707	-0.023 [0.125] N=542
Underweight WAZ below -2	-0.046 [0.037] N=1,271	**-0.077* [0.040] N=1,135**	-0.050 [0.044] N=963	-0.066 [0.053] N=780	-0.089 [0.054] N=590
Wasted WHZ below -2	**0.089** [0.042] N=1,102**	0.062 [0.048] N=978	**0.121** [0.052] N=838**	**0.102* [0.054] N=681**	**0.154** [0.063] N=521**
Overweight WHZ above +2	0.044 [0.064] N=1,102	0.049 [0.067] N=978	0.052 [0.078] N=838	0.020 [0.097] N=681	0.002 [0.128] N=521
Obese WHZ above +3	0.002 [0.048] N=1,102	0.009 [0.050] N=978	-0.018 [0.058] N=838	-0.008 [0.075] N=681	-0.068 [0.091] N=521
CIAF any stunting underweight or wasting	0.066 [0.076] N=1,082	0.007 [0.086] N=958	0.055 [0.099] N=819	0.026 [0.115] N=664	0.111 [0.132] N=509
Below-normal BMI-for-age BMIZ below -2	**0.095** [0.038] N=1,098**	**0.082* [0.047] N=976**	**0.143*** [0.052] N=836**	**0.161*** [0.054] N=678**	**0.203*** [0.057] N=524**

Specification 6 UNICEF age intervals female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.478	-0.031 [0.245] N=1,332	-0.036 [0.263] N=1,200	0.006 [0.289] N=1,050	-0.053 [0.310] N=887	-0.097 [0.327] N=722
WHO weight-for-age z score	0.018	0.106 [0.143] N=1,464	0.204 [0.151] N=1,328	0.024 [0.151] N=1,156	0.003 [0.163] N=973	0.003 [0.174] N=783
WHO weight-for-height z score	0.384	0.113 [0.204] N=1,269	0.167 [0.209] N=1,145	0.086 [0.222] N=1,005	0.035 [0.244] N=848	-0.084 [0.278] N=688
WHO BMI-for-age z score	0.353	0.004 [0.222] N=1,263	0.041 [0.240] N=1,141	-0.121 [0.258] N=1,001	-0.158 [0.276] N=843	-0.174 [0.309] N=689
Stunted HAZ below -2	0.238 P=317	0.036 [0.046] N=1,332	0.043 [0.049] N=1,200	0.036 [0.051] N=1,050	0.061 [0.055] N=887	0.077 [0.059] N=722
Underweight WAZ below -2	0.063 P=95	-0.025 [0.026] N=1,464	-0.041 [0.028] N=1,328	-0.020 [0.030] N=1,156	-0.020 [0.032] N=973	-0.014 [0.035] N=783
Wasted WHZ below -2	0.107 P=137	**0.057* [0.032] N=1,269**	0.042 [0.035] N=1,145	**0.073** [0.036] N=1,005**	0.057 [0.037] N=848	**0.079* [0.040] N=688**
Overweight WHZ above +2	0.194 P=247	0.049 [0.044] N=1,269	0.055 [0.045] N=1,145	0.056 [0.048] N=1,005	0.049 [0.053] N=848	0.049 [0.059] N=688
Obese WHZ above +3	0.100 P=130	0.020 [0.031] N=1,269	0.028 [0.032] N=1,145	0.015 [0.035] N=1,005	0.024 [0.040] N=848	0.012 [0.043] N=688
CIAF any stunting underweight or wasting	0.324 P=403	**0.101* [0.056] N=1,245**	0.078 [0.061] N=1,121	**0.116* [0.064] N=982**	**0.118* [0.069] N=827**	**0.174** [0.072] N=672**
Below-normal BMI-for-age BMIZ below -2	0.107 P=137	0.055 [0.038] N=1,263	0.044 [0.042] N=1,141	**0.075* [0.045] N=1,001**	0.074 [0.047] N=843	**0.088* [0.052] N=689**

Specification 6 UNICEF age intervals female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.265 [0.324] N=1,152	0.331 [0.357] N=1,020	0.537 [0.424] N=870	0.481 [0.515] N=707	0.521 [0.586] N=542
WHO weight-for-age z score	0.205 [0.210] N=1,271	**0.402* [0.219] N=1,135**	0.102 [0.237] N=963	0.095 [0.281] N=780	0.210 [0.353] N=590
WHO weight-for-height z score	0.027 [0.298] N=1,102	0.108 [0.305] N=978	-0.048 [0.339] N=838	-0.144 [0.421] N=681	-0.490 [0.589] N=521
WHO BMI-for-age z score	-0.192 [0.292] N=1,098	-0.162 [0.324] N=976	-0.533 [0.357] N=836	-0.672 [0.439] N=678	-0.829 [0.589] N=524
Stunted HAZ below -2	-0.012 [0.071] N=1,152	-0.012 [0.078] N=1,020	-0.047 [0.086] N=870	-0.011 [0.107] N=707	0.001 [0.125] N=542
Underweight WAZ below -2	-0.046 [0.036] N=1,271	**-0.076* [0.039] N=1,135**	-0.045 [0.044] N=963	-0.060 [0.051] N=780	-0.081 [0.053] N=590
Wasted WHZ below -2	**0.089** [0.042] N=1,102**	0.064 [0.048] N=978	**0.128** [0.052] N=838**	**0.101* [0.054] N=681**	**0.156** [0.063] N=521**
Overweight WHZ above +2	0.044 [0.064] N=1,102	0.051 [0.067] N=978	0.052 [0.077] N=838	0.029 [0.096] N=681	0.003 [0.128] N=521
Obese WHZ above +3	0.005 [0.048] N=1,102	0.012 [0.050] N=978	-0.015 [0.057] N=838	0.001 [0.075] N=681	-0.064 [0.092] N=521
CIAF any stunting underweight or wasting	0.069 [0.076] N=1,082	0.014 [0.087] N=958	0.067 [0.098] N=819	0.046 [0.114] N=664	0.134 [0.131] N=509
Below-normal BMI-for-age BMIZ below -2	**0.095** [0.038] N=1,098**	**0.083* [0.047] N=976**	**0.151*** [0.051] N=836**	**0.158*** [0.054] N=678**	**0.205*** [0.055] N=524**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. UNICEF child-age-interval fixed effects are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Neither aid control is combined with an income control. Aid specifications use only the six main questions; households with a blank or 99 on any question are excluded. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household; raw ages 60 to 62 months map to analysis month 59. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.