Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for the Oldest Eligible Child Using 10 October 2022 Child Age

Continuous z scores and prespecified adverse clinical classifications exclude WHO-flagged measurements. The earliest-born eligible child is selected per household using age on 10 October 2022. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 UNICEF age-interval fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.220	0.168 [0.201] N=1,522	0.168 [0.213] N=1,407	0.174 [0.217] N=1,264	0.151 [0.236] N=1,081	0.094 [0.263] N=890
WHO weight-for-age z score	0.237	0.103 [0.125] N=1,679	0.128 [0.133] N=1,550	0.046 [0.136] N=1,395	0.069 [0.155] N=1,190	0.023 [0.158] N=979
WHO weight-for-height z score	0.366	-0.135 [0.198] N=1,438	-0.129 [0.207] N=1,338	-0.290 [0.222] N=1,201	-0.323 [0.247] N=1,028	-0.266 [0.265] N=843
WHO BMI-for-age z score	0.335	-0.174 [0.220] N=1,445	-0.167 [0.233] N=1,343	-0.349 [0.247] N=1,200	-0.452 [0.279] N=1,024	-0.425 [0.301] N=844
Stunted HAZ below -2	0.202 P=297	-0.024 [0.039] N=1,522	-0.024 [0.042] N=1,407	-0.014 [0.044] N=1,264	-0.021 [0.048] N=1,081	0.006 [0.054] N=890
Underweight WAZ below -2	0.049 P=85	-0.025 [0.018] N=1,679	-0.031 [0.019] N=1,550	-0.021 [0.019] N=1,395	-0.019 [0.021] N=1,190	-0.009 [0.023] N=979
Wasted WHZ below -2	0.106 P=158	-0.002 [0.033] N=1,438	0.010 [0.034] N=1,338	0.033 [0.038] N=1,201	0.026 [0.042] N=1,028	0.012 [0.047] N=843
Overweight WHZ above +2	0.196 P=273	-0.005 [0.044] N=1,438	0.001 [0.045] N=1,338	-0.006 [0.048] N=1,201	-0.022 [0.052] N=1,028	-0.018 [0.058] N=843
Obese WHZ above +3	0.090 P=128	-0.017 [0.030] N=1,438	-0.020 [0.031] N=1,338	-0.023 [0.033] N=1,201	-0.031 [0.036] N=1,028	-0.035 [0.039] N=843
CIAF any stunting underweight or wasting	0.279 P=391	-0.056 [0.042] N=1,415	-0.047 [0.042] N=1,315	-0.011 [0.044] N=1,180	-0.029 [0.046] N=1,008	-0.009 [0.050] N=826
Below-normal BMI-for-age BMIZ below -2	0.111 P=166	0.015 [0.037] N=1,445	0.022 [0.039] N=1,343	0.049 [0.043] N=1,200	0.060 [0.048] N=1,024	0.063 [0.053] N=844

Specification 1 UNICEF age-interval fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.104 [0.310] N=1,287	0.096 [0.347] N=1,172	0.093 [0.369] N=1,029	0.063 [0.432] N=846	-0.145 [0.585] N=655
WHO weight-for-age z score	0.108 [0.175] N=1,424	0.150 [0.191] N=1,295	0.005 [0.203] N=1,140	0.080 [0.272] N=935	-0.021 [0.324] N=724
WHO weight-for-height z score	-0.087 [0.269] N=1,218	-0.076 [0.293] N=1,118	-0.373 [0.340] N=981	-0.457 [0.421] N=808	-0.307 [0.514] N=623
WHO BMI-for-age z score	0.026 [0.286] N=1,226	0.064 [0.314] N=1,124	-0.227 [0.353] N=981	-0.401 [0.442] N=805	-0.204 [0.532] N=625
Stunted HAZ below -2	-0.002 [0.054] N=1,287	-0.000 [0.061] N=1,172	0.024 [0.064] N=1,029	0.015 [0.073] N=846	0.110 [0.096] N=655
Underweight WAZ below -2	-0.036 [0.027] N=1,424	-0.046 [0.029] N=1,295	-0.032 [0.032] N=1,140	-0.031 [0.036] N=935	-0.007 [0.049] N=724
Wasted WHZ below -2	0.004 [0.036] N=1,218	0.025 [0.037] N=1,118	0.073 [0.045] N=981	0.067 [0.055] N=808	0.041 [0.061] N=623
Overweight WHZ above +2	0.009 [0.057] N=1,218	0.017 [0.061] N=1,118	0.008 [0.067] N=981	-0.026 [0.082] N=808	0.001 [0.100] N=623
Obese WHZ above +3	0.003 [0.040] N=1,218	-0.001 [0.043] N=1,118	0.002 [0.048] N=981	-0.006 [0.060] N=808	-0.007 [0.082] N=623
CIAF any stunting underweight or wasting	-0.032 [0.058] N=1,201	-0.015 [0.060] N=1,101	0.059 [0.065] N=966	0.034 [0.073] N=794	0.117 [0.092] N=612
Below-normal BMI-for-age BMIZ below -2	-0.010 [0.038] N=1,226	-0.003 [0.040] N=1,124	0.043 [0.046] N=981	0.060 [0.054] N=805	0.056 [0.065] N=625

Specification 2 UNICEF age intervals and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.220	0.172 [0.201] N=1,522	0.173 [0.213] N=1,407	0.182 [0.217] N=1,264	0.160 [0.237] N=1,081	0.102 [0.264] N=890
WHO weight-for-age z score	0.237	0.105 [0.125] N=1,679	0.130 [0.133] N=1,550	0.049 [0.136] N=1,395	0.071 [0.156] N=1,190	0.025 [0.159] N=979
WHO weight-for-height z score	0.366	-0.137 [0.199] N=1,438	-0.134 [0.207] N=1,338	-0.296 [0.223] N=1,201	-0.325 [0.248] N=1,028	-0.271 [0.266] N=843
WHO BMI-for-age z score	0.335	-0.178 [0.219] N=1,445	-0.174 [0.232] N=1,343	-0.358 [0.246] N=1,200	-0.455 [0.278] N=1,024	-0.432 [0.298] N=844
Stunted HAZ below -2	0.202 P=297	-0.024 [0.040] N=1,522	-0.025 [0.042] N=1,407	-0.015 [0.044] N=1,264	-0.022 [0.048] N=1,081	0.004 [0.054] N=890
Underweight WAZ below -2	0.049 P=85	-0.026 [0.018] N=1,679	-0.031 [0.019] N=1,550	-0.022 [0.019] N=1,395	-0.019 [0.021] N=1,190	-0.009 [0.023] N=979
Wasted WHZ below -2	0.106 P=158	-0.001 [0.033] N=1,438	0.011 [0.034] N=1,338	0.036 [0.037] N=1,201	0.027 [0.041] N=1,028	0.013 [0.046] N=843
Overweight WHZ above +2	0.196 P=273	-0.004 [0.044] N=1,438	0.001 [0.045] N=1,338	-0.005 [0.048] N=1,201	-0.021 [0.052] N=1,028	-0.017 [0.058] N=843
Obese WHZ above +3	0.090 P=128	-0.017 [0.030] N=1,438	-0.020 [0.030] N=1,338	-0.022 [0.032] N=1,201	-0.031 [0.036] N=1,028	-0.035 [0.039] N=843
CIAF any stunting underweight or wasting	0.279 P=391	-0.056 [0.041] N=1,415	-0.046 [0.042] N=1,315	-0.010 [0.044] N=1,180	-0.028 [0.045] N=1,008	-0.008 [0.050] N=826
Below-normal BMI-for-age BMIZ below -2	0.111 P=166	0.016 [0.037] N=1,445	0.023 [0.038] N=1,343	0.051 [0.042] N=1,200	0.061 [0.046] N=1,024	0.064 [0.052] N=844

Specification 2 UNICEF age intervals and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.110 [0.309] N=1,287	0.106 [0.346] N=1,172	0.114 [0.368] N=1,029	0.092 [0.428] N=846	-0.109 [0.583] N=655
WHO weight-for-age z score	0.109 [0.174] N=1,424	0.151 [0.191] N=1,295	0.009 [0.203] N=1,140	0.083 [0.271] N=935	-0.018 [0.324] N=724
WHO weight-for-height z score	-0.093 [0.270] N=1,218	-0.087 [0.294] N=1,118	-0.393 [0.342] N=981	-0.471 [0.423] N=808	-0.340 [0.522] N=623
WHO BMI-for-age z score	0.020 [0.286] N=1,226	0.050 [0.316] N=1,124	-0.249 [0.354] N=981	-0.414 [0.445] N=805	-0.246 [0.544] N=625
Stunted HAZ below -2	-0.003 [0.054] N=1,287	-0.001 [0.061] N=1,172	0.023 [0.064] N=1,029	0.014 [0.074] N=846	0.107 [0.097] N=655
Underweight WAZ below -2	-0.036 [0.027] N=1,424	-0.046 [0.029] N=1,295	-0.033 [0.032] N=1,140	-0.031 [0.037] N=935	-0.008 [0.049] N=724
Wasted WHZ below -2	0.005 [0.037] N=1,218	0.027 [0.038] N=1,118	**0.078* [0.046] N=981**	0.071 [0.055] N=808	0.049 [0.063] N=623
Overweight WHZ above +2	0.010 [0.057] N=1,218	0.017 [0.061] N=1,118	0.010 [0.067] N=981	-0.024 [0.082] N=808	0.005 [0.099] N=623
Obese WHZ above +3	0.004 [0.040] N=1,218	-0.000 [0.042] N=1,118	0.003 [0.048] N=981	-0.006 [0.060] N=808	-0.006 [0.082] N=623
CIAF any stunting underweight or wasting	-0.032 [0.058] N=1,201	-0.012 [0.060] N=1,101	0.063 [0.066] N=966	0.037 [0.073] N=794	0.123 [0.091] N=612
Below-normal BMI-for-age BMIZ below -2	-0.009 [0.039] N=1,226	-0.000 [0.040] N=1,124	0.047 [0.047] N=981	0.063 [0.055] N=805	0.063 [0.068] N=625

Specification 3 UNICEF age intervals female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.220	0.122 [0.199] N=1,522	0.109 [0.211] N=1,407	0.134 [0.216] N=1,264	0.091 [0.238] N=1,081	0.025 [0.264] N=890
WHO weight-for-age z score	0.237	0.040 [0.123] N=1,679	0.066 [0.132] N=1,550	-0.013 [0.136] N=1,395	-0.001 [0.157] N=1,190	-0.056 [0.164] N=979
WHO weight-for-height z score	0.366	-0.179 [0.197] N=1,438	-0.169 [0.206] N=1,338	-0.330 [0.221] N=1,201	-0.364 [0.247] N=1,028	-0.314 [0.268] N=843
WHO BMI-for-age z score	0.335	-0.198 [0.219] N=1,445	-0.191 [0.232] N=1,343	-0.374 [0.245] N=1,200	**-0.475* [0.277] N=1,024**	-0.447 [0.302] N=844
Stunted HAZ below -2	0.202 P=297	-0.014 [0.039] N=1,522	-0.010 [0.041] N=1,407	-0.005 [0.043] N=1,264	-0.012 [0.047] N=1,081	0.017 [0.053] N=890
Underweight WAZ below -2	0.049 P=85	-0.017 [0.018] N=1,679	-0.020 [0.019] N=1,550	-0.012 [0.020] N=1,395	-0.009 [0.020] N=1,190	0.003 [0.023] N=979
Wasted WHZ below -2	0.106 P=158	0.009 [0.032] N=1,438	0.023 [0.033] N=1,338	0.045 [0.036] N=1,201	0.037 [0.040] N=1,028	0.023 [0.045] N=843
Overweight WHZ above +2	0.196 P=273	-0.007 [0.044] N=1,438	0.001 [0.046] N=1,338	-0.006 [0.049] N=1,201	-0.025 [0.053] N=1,028	-0.023 [0.059] N=843
Obese WHZ above +3	0.090 P=128	-0.016 [0.030] N=1,438	-0.018 [0.030] N=1,338	-0.021 [0.032] N=1,201	-0.032 [0.035] N=1,028	-0.038 [0.040] N=843
CIAF any stunting underweight or wasting	0.279 P=391	-0.036 [0.041] N=1,415	-0.021 [0.040] N=1,315	0.010 [0.042] N=1,180	-0.009 [0.043] N=1,008	0.014 [0.048] N=826
Below-normal BMI-for-age BMIZ below -2	0.111 P=166	0.023 [0.037] N=1,445	0.033 [0.038] N=1,343	0.059 [0.041] N=1,200	0.069 [0.045] N=1,024	0.072 [0.051] N=844

Specification 3 UNICEF age intervals female and household income groups 12 month donut hole

Outcome	**8 years**	**7 years**	**6 years**	**5 years**	**4 years**
WHO height-for-age z score	0.072 [0.303] N=1,287	0.043 [0.338] N=1,172	0.065 [0.363] N=1,029	0.000 [0.425] N=846	-0.234 [0.568] N=655
WHO weight-for-age z score	0.051 [0.169] N=1,424	0.090 [0.186] N=1,295	-0.050 [0.199] N=1,140	-0.002 [0.268] N=935	-0.132 [0.330] N=724
WHO weight-for-height z score	-0.120 [0.269] N=1,218	-0.105 [0.295] N=1,118	-0.404 [0.343] N=981	-0.502 [0.430] N=808	-0.384 [0.538] N=623
WHO BMI-for-age z score	0.009 [0.286] N=1,226	0.040 [0.315] N=1,124	-0.255 [0.355] N=981	-0.450 [0.453] N=805	-0.271 [0.564] N=625
Stunted HAZ below -2	0.004 [0.053] N=1,287	0.013 [0.058] N=1,172	0.032 [0.063] N=1,029	0.021 [0.072] N=846	0.118 [0.093] N=655
Underweight WAZ below -2	-0.028 [0.027] N=1,424	-0.035 [0.030] N=1,295	-0.024 [0.032] N=1,140	-0.023 [0.037] N=935	0.004 [0.048] N=724
Wasted WHZ below -2	0.013 [0.037] N=1,218	0.038 [0.038] N=1,118	**0.087* [0.045] N=981**	0.085 [0.055] N=808	0.060 [0.065] N=623
Overweight WHZ above +2	0.011 [0.057] N=1,218	0.023 [0.061] N=1,118	0.015 [0.068] N=981	-0.023 [0.083] N=808	0.003 [0.101] N=623
Obese WHZ above +3	0.006 [0.040] N=1,218	0.005 [0.042] N=1,118	0.007 [0.048] N=981	-0.005 [0.060] N=808	-0.014 [0.082] N=623
CIAF any stunting underweight or wasting	-0.016 [0.058] N=1,201	0.012 [0.058] N=1,101	0.081 [0.064] N=966	0.055 [0.071] N=794	0.144 [0.090] N=612
Below-normal BMI-for-age BMIZ below -2	-0.004 [0.039] N=1,226	0.009 [0.041] N=1,124	0.054 [0.047] N=981	0.076 [0.055] N=805	0.073 [0.069] N=625

Specification 4 UNICEF age intervals female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.220	0.056 [0.199] N=1,522	0.036 [0.211] N=1,407	0.061 [0.220] N=1,264	0.011 [0.236] N=1,081	-0.086 [0.262] N=890
WHO weight-for-age z score	0.237	0.047 [0.122] N=1,679	0.066 [0.130] N=1,550	-0.011 [0.137] N=1,395	-0.007 [0.159] N=1,190	-0.084 [0.163] N=979
WHO weight-for-height z score	0.366	-0.100 [0.197] N=1,438	-0.088 [0.205] N=1,338	-0.261 [0.222] N=1,201	-0.304 [0.249] N=1,028	-0.236 [0.267] N=843
WHO BMI-for-age z score	0.335	-0.134 [0.220] N=1,445	-0.123 [0.233] N=1,343	-0.319 [0.249] N=1,200	-0.417 [0.284] N=1,024	-0.382 [0.310] N=844
Stunted HAZ below -2	0.202 P=297	-0.003 [0.039] N=1,522	0.001 [0.042] N=1,407	0.008 [0.044] N=1,264	0.008 [0.047] N=1,081	0.040 [0.054] N=890
Underweight WAZ below -2	0.049 P=85	-0.016 [0.019] N=1,679	-0.019 [0.020] N=1,550	-0.010 [0.021] N=1,395	-0.005 [0.021] N=1,190	0.009 [0.024] N=979
Wasted WHZ below -2	0.106 P=158	0.004 [0.033] N=1,438	0.018 [0.034] N=1,338	0.042 [0.037] N=1,201	0.036 [0.041] N=1,028	0.020 [0.046] N=843
Overweight WHZ above +2	0.196 P=273	0.004 [0.044] N=1,438	0.009 [0.046] N=1,338	-0.000 [0.049] N=1,201	-0.017 [0.054] N=1,028	-0.015 [0.059] N=843
Obese WHZ above +3	0.090 P=128	-0.006 [0.030] N=1,438	-0.007 [0.030] N=1,338	-0.011 [0.033] N=1,201	-0.018 [0.036] N=1,028	-0.020 [0.040] N=843
CIAF any stunting underweight or wasting	0.279 P=391	-0.030 [0.042] N=1,415	-0.017 [0.042] N=1,315	0.016 [0.044] N=1,180	0.007 [0.046] N=1,008	0.031 [0.051] N=826
Below-normal BMI-for-age BMIZ below -2	0.111 P=166	0.018 [0.037] N=1,445	0.027 [0.039] N=1,343	0.055 [0.042] N=1,200	0.066 [0.047] N=1,024	0.067 [0.053] N=844

Specification 4 UNICEF age intervals female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.048 [0.298] N=1,287	0.006 [0.331] N=1,172	0.042 [0.360] N=1,029	0.014 [0.413] N=846	-0.294 [0.553] N=655
WHO weight-for-age z score	0.057 [0.169] N=1,424	0.093 [0.186] N=1,295	-0.048 [0.203] N=1,140	0.004 [0.274] N=935	-0.120 [0.332] N=724
WHO weight-for-height z score	-0.088 [0.270] N=1,218	-0.051 [0.292] N=1,118	-0.370 [0.342] N=981	-0.480 [0.433] N=808	-0.282 [0.536] N=623
WHO BMI-for-age z score	0.030 [0.287] N=1,226	0.086 [0.314] N=1,124	-0.229 [0.357] N=981	-0.429 [0.453] N=805	-0.193 [0.560] N=625
Stunted HAZ below -2	0.011 [0.053] N=1,287	0.020 [0.058] N=1,172	0.036 [0.063] N=1,029	0.025 [0.071] N=846	0.136 [0.093] N=655
Underweight WAZ below -2	-0.027 [0.027] N=1,424	-0.034 [0.029] N=1,295	-0.021 [0.033] N=1,140	-0.016 [0.038] N=935	0.012 [0.051] N=724
Wasted WHZ below -2	0.015 [0.037] N=1,218	0.037 [0.039] N=1,118	**0.087* [0.046] N=981**	0.086 [0.055] N=808	0.056 [0.065] N=623
Overweight WHZ above +2	0.018 [0.058] N=1,218	0.030 [0.061] N=1,118	0.019 [0.068] N=981	-0.014 [0.085] N=808	0.027 [0.103] N=623
Obese WHZ above +3	0.011 [0.040] N=1,218	0.012 [0.042] N=1,118	0.014 [0.048] N=981	0.004 [0.059] N=808	0.006 [0.081] N=623
CIAF any stunting underweight or wasting	-0.006 [0.058] N=1,201	0.019 [0.059] N=1,101	0.087 [0.065] N=966	0.065 [0.071] N=794	**0.163* [0.090] N=612**
Below-normal BMI-for-age BMIZ below -2	-0.002 [0.039] N=1,226	0.006 [0.042] N=1,124	0.053 [0.048] N=981	0.075 [0.055] N=805	0.066 [0.069] N=625

Specification 5 UNICEF age intervals female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.233	0.107 [0.213] N=1,433	0.090 [0.226] N=1,330	0.106 [0.227] N=1,192	0.141 [0.242] N=1,022	0.051 [0.282] N=838
WHO weight-for-age z score	0.227	0.099 [0.131] N=1,581	0.121 [0.137] N=1,465	0.037 [0.144] N=1,317	0.058 [0.166] N=1,127	0.009 [0.175] N=924
WHO weight-for-height z score	0.371	-0.068 [0.206] N=1,356	-0.064 [0.212] N=1,268	-0.234 [0.230] N=1,137	-0.283 [0.255] N=976	-0.199 [0.282] N=798
WHO BMI-for-age z score	0.348	-0.104 [0.223] N=1,360	-0.101 [0.233] N=1,270	-0.273 [0.247] N=1,134	-0.398 [0.279] N=970	-0.346 [0.307] N=797
Stunted HAZ below -2	0.204 P=281	-0.014 [0.042] N=1,433	-0.014 [0.045] N=1,330	-0.007 [0.046] N=1,192	-0.025 [0.049] N=1,022	0.007 [0.056] N=838
Underweight WAZ below -2	0.051 P=84	-0.025 [0.020] N=1,581	-0.030 [0.020] N=1,465	-0.020 [0.021] N=1,317	-0.018 [0.023] N=1,127	-0.006 [0.026] N=924
Wasted WHZ below -2	0.106 P=150	0.003 [0.035] N=1,356	0.012 [0.035] N=1,268	0.040 [0.039] N=1,137	0.032 [0.044] N=976	0.017 [0.049] N=798
Overweight WHZ above +2	0.198 P=258	0.001 [0.046] N=1,356	0.003 [0.047] N=1,268	-0.004 [0.051] N=1,137	-0.027 [0.054] N=976	-0.024 [0.062] N=798
Obese WHZ above +3	0.092 P=124	-0.012 [0.031] N=1,356	-0.015 [0.032] N=1,268	-0.018 [0.034] N=1,137	-0.028 [0.037] N=976	-0.032 [0.042] N=798
CIAF any stunting underweight or wasting	0.282 P=372	-0.039 [0.045] N=1,334	-0.033 [0.045] N=1,246	0.001 [0.047] N=1,116	-0.026 [0.050] N=956	-0.004 [0.056] N=781
Below-normal BMI-for-age BMIZ below -2	0.110 P=155	0.016 [0.037] N=1,360	0.020 [0.038] N=1,270	0.049 [0.041] N=1,134	0.059 [0.046] N=970	0.059 [0.051] N=797

Specification 5 UNICEF age intervals female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.008 [0.319] N=1,213	-0.041 [0.356] N=1,110	-0.035 [0.373] N=972	0.062 [0.420] N=802	-0.224 [0.604] N=618
WHO weight-for-age z score	0.090 [0.179] N=1,342	0.124 [0.191] N=1,226	-0.036 [0.212] N=1,078	0.042 [0.282] N=888	-0.079 [0.351] N=685
WHO weight-for-height z score	-0.009 [0.272] N=1,148	-0.004 [0.290] N=1,060	-0.305 [0.338] N=929	-0.405 [0.417] N=768	-0.168 [0.508] N=590
WHO BMI-for-age z score	0.065 [0.289] N=1,155	0.089 [0.312] N=1,065	-0.184 [0.353] N=929	-0.385 [0.438] N=765	-0.134 [0.538] N=592
Stunted HAZ below -2	0.017 [0.058] N=1,213	0.021 [0.064] N=1,110	0.040 [0.069] N=972	0.010 [0.076] N=802	0.105 [0.103] N=618
Underweight WAZ below -2	-0.040 [0.028] N=1,342	**-0.050* [0.029] N=1,226**	-0.032 [0.033] N=1,078	-0.034 [0.038] N=888	-0.005 [0.051] N=685
Wasted WHZ below -2	0.004 [0.037] N=1,148	0.021 [0.037] N=1,060	**0.079* [0.045] N=929**	0.068 [0.055] N=768	0.034 [0.061] N=590
Overweight WHZ above +2	0.024 [0.058] N=1,148	0.025 [0.061] N=1,060	0.026 [0.068] N=929	-0.019 [0.083] N=768	0.025 [0.104] N=590
Obese WHZ above +3	0.015 [0.042] N=1,148	0.013 [0.044] N=1,060	0.017 [0.049] N=929	0.008 [0.061] N=768	0.023 [0.085] N=590
CIAF any stunting underweight or wasting	-0.005 [0.061] N=1,132	0.009 [0.062] N=1,044	0.086 [0.069] N=914	0.040 [0.075] N=754	0.113 [0.099] N=579
Below-normal BMI-for-age BMIZ below -2	-0.005 [0.038] N=1,155	-0.001 [0.038] N=1,065	0.049 [0.045] N=929	0.065 [0.053] N=765	0.052 [0.066] N=592

Specification 6 UNICEF age intervals female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.233	0.105 [0.211] N=1,433	0.089 [0.223] N=1,330	0.071 [0.228] N=1,192	0.090 [0.247] N=1,022	-0.024 [0.285] N=838
WHO weight-for-age z score	0.227	0.102 [0.131] N=1,581	0.123 [0.138] N=1,465	0.035 [0.143] N=1,317	0.064 [0.164] N=1,127	0.013 [0.171] N=924
WHO weight-for-height z score	0.371	-0.068 [0.205] N=1,356	-0.063 [0.211] N=1,268	-0.227 [0.228] N=1,137	-0.268 [0.254] N=976	-0.182 [0.278] N=798
WHO BMI-for-age z score	0.348	-0.106 [0.222] N=1,360	-0.103 [0.232] N=1,270	-0.274 [0.246] N=1,134	-0.383 [0.280] N=970	-0.329 [0.307] N=797
Stunted HAZ below -2	0.204 P=281	-0.014 [0.042] N=1,433	-0.014 [0.044] N=1,330	-0.004 [0.046] N=1,192	-0.020 [0.049] N=1,022	0.011 [0.056] N=838
Underweight WAZ below -2	0.051 P=84	-0.028 [0.020] N=1,581	-0.032 [0.020] N=1,465	-0.023 [0.021] N=1,317	-0.021 [0.023] N=1,127	-0.009 [0.026] N=924
Wasted WHZ below -2	0.106 P=150	0.002 [0.035] N=1,356	0.011 [0.035] N=1,268	0.039 [0.039] N=1,137	0.030 [0.044] N=976	0.011 [0.049] N=798
Overweight WHZ above +2	0.198 P=258	0.001 [0.046] N=1,356	0.003 [0.047] N=1,268	-0.000 [0.050] N=1,137	-0.022 [0.054] N=976	-0.017 [0.061] N=798
Obese WHZ above +3	0.092 P=124	-0.012 [0.031] N=1,356	-0.014 [0.031] N=1,268	-0.016 [0.034] N=1,137	-0.025 [0.036] N=976	-0.027 [0.041] N=798
CIAF any stunting underweight or wasting	0.282 P=372	-0.040 [0.044] N=1,334	-0.034 [0.045] N=1,246	0.002 [0.047] N=1,116	-0.026 [0.049] N=956	-0.010 [0.054] N=781
Below-normal BMI-for-age BMIZ below -2	0.110 P=155	0.016 [0.037] N=1,360	0.020 [0.038] N=1,270	0.048 [0.042] N=1,134	0.058 [0.047] N=970	0.055 [0.052] N=797

Specification 6 UNICEF age intervals female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.070 [0.322] N=1,213	0.038 [0.357] N=1,110	-0.004 [0.380] N=972	0.082 [0.442] N=802	-0.253 [0.629] N=618
WHO weight-for-age z score	0.106 [0.179] N=1,342	0.143 [0.193] N=1,226	-0.012 [0.209] N=1,078	0.078 [0.275] N=888	-0.031 [0.339] N=685
WHO weight-for-height z score	-0.012 [0.271] N=1,148	-0.001 [0.289] N=1,060	-0.304 [0.336] N=929	-0.400 [0.420] N=768	-0.159 [0.514] N=590
WHO BMI-for-age z score	0.068 [0.289] N=1,155	0.097 [0.312] N=1,065	-0.182 [0.350] N=929	-0.379 [0.443] N=765	-0.131 [0.546] N=592
Stunted HAZ below -2	0.009 [0.058] N=1,213	0.012 [0.064] N=1,110	0.037 [0.069] N=972	0.003 [0.078] N=802	0.103 [0.103] N=618
Underweight WAZ below -2	-0.041 [0.028] N=1,342	**-0.051* [0.030] N=1,226**	-0.037 [0.033] N=1,078	-0.041 [0.038] N=888	-0.016 [0.051] N=685
Wasted WHZ below -2	0.007 [0.037] N=1,148	0.023 [0.037] N=1,060	**0.081* [0.045] N=929**	0.069 [0.055] N=768	0.029 [0.061] N=590
Overweight WHZ above +2	0.023 [0.058] N=1,148	0.027 [0.060] N=1,060	0.026 [0.068] N=929	-0.016 [0.083] N=768	0.029 [0.104] N=590
Obese WHZ above +3	0.014 [0.042] N=1,148	0.011 [0.044] N=1,060	0.019 [0.050] N=929	0.009 [0.061] N=768	0.026 [0.085] N=590
CIAF any stunting underweight or wasting	-0.006 [0.061] N=1,132	0.007 [0.063] N=1,044	0.084 [0.070] N=914	0.034 [0.076] N=754	0.101 [0.098] N=579
Below-normal BMI-for-age BMIZ below -2	-0.004 [0.038] N=1,155	0.000 [0.038] N=1,065	0.049 [0.045] N=929	0.064 [0.054] N=765	0.047 [0.067] N=592

Mother RDD secondary analysis

Specification 1 UNICEF age-interval fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.215	-0.123 [0.240] N=1,415	-0.143 [0.254] N=1,278	-0.141 [0.279] N=1,119	-0.195 [0.301] N=946	-0.170 [0.326] N=766
WHO weight-for-age z score	0.229	0.100 [0.137] N=1,552	0.179 [0.146] N=1,411	0.031 [0.145] N=1,229	-0.010 [0.157] N=1,035	0.040 [0.170] N=831
WHO weight-for-height z score	0.385	0.019 [0.191] N=1,341	0.074 [0.194] N=1,214	0.044 [0.204] N=1,066	0.000 [0.224] N=899	-0.025 [0.246] N=727
WHO BMI-for-age z score	0.331	-0.104 [0.220] N=1,340	-0.033 [0.233] N=1,214	-0.126 [0.250] N=1,066	-0.143 [0.271] N=898	-0.093 [0.296] N=732
Stunted HAZ below -2	0.204 P=284	0.035 [0.046] N=1,415	0.046 [0.049] N=1,278	0.042 [0.050] N=1,119	0.042 [0.055] N=946	0.049 [0.062] N=766
Underweight WAZ below -2	0.047 P=75	-0.003 [0.019] N=1,552	-0.018 [0.020] N=1,411	0.001 [0.021] N=1,229	-0.006 [0.022] N=1,035	-0.015 [0.024] N=831
Wasted WHZ below -2	0.104 P=139	0.047 [0.032] N=1,341	0.029 [0.035] N=1,214	0.059 [0.036] N=1,066	0.041 [0.036] N=899	0.049 [0.040] N=727
Overweight WHZ above +2	0.193 P=262	0.022 [0.040] N=1,341	0.027 [0.041] N=1,214	0.045 [0.044] N=1,066	0.031 [0.049] N=899	0.040 [0.051] N=727
Obese WHZ above +3	0.095 P=132	0.019 [0.030] N=1,341	0.022 [0.031] N=1,214	0.020 [0.033] N=1,066	0.018 [0.038] N=899	0.003 [0.038] N=727
CIAF any stunting underweight or wasting	0.280 P=366	0.080 [0.049] N=1,319	0.057 [0.052] N=1,192	**0.095* [0.054] N=1,045**	0.078 [0.057] N=881	**0.104* [0.060] N=712**
Below-normal BMI-for-age BMIZ below -2	0.108 P=145	0.044 [0.038] N=1,340	0.022 [0.041] N=1,214	0.053 [0.043] N=1,066	0.046 [0.046] N=898	0.057 [0.050] N=732

Specification 1 UNICEF age-interval fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.225 [0.315] N=1,226	0.268 [0.342] N=1,089	0.371 [0.404] N=930	0.393 [0.501] N=757	0.770 [0.575] N=577
WHO weight-for-age z score	0.153 [0.201] N=1,348	0.315 [0.215] N=1,207	0.062 [0.226] N=1,025	0.003 [0.265] N=831	0.148 [0.321] N=627
WHO weight-for-height z score	-0.219 [0.292] N=1,166	-0.161 [0.301] N=1,039	-0.236 [0.334] N=891	-0.388 [0.413] N=724	-0.692 [0.530] N=552
WHO BMI-for-age z score	-0.474 [0.289] N=1,166	-0.404 [0.317] N=1,040	**-0.643* [0.354] N=892**	**-0.805* [0.438] N=724**	**-0.938* [0.547] N=558**
Stunted HAZ below -2	0.007 [0.068] N=1,226	0.020 [0.076] N=1,089	0.003 [0.083] N=930	-0.013 [0.102] N=757	-0.022 [0.131] N=577
Underweight WAZ below -2	-0.007 [0.028] N=1,348	-0.035 [0.031] N=1,207	-0.001 [0.036] N=1,025	-0.024 [0.039] N=831	-0.058 [0.045] N=627
Wasted WHZ below -2	**0.093** [0.041] N=1,166**	0.068 [0.047] N=1,039	**0.132** [0.050] N=891**	**0.107** [0.050] N=724**	**0.158*** [0.057] N=552**
Overweight WHZ above +2	-0.008 [0.062] N=1,166	-0.008 [0.066] N=1,039	0.021 [0.074] N=891	-0.021 [0.094] N=724	-0.044 [0.111] N=552
Obese WHZ above +3	0.003 [0.046] N=1,166	0.003 [0.048] N=1,039	-0.003 [0.057] N=891	-0.012 [0.073] N=724	-0.086 [0.082] N=552
CIAF any stunting underweight or wasting	0.081 [0.071] N=1,148	0.038 [0.080] N=1,021	0.102 [0.090] N=874	0.056 [0.104] N=710	0.108 [0.122] N=541
Below-normal BMI-for-age BMIZ below -2	**0.094** [0.040] N=1,166**	0.063 [0.046] N=1,040	**0.129** [0.049] N=892**	**0.128** [0.050] N=724**	**0.181*** [0.054] N=558**

Specification 2 UNICEF age intervals and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.215	-0.127 [0.239] N=1,415	-0.146 [0.253] N=1,278	-0.142 [0.278] N=1,119	-0.194 [0.301] N=946	-0.151 [0.326] N=766
WHO weight-for-age z score	0.229	0.100 [0.137] N=1,552	0.179 [0.145] N=1,411	0.031 [0.145] N=1,229	-0.009 [0.157] N=1,035	0.042 [0.171] N=831
WHO weight-for-height z score	0.385	0.021 [0.191] N=1,341	0.075 [0.195] N=1,214	0.046 [0.205] N=1,066	-0.003 [0.225] N=899	-0.046 [0.249] N=727
WHO BMI-for-age z score	0.331	-0.101 [0.220] N=1,340	-0.032 [0.233] N=1,214	-0.126 [0.250] N=1,066	-0.146 [0.272] N=898	-0.119 [0.299] N=732
Stunted HAZ below -2	0.204 P=284	0.035 [0.046] N=1,415	0.046 [0.049] N=1,278	0.043 [0.050] N=1,119	0.042 [0.055] N=946	0.048 [0.062] N=766
Underweight WAZ below -2	0.047 P=75	-0.003 [0.019] N=1,552	-0.018 [0.020] N=1,411	0.001 [0.021] N=1,229	-0.006 [0.022] N=1,035	-0.014 [0.024] N=831
Wasted WHZ below -2	0.104 P=139	0.046 [0.032] N=1,341	0.029 [0.035] N=1,214	0.059 [0.036] N=1,066	0.042 [0.037] N=899	0.055 [0.041] N=727
Overweight WHZ above +2	0.193 P=262	0.022 [0.040] N=1,341	0.027 [0.041] N=1,214	0.045 [0.044] N=1,066	0.031 [0.049] N=899	0.041 [0.052] N=727
Obese WHZ above +3	0.095 P=132	0.019 [0.030] N=1,341	0.022 [0.030] N=1,214	0.019 [0.033] N=1,066	0.018 [0.038] N=899	0.006 [0.038] N=727
CIAF any stunting underweight or wasting	0.280 P=366	0.080 [0.049] N=1,319	0.057 [0.053] N=1,192	**0.094* [0.055] N=1,045**	0.079 [0.058] N=881	**0.108* [0.061] N=712**
Below-normal BMI-for-age BMIZ below -2	0.108 P=145	0.043 [0.038] N=1,340	0.022 [0.041] N=1,214	0.053 [0.044] N=1,066	0.047 [0.046] N=898	0.062 [0.051] N=732

Specification 2 UNICEF age intervals and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.211 [0.312] N=1,226	0.255 [0.340] N=1,089	0.365 [0.404] N=930	0.369 [0.496] N=757	0.754 [0.567] N=577
WHO weight-for-age z score	0.152 [0.199] N=1,348	0.313 [0.213] N=1,207	0.060 [0.224] N=1,025	0.001 [0.262] N=831	0.148 [0.321] N=627
WHO weight-for-height z score	-0.202 [0.292] N=1,166	-0.143 [0.301] N=1,039	-0.220 [0.333] N=891	-0.364 [0.412] N=724	-0.677 [0.536] N=552
WHO BMI-for-age z score	-0.450 [0.290] N=1,166	-0.381 [0.318] N=1,040	**-0.624* [0.354] N=892**	**-0.775* [0.439] N=724**	**-0.925* [0.555] N=558**
Stunted HAZ below -2	0.010 [0.068] N=1,226	0.023 [0.075] N=1,089	0.004 [0.083] N=930	-0.010 [0.102] N=757	-0.020 [0.131] N=577
Underweight WAZ below -2	-0.007 [0.029] N=1,348	-0.035 [0.031] N=1,207	-0.001 [0.036] N=1,025	-0.024 [0.040] N=831	-0.058 [0.045] N=627
Wasted WHZ below -2	**0.088** [0.041] N=1,166**	0.062 [0.047] N=1,039	**0.124** [0.051] N=891**	**0.098* [0.050] N=724**	**0.154*** [0.058] N=552**
Overweight WHZ above +2	-0.008 [0.062] N=1,166	-0.008 [0.065] N=1,039	0.020 [0.074] N=891	-0.021 [0.094] N=724	-0.043 [0.111] N=552
Obese WHZ above +3	0.001 [0.046] N=1,166	-0.000 [0.048] N=1,039	-0.008 [0.056] N=891	-0.016 [0.073] N=724	-0.088 [0.081] N=552
CIAF any stunting underweight or wasting	0.080 [0.072] N=1,148	0.035 [0.081] N=1,021	0.096 [0.092] N=874	0.049 [0.106] N=710	0.105 [0.125] N=541
Below-normal BMI-for-age BMIZ below -2	**0.089** [0.040] N=1,166**	0.057 [0.046] N=1,040	**0.123** [0.049] N=892**	**0.120** [0.050] N=724**	**0.178*** [0.054] N=558**

Specification 3 UNICEF age intervals female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.215	-0.153 [0.236] N=1,415	-0.160 [0.251] N=1,278	-0.158 [0.277] N=1,119	-0.201 [0.298] N=946	-0.179 [0.315] N=766
WHO weight-for-age z score	0.229	0.083 [0.137] N=1,552	0.171 [0.146] N=1,411	0.023 [0.147] N=1,229	-0.021 [0.158] N=1,035	0.028 [0.170] N=831
WHO weight-for-height z score	0.385	0.018 [0.195] N=1,341	0.075 [0.199] N=1,214	0.059 [0.212] N=1,066	0.012 [0.233] N=899	-0.019 [0.259] N=727
WHO BMI-for-age z score	0.331	-0.094 [0.221] N=1,340	-0.026 [0.235] N=1,214	-0.105 [0.254] N=1,066	-0.132 [0.277] N=898	-0.094 [0.305] N=732
Stunted HAZ below -2	0.204 P=284	0.040 [0.046] N=1,415	0.049 [0.048] N=1,278	0.047 [0.051] N=1,119	0.045 [0.055] N=946	0.053 [0.061] N=766
Underweight WAZ below -2	0.047 P=75	-0.000 [0.019] N=1,552	-0.016 [0.021] N=1,411	0.004 [0.021] N=1,229	-0.004 [0.022] N=1,035	-0.010 [0.023] N=831
Wasted WHZ below -2	0.104 P=139	0.046 [0.032] N=1,341	0.027 [0.035] N=1,214	0.055 [0.037] N=1,066	0.037 [0.037] N=899	0.048 [0.040] N=727
Overweight WHZ above +2	0.193 P=262	0.022 [0.040] N=1,341	0.027 [0.041] N=1,214	0.047 [0.044] N=1,066	0.031 [0.050] N=899	0.041 [0.052] N=727
Obese WHZ above +3	0.095 P=132	0.020 [0.030] N=1,341	0.023 [0.031] N=1,214	0.021 [0.034] N=1,066	0.019 [0.039] N=899	0.008 [0.039] N=727
CIAF any stunting underweight or wasting	0.280 P=366	**0.085* [0.048] N=1,319**	0.059 [0.053] N=1,192	**0.096* [0.055] N=1,045**	0.077 [0.058] N=881	**0.105* [0.060] N=712**
Below-normal BMI-for-age BMIZ below -2	0.108 P=145	0.042 [0.038] N=1,340	0.021 [0.041] N=1,214	0.050 [0.044] N=1,066	0.044 [0.046] N=898	0.058 [0.051] N=732

Specification 3 UNICEF age intervals female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.244 [0.313] N=1,226	0.306 [0.341] N=1,089	0.450 [0.406] N=930	0.488 [0.503] N=757	0.935 [0.566] N=577
WHO weight-for-age z score	0.145 [0.197] N=1,348	0.323 [0.212] N=1,207	0.073 [0.226] N=1,025	-0.007 [0.269] N=831	0.102 [0.337] N=627
WHO weight-for-height z score	-0.235 [0.296] N=1,166	-0.168 [0.307] N=1,039	-0.256 [0.343] N=891	-0.446 [0.433] N=724	-0.831 [0.555] N=552
WHO BMI-for-age z score	-0.460 [0.292] N=1,166	-0.390 [0.322] N=1,040	**-0.649* [0.361] N=892**	**-0.859* [0.455] N=724**	**-1.078* [0.564] N=558**
Stunted HAZ below -2	0.001 [0.069] N=1,226	0.010 [0.076] N=1,089	-0.012 [0.085] N=930	-0.036 [0.105] N=757	-0.061 [0.133] N=577
Underweight WAZ below -2	-0.010 [0.028] N=1,348	-0.040 [0.031] N=1,207	-0.008 [0.036] N=1,025	-0.030 [0.040] N=831	-0.058 [0.045] N=627
Wasted WHZ below -2	**0.087** [0.042] N=1,166**	0.060 [0.048] N=1,039	**0.122** [0.052] N=891**	**0.095* [0.052] N=724**	**0.153** [0.059] N=552**
Overweight WHZ above +2	-0.012 [0.064] N=1,166	-0.011 [0.068] N=1,039	0.015 [0.079] N=891	-0.039 [0.100] N=724	-0.074 [0.119] N=552
Obese WHZ above +3	-0.004 [0.046] N=1,166	-0.006 [0.049] N=1,039	-0.013 [0.058] N=891	-0.027 [0.075] N=724	-0.110 [0.081] N=552
CIAF any stunting underweight or wasting	0.073 [0.074] N=1,148	0.025 [0.084] N=1,021	0.085 [0.096] N=874	0.031 [0.111] N=710	0.081 [0.131] N=541
Below-normal BMI-for-age BMIZ below -2	**0.086** [0.040] N=1,166**	0.054 [0.046] N=1,040	**0.121** [0.049] N=892**	**0.117** [0.051] N=724**	**0.179*** [0.052] N=558**

Specification 4 UNICEF age intervals female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.215	-0.146 [0.234] N=1,415	-0.153 [0.249] N=1,278	-0.154 [0.277] N=1,119	-0.184 [0.298] N=946	-0.155 [0.314] N=766
WHO weight-for-age z score	0.229	0.096 [0.135] N=1,552	0.183 [0.145] N=1,411	0.034 [0.144] N=1,229	0.000 [0.156] N=1,035	0.036 [0.165] N=831
WHO weight-for-height z score	0.385	0.030 [0.190] N=1,341	0.087 [0.192] N=1,214	0.063 [0.201] N=1,066	0.005 [0.220] N=899	-0.040 [0.247] N=727
WHO BMI-for-age z score	0.331	-0.085 [0.216] N=1,340	-0.015 [0.229] N=1,214	-0.107 [0.245] N=1,066	-0.136 [0.265] N=898	-0.102 [0.292] N=732
Stunted HAZ below -2	0.204 P=284	0.041 [0.046] N=1,415	0.048 [0.049] N=1,278	0.045 [0.051] N=1,119	0.041 [0.055] N=946	0.049 [0.061] N=766
Underweight WAZ below -2	0.047 P=75	-0.001 [0.019] N=1,552	-0.018 [0.020] N=1,411	0.001 [0.021] N=1,229	-0.007 [0.022] N=1,035	-0.013 [0.024] N=831
Wasted WHZ below -2	0.104 P=139	0.046 [0.032] N=1,341	0.028 [0.035] N=1,214	0.058 [0.036] N=1,066	0.042 [0.037] N=899	0.054 [0.040] N=727
Overweight WHZ above +2	0.193 P=262	0.024 [0.040] N=1,341	0.030 [0.041] N=1,214	0.049 [0.044] N=1,066	0.034 [0.049] N=899	0.041 [0.051] N=727
Obese WHZ above +3	0.095 P=132	0.022 [0.030] N=1,341	0.024 [0.030] N=1,214	0.023 [0.033] N=1,066	0.020 [0.037] N=899	0.009 [0.038] N=727
CIAF any stunting underweight or wasting	0.280 P=366	**0.086* [0.049] N=1,319**	0.059 [0.053] N=1,192	**0.097* [0.055] N=1,045**	0.080 [0.059] N=881	**0.107* [0.061] N=712**
Below-normal BMI-for-age BMIZ below -2	0.108 P=145	0.041 [0.038] N=1,340	0.020 [0.041] N=1,214	0.052 [0.043] N=1,066	0.047 [0.046] N=898	0.057 [0.050] N=732

Specification 4 UNICEF age intervals female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.210 [0.305] N=1,226	0.275 [0.332] N=1,089	0.389 [0.402] N=930	0.450 [0.492] N=757	0.813 [0.546] N=577
WHO weight-for-age z score	0.153 [0.191] N=1,348	0.325 [0.207] N=1,207	0.080 [0.214] N=1,025	0.039 [0.251] N=831	0.133 [0.293] N=627
WHO weight-for-height z score	-0.209 [0.289] N=1,166	-0.148 [0.294] N=1,039	-0.227 [0.322] N=891	-0.398 [0.402] N=724	-0.734 [0.520] N=552
WHO BMI-for-age z score	-0.449 [0.285] N=1,166	-0.381 [0.312] N=1,040	**-0.629* [0.347] N=892**	**-0.815* [0.431] N=724**	**-0.967* [0.537] N=558**
Stunted HAZ below -2	0.011 [0.068] N=1,226	0.019 [0.076] N=1,089	-0.002 [0.085] N=930	-0.028 [0.103] N=757	-0.041 [0.131] N=577
Underweight WAZ below -2	-0.007 [0.028] N=1,348	-0.037 [0.031] N=1,207	-0.005 [0.035] N=1,025	-0.033 [0.038] N=831	-0.061 [0.043] N=627
Wasted WHZ below -2	**0.089** [0.042] N=1,166**	0.062 [0.048] N=1,039	**0.124** [0.051] N=891**	**0.097* [0.051] N=724**	**0.150*** [0.057] N=552**
Overweight WHZ above +2	-0.009 [0.061] N=1,166	-0.010 [0.065] N=1,039	0.018 [0.074] N=891	-0.029 [0.094] N=724	-0.060 [0.112] N=552
Obese WHZ above +3	0.002 [0.046] N=1,166	-0.000 [0.048] N=1,039	-0.009 [0.057] N=891	-0.022 [0.072] N=724	-0.095 [0.080] N=552
CIAF any stunting underweight or wasting	0.085 [0.073] N=1,148	0.034 [0.084] N=1,021	0.091 [0.095] N=874	0.035 [0.109] N=710	0.085 [0.127] N=541
Below-normal BMI-for-age BMIZ below -2	**0.088** [0.041] N=1,166**	0.056 [0.047] N=1,040	**0.122** [0.050] N=892**	**0.119** [0.051] N=724**	**0.171*** [0.053] N=558**

Specification 5 UNICEF age intervals female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.233	-0.181 [0.239] N=1,336	-0.205 [0.256] N=1,203	-0.205 [0.281] N=1,053	-0.228 [0.295] N=889	-0.236 [0.316] N=721
WHO weight-for-age z score	0.215	0.146 [0.139] N=1,466	0.240 [0.148] N=1,329	0.073 [0.148] N=1,157	0.027 [0.161] N=974	0.032 [0.171] N=784
WHO weight-for-height z score	0.384	0.072 [0.202] N=1,268	0.144 [0.206] N=1,145	0.093 [0.219] N=1,005	0.008 [0.244] N=848	-0.083 [0.270] N=687
WHO BMI-for-age z score	0.334	-0.038 [0.225] N=1,264	0.048 [0.242] N=1,142	-0.072 [0.258] N=1,002	-0.135 [0.279] N=844	-0.133 [0.310] N=690
Stunted HAZ below -2	0.207 P=271	0.046 [0.047] N=1,336	0.060 [0.049] N=1,203	0.054 [0.051] N=1,053	0.054 [0.055] N=889	0.075 [0.060] N=721
Underweight WAZ below -2	0.049 P=74	-0.002 [0.020] N=1,466	-0.021 [0.022] N=1,329	-0.002 [0.023] N=1,157	-0.005 [0.024] N=974	-0.008 [0.027] N=784
Wasted WHZ below -2	0.106 P=134	0.050 [0.033] N=1,268	0.029 [0.036] N=1,145	0.059 [0.037] N=1,005	0.051 [0.039] N=848	**0.072* [0.043] N=687**
Overweight WHZ above +2	0.193 P=247	0.028 [0.043] N=1,268	0.035 [0.044] N=1,145	0.051 [0.047] N=1,005	0.034 [0.053] N=848	0.042 [0.057] N=687
Obese WHZ above +3	0.098 P=128	0.019 [0.032] N=1,268	0.022 [0.032] N=1,145	0.021 [0.035] N=1,005	0.021 [0.041] N=848	0.015 [0.041] N=687
CIAF any stunting underweight or wasting	0.286 P=352	**0.088* [0.052] N=1,247**	0.066 [0.056] N=1,124	**0.101* [0.058] N=985**	0.092 [0.061] N=830	**0.142** [0.063] N=672**
Below-normal BMI-for-age BMIZ below -2	0.109 P=137	0.049 [0.039] N=1,264	0.025 [0.043] N=1,142	0.057 [0.045] N=1,002	0.065 [0.047] N=844	0.085 [0.052] N=690

Specification 5 UNICEF age intervals female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.216 [0.335] N=1,158	0.269 [0.370] N=1,025	0.389 [0.450] N=875	0.477 [0.522] N=711	0.734 [0.593] N=543
WHO weight-for-age z score	0.226 [0.203] N=1,274	**0.415* [0.215] N=1,137**	0.140 [0.228] N=965	0.092 [0.270] N=782	0.165 [0.324] N=592
WHO weight-for-height z score	-0.080 [0.301] N=1,101	0.009 [0.311] N=978	-0.088 [0.347] N=838	-0.281 [0.427] N=681	-0.624 [0.574] N=520
WHO BMI-for-age z score	-0.340 [0.300] N=1,099	-0.244 [0.334] N=977	-0.529 [0.375] N=837	-0.748 [0.457] N=679	-0.897 [0.599] N=525
Stunted HAZ below -2	0.002 [0.073] N=1,158	0.016 [0.082] N=1,025	-0.015 [0.093] N=875	-0.039 [0.111] N=711	-0.017 [0.141] N=543
Underweight WAZ below -2	-0.008 [0.030] N=1,274	-0.039 [0.033] N=1,137	-0.010 [0.037] N=965	-0.028 [0.041] N=782	-0.054 [0.047] N=592
Wasted WHZ below -2	**0.086** [0.043] N=1,101**	0.053 [0.049] N=978	**0.111** [0.052] N=838**	**0.101* [0.053] N=681**	**0.164** [0.063] N=520**
Overweight WHZ above +2	0.007 [0.066] N=1,101	0.008 [0.070] N=978	0.033 [0.079] N=838	-0.019 [0.101] N=681	-0.037 [0.121] N=520
Obese WHZ above +3	-0.002 [0.048] N=1,101	-0.005 [0.051] N=978	-0.012 [0.058] N=838	-0.018 [0.076] N=681	-0.068 [0.086] N=520
CIAF any stunting underweight or wasting	0.073 [0.078] N=1,084	0.023 [0.089] N=961	0.071 [0.102] N=822	0.023 [0.119] N=667	0.119 [0.138] N=509
Below-normal BMI-for-age BMIZ below -2	**0.087** [0.042] N=1,099**	0.049 [0.048] N=977	**0.113** [0.051] N=837**	**0.133** [0.053] N=679**	**0.193*** [0.060] N=525**

Specification 6 UNICEF age intervals female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.233	-0.185 [0.242] N=1,336	-0.206 [0.258] N=1,203	-0.197 [0.283] N=1,053	-0.236 [0.302] N=889	-0.255 [0.318] N=721
WHO weight-for-age z score	0.215	0.149 [0.141] N=1,466	0.242 [0.150] N=1,329	0.073 [0.150] N=1,157	0.033 [0.165] N=974	0.036 [0.176] N=784
WHO weight-for-height z score	0.384	0.076 [0.200] N=1,268	0.145 [0.205] N=1,145	0.088 [0.217] N=1,005	0.018 [0.241] N=848	-0.071 [0.270] N=687
WHO BMI-for-age z score	0.334	-0.035 [0.224] N=1,264	0.049 [0.241] N=1,142	-0.075 [0.257] N=1,002	-0.125 [0.279] N=844	-0.122 [0.311] N=690
Stunted HAZ below -2	0.207 P=271	0.045 [0.047] N=1,336	0.058 [0.049] N=1,203	0.051 [0.051] N=1,053	0.054 [0.056] N=889	0.074 [0.061] N=721
Underweight WAZ below -2	0.049 P=74	-0.003 [0.020] N=1,466	-0.020 [0.021] N=1,329	-0.000 [0.022] N=1,157	-0.005 [0.023] N=974	-0.008 [0.025] N=784
Wasted WHZ below -2	0.106 P=134	0.049 [0.033] N=1,268	0.029 [0.036] N=1,145	0.060 [0.038] N=1,005	0.049 [0.039] N=848	0.070 [0.043] N=687
Overweight WHZ above +2	0.193 P=247	0.028 [0.043] N=1,268	0.034 [0.044] N=1,145	0.048 [0.047] N=1,005	0.033 [0.053] N=848	0.041 [0.057] N=687
Obese WHZ above +3	0.098 P=128	0.018 [0.031] N=1,268	0.022 [0.032] N=1,145	0.019 [0.035] N=1,005	0.021 [0.040] N=848	0.015 [0.041] N=687
CIAF any stunting underweight or wasting	0.286 P=352	**0.086* [0.052] N=1,247**	0.063 [0.056] N=1,124	**0.099* [0.059] N=985**	0.090 [0.062] N=830	**0.138** [0.063] N=672**
Below-normal BMI-for-age BMIZ below -2	0.109 P=137	0.048 [0.039] N=1,264	0.025 [0.043] N=1,142	0.059 [0.045] N=1,002	0.063 [0.048] N=844	0.083 [0.052] N=690

Specification 6 UNICEF age intervals female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.196 [0.340] N=1,158	0.245 [0.374] N=1,025	0.382 [0.449] N=875	0.419 [0.542] N=711	0.653 [0.613] N=543
WHO weight-for-age z score	0.228 [0.203] N=1,274	**0.417* [0.216] N=1,137**	0.134 [0.230] N=965	0.100 [0.273] N=782	0.171 [0.334] N=592
WHO weight-for-height z score	-0.075 [0.300] N=1,101	0.016 [0.309] N=978	-0.100 [0.342] N=838	-0.261 [0.430] N=681	-0.629 [0.575] N=520
WHO BMI-for-age z score	-0.334 [0.299] N=1,099	-0.233 [0.330] N=977	-0.530 [0.367] N=837	-0.717 [0.457] N=679	-0.878 [0.596] N=525
Stunted HAZ below -2	0.004 [0.074] N=1,158	0.019 [0.082] N=1,025	-0.012 [0.092] N=875	-0.027 [0.113] N=711	-0.004 [0.141] N=543
Underweight WAZ below -2	-0.009 [0.030] N=1,274	-0.039 [0.033] N=1,137	-0.006 [0.037] N=965	-0.026 [0.040] N=782	-0.049 [0.046] N=592
Wasted WHZ below -2	**0.087** [0.043] N=1,101**	0.056 [0.049] N=978	**0.119** [0.051] N=838**	**0.102* [0.053] N=681**	**0.170*** [0.063] N=520**
Overweight WHZ above +2	0.007 [0.065] N=1,101	0.008 [0.069] N=978	0.031 [0.078] N=838	-0.012 [0.100] N=681	-0.037 [0.121] N=520
Obese WHZ above +3	0.000 [0.048] N=1,101	-0.002 [0.050] N=978	-0.011 [0.058] N=838	-0.012 [0.076] N=681	-0.068 [0.086] N=520
CIAF any stunting underweight or wasting	0.076 [0.078] N=1,084	0.030 [0.089] N=961	0.085 [0.102] N=822	0.042 [0.118] N=667	0.137 [0.138] N=509
Below-normal BMI-for-age BMIZ below -2	**0.088** [0.042] N=1,099**	0.051 [0.048] N=977	**0.120** [0.050] N=837**	**0.131** [0.053] N=679**	**0.196*** [0.059] N=525**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. UNICEF child-age-interval fixed effects are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Neither aid control is combined with an income control. Aid specifications use only the six main questions; households with a blank or 99 on any question are excluded. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household; all complete-DOB development respondents are included; raw ages below 24 map to 24 and above 59 map to 59. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.