Table 15 Reduced Form Effects on WHO Anthropometric Outcomes for the Oldest Eligible Child With Monthly Child Age Controls Using 16 December 2022 Child Age

Continuous z scores and prespecified adverse clinical classifications exclude WHO-flagged measurements. The earliest-born eligible child is selected per household using age on 16 December 2022. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.487	0.126 [0.211] N=1,525	0.175 [0.220] N=1,411	0.196 [0.230] N=1,268	0.180 [0.250] N=1,083	0.160 [0.283] N=891
WHO weight-for-age z score	0.038	0.093 [0.124] N=1,677	0.120 [0.134] N=1,548	0.054 [0.136] N=1,393	0.072 [0.154] N=1,188	0.020 [0.156] N=977
WHO weight-for-height z score	0.364	-0.082 [0.201] N=1,437	-0.095 [0.208] N=1,337	-0.257 [0.224] N=1,200	-0.292 [0.251] N=1,027	-0.210 [0.264] N=842
WHO BMI-for-age z score	0.355	-0.095 [0.221] N=1,448	-0.084 [0.234] N=1,345	-0.242 [0.251] N=1,202	-0.352 [0.290] N=1,025	-0.292 [0.310] N=845
Stunted HAZ below -2	0.236 P=351	-0.036 [0.039] N=1,525	-0.048 [0.040] N=1,411	-0.044 [0.042] N=1,268	-0.044 [0.046] N=1,083	-0.006 [0.050] N=891
Underweight WAZ below -2	0.061 P=102	-0.024 [0.020] N=1,677	-0.027 [0.021] N=1,548	-0.022 [0.022] N=1,393	-0.019 [0.023] N=1,188	-0.007 [0.025] N=977
Wasted WHZ below -2	0.106 P=158	-0.001 [0.033] N=1,437	0.010 [0.033] N=1,337	0.032 [0.037] N=1,200	0.020 [0.042] N=1,027	0.003 [0.045] N=842
Overweight WHZ above +2	0.196 P=272	0.014 [0.044] N=1,437	0.015 [0.046] N=1,337	0.004 [0.049] N=1,200	-0.013 [0.053] N=1,027	-0.008 [0.060] N=842
Obese WHZ above +3	0.089 P=127	-0.003 [0.030] N=1,437	-0.007 [0.031] N=1,337	-0.010 [0.033] N=1,200	-0.015 [0.037] N=1,027	-0.020 [0.042] N=842
CIAF any stunting underweight or wasting	0.315 P=443	-0.056 [0.041] N=1,415	-0.057 [0.042] N=1,315	-0.030 [0.046] N=1,180	-0.038 [0.048] N=1,008	-0.010 [0.052] N=825
Below-normal BMI-for-age BMIZ below -2	0.109 P=164	0.014 [0.036] N=1,448	0.020 [0.037] N=1,345	0.045 [0.041] N=1,202	0.046 [0.046] N=1,025	0.046 [0.050] N=845

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.035 [0.311] N=1,292	0.012 [0.346] N=1,178	0.024 [0.376] N=1,035	-0.009 [0.429] N=850	-0.160 [0.580] N=658
WHO weight-for-age z score	0.094 [0.180] N=1,423	0.137 [0.199] N=1,294	0.005 [0.209] N=1,139	0.040 [0.272] N=934	-0.113 [0.321] N=723
WHO weight-for-height z score	-0.040 [0.276] N=1,218	-0.058 [0.299] N=1,118	-0.378 [0.355] N=981	-0.476 [0.437] N=808	-0.338 [0.530] N=623
WHO BMI-for-age z score	0.096 [0.286] N=1,229	0.150 [0.316] N=1,126	-0.093 [0.368] N=983	-0.319 [0.464] N=806	-0.127 [0.555] N=626
Stunted HAZ below -2	-0.029 [0.059] N=1,292	-0.050 [0.066] N=1,178	-0.048 [0.070] N=1,035	-0.059 [0.083] N=850	0.034 [0.108] N=658
Underweight WAZ below -2	**-0.050* [0.030] N=1,423**	**-0.060* [0.033] N=1,294**	-0.054 [0.037] N=1,139	-0.058 [0.040] N=934	-0.029 [0.053] N=723
Wasted WHZ below -2	0.012 [0.036] N=1,218	0.033 [0.036] N=1,118	**0.078* [0.045] N=981**	0.064 [0.056] N=808	0.039 [0.060] N=623
Overweight WHZ above +2	0.032 [0.058] N=1,218	0.035 [0.062] N=1,118	0.013 [0.070] N=981	-0.015 [0.084] N=808	0.009 [0.110] N=623
Obese WHZ above +3	0.016 [0.042] N=1,218	0.007 [0.046] N=1,118	0.009 [0.053] N=981	0.004 [0.066] N=808	0.009 [0.094] N=623
CIAF any stunting underweight or wasting	-0.034 [0.060] N=1,203	-0.033 [0.065] N=1,103	0.017 [0.074] N=968	0.005 [0.086] N=796	0.097 [0.105] N=613
Below-normal BMI-for-age BMIZ below -2	-0.005 [0.038] N=1,229	0.002 [0.039] N=1,126	0.041 [0.046] N=983	0.042 [0.054] N=806	0.039 [0.064] N=626

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.487	0.128 [0.211] N=1,525	0.178 [0.220] N=1,411	0.201 [0.229] N=1,268	0.186 [0.248] N=1,083	0.165 [0.279] N=891
WHO weight-for-age z score	0.038	0.095 [0.124] N=1,677	0.121 [0.134] N=1,548	0.055 [0.136] N=1,393	0.073 [0.154] N=1,188	0.020 [0.157] N=977
WHO weight-for-height z score	0.364	-0.083 [0.201] N=1,437	-0.098 [0.208] N=1,337	-0.262 [0.224] N=1,200	-0.292 [0.250] N=1,027	-0.212 [0.264] N=842
WHO BMI-for-age z score	0.355	-0.097 [0.220] N=1,448	-0.088 [0.233] N=1,345	-0.246 [0.250] N=1,202	-0.351 [0.288] N=1,025	-0.294 [0.306] N=845
Stunted HAZ below -2	0.236 P=351	-0.036 [0.039] N=1,525	-0.047 [0.040] N=1,411	-0.045 [0.042] N=1,268	-0.044 [0.047] N=1,083	-0.007 [0.050] N=891
Underweight WAZ below -2	0.061 P=102	-0.024 [0.020] N=1,677	-0.027 [0.021] N=1,548	-0.022 [0.022] N=1,393	-0.019 [0.023] N=1,188	-0.007 [0.025] N=977
Wasted WHZ below -2	0.106 P=158	0.000 [0.032] N=1,437	0.011 [0.033] N=1,337	0.033 [0.036] N=1,200	0.020 [0.041] N=1,027	0.004 [0.044] N=842
Overweight WHZ above +2	0.196 P=272	0.014 [0.044] N=1,437	0.015 [0.045] N=1,337	0.005 [0.049] N=1,200	-0.013 [0.053] N=1,027	-0.008 [0.060] N=842
Obese WHZ above +3	0.089 P=127	-0.003 [0.030] N=1,437	-0.007 [0.031] N=1,337	-0.009 [0.033] N=1,200	-0.015 [0.037] N=1,027	-0.020 [0.042] N=842
CIAF any stunting underweight or wasting	0.315 P=443	-0.056 [0.041] N=1,415	-0.055 [0.042] N=1,315	-0.029 [0.045] N=1,180	-0.037 [0.047] N=1,008	-0.009 [0.052] N=825
Below-normal BMI-for-age BMIZ below -2	0.109 P=164	0.014 [0.036] N=1,448	0.021 [0.037] N=1,345	0.046 [0.040] N=1,202	0.045 [0.045] N=1,025	0.046 [0.049] N=845

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	**8 years**	**7 years**	**6 years**	**5 years**	**4 years**
WHO height-for-age z score	-0.031 [0.310] N=1,292	0.019 [0.344] N=1,178	0.040 [0.373] N=1,035	0.012 [0.420] N=850	-0.134 [0.566] N=658
WHO weight-for-age z score	0.096 [0.180] N=1,423	0.138 [0.199] N=1,294	0.007 [0.210] N=1,139	0.042 [0.271] N=934	-0.113 [0.320] N=723
WHO weight-for-height z score	-0.045 [0.277] N=1,218	-0.068 [0.300] N=1,118	-0.395 [0.356] N=981	-0.485 [0.438] N=808	-0.369 [0.535] N=623
WHO BMI-for-age z score	0.091 [0.287] N=1,229	0.139 [0.317] N=1,126	-0.110 [0.368] N=983	-0.327 [0.465] N=806	-0.156 [0.559] N=626
Stunted HAZ below -2	-0.029 [0.059] N=1,292	-0.050 [0.066] N=1,178	-0.049 [0.070] N=1,035	-0.060 [0.083] N=850	0.031 [0.107] N=658
Underweight WAZ below -2	**-0.050* [0.030]** **N=1,423**	**-0.060* [0.033]** **N=1,294**	-0.054 [0.037] N=1,139	-0.058 [0.040] N=934	-0.029 [0.053] N=723
Wasted WHZ below -2	0.013 [0.037] N=1,218	0.036 [0.037] N=1,118	**0.082* [0.046]** **N=981**	0.067 [0.056] N=808	0.046 [0.062] N=623
Overweight WHZ above +2	0.032 [0.058] N=1,218	0.036 [0.062] N=1,118	0.014 [0.070] N=981	-0.014 [0.084] N=808	0.012 [0.110] N=623
Obese WHZ above +3	0.016 [0.042] N=1,218	0.008 [0.046] N=1,118	0.010 [0.053] N=981	0.005 [0.067] N=808	0.010 [0.094] N=623
CIAF any stunting underweight or wasting	-0.033 [0.060] N=1,203	-0.030 [0.065] N=1,103	0.020 [0.074] N=968	0.007 [0.086] N=796	0.101 [0.105] N=613
Below-normal BMI-for-age BMIZ below -2	-0.004 [0.038] N=1,229	0.004 [0.039] N=1,126	0.044 [0.046] N=983	0.044 [0.054] N=806	0.045 [0.066] N=626

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.487	0.076 [0.208] N=1,525	0.110 [0.217] N=1,411	0.152 [0.228] N=1,268	0.115 [0.247] N=1,083	0.087 [0.276] N=891
WHO weight-for-age z score	0.038	0.032 [0.122] N=1,677	0.057 [0.132] N=1,548	-0.005 [0.136] N=1,393	0.002 [0.154] N=1,188	-0.056 [0.160] N=977
WHO weight-for-height z score	0.364	-0.126 [0.198] N=1,437	-0.137 [0.206] N=1,337	-0.298 [0.221] N=1,200	-0.330 [0.247] N=1,027	-0.250 [0.263] N=842
WHO BMI-for-age z score	0.355	-0.124 [0.218] N=1,448	-0.113 [0.231] N=1,345	-0.271 [0.247] N=1,202	-0.378 [0.284] N=1,025	-0.314 [0.307] N=845
Stunted HAZ below -2	0.236 P=351	-0.024 [0.037] N=1,525	-0.032 [0.039] N=1,411	-0.033 [0.040] N=1,268	-0.031 [0.045] N=1,083	0.008 [0.048] N=891
Underweight WAZ below -2	0.061 P=102	-0.014 [0.020] N=1,677	-0.014 [0.021] N=1,548	-0.010 [0.022] N=1,393	-0.006 [0.023] N=1,188	0.008 [0.025] N=977
Wasted WHZ below -2	0.106 P=158	0.010 [0.032] N=1,437	0.023 [0.032] N=1,337	0.043 [0.035] N=1,200	0.029 [0.039] N=1,027	0.012 [0.043] N=842
Overweight WHZ above +2	0.196 P=272	0.012 [0.044] N=1,437	0.013 [0.046] N=1,337	0.002 [0.049] N=1,200	-0.017 [0.054] N=1,027	-0.012 [0.060] N=842
Obese WHZ above +3	0.089 P=127	-0.003 [0.030] N=1,437	-0.006 [0.031] N=1,337	-0.009 [0.033] N=1,200	-0.017 [0.036] N=1,027	-0.024 [0.043] N=842
CIAF any stunting underweight or wasting	0.315 P=443	-0.035 [0.039] N=1,415	-0.030 [0.040] N=1,315	-0.010 [0.042] N=1,180	-0.018 [0.044] N=1,008	0.013 [0.047] N=825
Below-normal BMI-for-age BMIZ below -2	0.109 P=164	0.023 [0.035] N=1,448	0.033 [0.036] N=1,345	0.056 [0.039] N=1,202	0.056 [0.044] N=1,025	0.056 [0.048] N=845

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.065 [0.304] N=1,292	-0.037 [0.337] N=1,178	0.007 [0.370] N=1,035	-0.058 [0.415] N=850	-0.204 [0.550] N=658
WHO weight-for-age z score	0.040 [0.175] N=1,423	0.076 [0.193] N=1,294	-0.048 [0.206] N=1,139	-0.031 [0.267] N=934	-0.196 [0.325] N=723
WHO weight-for-height z score	-0.072 [0.275] N=1,218	-0.089 [0.299] N=1,118	-0.411 [0.354] N=981	-0.507 [0.442] N=808	-0.406 [0.542] N=623
WHO BMI-for-age z score	0.080 [0.285] N=1,229	0.128 [0.315] N=1,126	-0.118 [0.366] N=983	-0.356 [0.469] N=806	-0.176 [0.573] N=626
Stunted HAZ below -2	-0.023 [0.057] N=1,292	-0.038 [0.063] N=1,178	-0.043 [0.068] N=1,035	-0.051 [0.081] N=850	0.037 [0.105] N=658
Underweight WAZ below -2	-0.041 [0.030] N=1,423	-0.046 [0.033] N=1,294	-0.043 [0.037] N=1,139	-0.046 [0.039] N=934	-0.016 [0.053] N=723
Wasted WHZ below -2	0.021 [0.036] N=1,218	0.046 [0.037] N=1,118	**0.089** [0.045] N=981**	0.077 [0.056] N=808	0.052 [0.063] N=623
Overweight WHZ above +2	0.034 [0.058] N=1,218	0.040 [0.063] N=1,118	0.017 [0.071] N=981	-0.013 [0.085] N=808	0.011 [0.111] N=623
Obese WHZ above +3	0.018 [0.042] N=1,218	0.012 [0.045] N=1,118	0.011 [0.053] N=981	0.005 [0.066] N=808	0.000 [0.094] N=623
CIAF any stunting underweight or wasting	-0.018 [0.059] N=1,203	-0.008 [0.063] N=1,103	0.032 [0.072] N=968	0.025 [0.083] N=796	0.114 [0.103] N=613
Below-normal BMI-for-age BMIZ below -2	0.001 [0.038] N=1,229	0.014 [0.039] N=1,126	0.050 [0.045] N=983	0.056 [0.054] N=806	0.050 [0.067] N=626

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.487	0.030 [0.207] N=1,525	0.060 [0.217] N=1,411	0.106 [0.232] N=1,268	0.067 [0.249] N=1,083	0.018 [0.279] N=891
WHO weight-for-age z score	0.038	0.040 [0.121] N=1,677	0.059 [0.130] N=1,548	-0.001 [0.136] N=1,393	0.001 [0.155] N=1,188	-0.074 [0.157] N=977
WHO weight-for-height z score	0.364	-0.050 [0.200] N=1,437	-0.059 [0.207] N=1,337	-0.233 [0.224] N=1,200	-0.275 [0.253] N=1,027	-0.180 [0.267] N=842
WHO BMI-for-age z score	0.355	-0.061 [0.221] N=1,448	-0.045 [0.233] N=1,345	-0.214 [0.252] N=1,202	-0.324 [0.294] N=1,025	-0.259 [0.316] N=845
Stunted HAZ below -2	0.236 P=351	-0.016 [0.038] N=1,525	-0.024 [0.039] N=1,411	-0.026 [0.041] N=1,268	-0.020 [0.045] N=1,083	0.019 [0.048] N=891
Underweight WAZ below -2	0.061 P=102	-0.014 [0.021] N=1,677	-0.014 [0.022] N=1,548	-0.009 [0.023] N=1,393	-0.004 [0.024] N=1,188	0.011 [0.026] N=977
Wasted WHZ below -2	0.106 P=158	0.005 [0.032] N=1,437	0.018 [0.033] N=1,337	0.040 [0.036] N=1,200	0.029 [0.041] N=1,027	0.010 [0.044] N=842
Overweight WHZ above +2	0.196 P=272	0.021 [0.044] N=1,437	0.021 [0.046] N=1,337	0.008 [0.049] N=1,200	-0.010 [0.055] N=1,027	-0.005 [0.061] N=842
Obese WHZ above +3	0.089 P=127	0.006 [0.030] N=1,437	0.003 [0.031] N=1,337	-0.001 [0.034] N=1,200	-0.005 [0.038] N=1,027	-0.008 [0.044] N=842
CIAF any stunting underweight or wasting	0.315 P=443	-0.030 [0.040] N=1,415	-0.026 [0.041] N=1,315	-0.004 [0.045] N=1,180	-0.005 [0.045] N=1,008	0.023 [0.050] N=825
Below-normal BMI-for-age BMIZ below -2	0.109 P=164	0.018 [0.036] N=1,448	0.027 [0.037] N=1,345	0.052 [0.040] N=1,202	0.053 [0.046] N=1,025	0.051 [0.050] N=845

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	**8 years**	**7 years**	**6 years**	**5 years**	**4 years**
WHO height-for-age z score	-0.089 [0.300] N=1,292	-0.073 [0.330] N=1,178	-0.014 [0.367] N=1,035	-0.064 [0.403] N=850	-0.262 [0.531] N=658
WHO weight-for-age z score	0.048 [0.176] N=1,423	0.084 [0.194] N=1,294	-0.043 [0.209] N=1,139	-0.025 [0.272] N=934	-0.184 [0.329] N=723
WHO weight-for-height z score	-0.043 [0.277] N=1,218	-0.035 [0.299] N=1,118	-0.377 [0.356] N=981	-0.496 [0.443] N=808	-0.354 [0.538] N=623
WHO BMI-for-age z score	0.096 [0.287] N=1,229	0.173 [0.315] N=1,126	-0.094 [0.370] N=983	-0.340 [0.467] N=806	-0.129 [0.566] N=626
Stunted HAZ below -2	-0.016 [0.058] N=1,292	-0.031 [0.064] N=1,178	-0.039 [0.070] N=1,035	-0.047 [0.082] N=850	0.058 [0.106] N=658
Underweight WAZ below -2	-0.040 [0.030] N=1,423	-0.046 [0.033] N=1,294	-0.040 [0.037] N=1,139	-0.039 [0.040] N=934	-0.008 [0.055] N=723
Wasted WHZ below -2	0.022 [0.036] N=1,218	0.044 [0.037] N=1,118	**0.089* [0.046] N=981**	0.079 [0.056] N=808	0.049 [0.063] N=623
Overweight WHZ above +2	0.039 [0.059] N=1,218	0.046 [0.063] N=1,118	0.021 [0.071] N=981	-0.007 [0.086] N=808	0.024 [0.111] N=623
Obese WHZ above +3	0.022 [0.042] N=1,218	0.019 [0.046] N=1,118	0.018 [0.053] N=981	0.012 [0.066] N=808	0.013 [0.093] N=623
CIAF any stunting underweight or wasting	-0.008 [0.059] N=1,203	-0.002 [0.064] N=1,103	0.038 [0.073] N=968	0.032 [0.082] N=796	0.132 [0.103] N=613
Below-normal BMI-for-age BMIZ below -2	0.003 [0.038] N=1,229	0.010 [0.040] N=1,126	0.050 [0.046] N=983	0.055 [0.055] N=806	0.047 [0.066] N=626

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.495	0.078 [0.215] N=1,436	0.114 [0.224] N=1,334	0.149 [0.228] N=1,196	0.188 [0.236] N=1,024	0.140 [0.272] N=839
WHO weight-for-age z score	0.027	0.094 [0.128] N=1,579	0.115 [0.136] N=1,463	0.049 [0.143] N=1,315	0.067 [0.164] N=1,125	0.008 [0.171] N=922
WHO weight-for-height z score	0.369	-0.015 [0.211] N=1,355	-0.038 [0.217] N=1,267	-0.213 [0.234] N=1,136	-0.257 [0.260] N=975	-0.158 [0.283] N=797
WHO BMI-for-age z score	0.367	-0.032 [0.225] N=1,363	-0.033 [0.237] N=1,272	-0.182 [0.259] N=1,136	-0.301 [0.295] N=971	-0.223 [0.322] N=798
Stunted HAZ below -2	0.237 P=330	-0.018 [0.041] N=1,436	-0.030 [0.043] N=1,334	-0.029 [0.043] N=1,196	-0.042 [0.047] N=1,024	0.001 [0.051] N=839
Underweight WAZ below -2	0.063 P=100	-0.023 [0.021] N=1,579	-0.025 [0.022] N=1,463	-0.019 [0.024] N=1,315	-0.019 [0.025] N=1,125	-0.006 [0.028] N=922
Wasted WHZ below -2	0.106 P=150	0.004 [0.035] N=1,355	0.013 [0.035] N=1,267	0.040 [0.039] N=1,136	0.026 [0.044] N=975	0.008 [0.048] N=797
Overweight WHZ above +2	0.198 P=257	0.018 [0.046] N=1,355	0.015 [0.048] N=1,267	0.003 [0.052] N=1,136	-0.020 [0.056] N=975	-0.015 [0.064] N=797
Obese WHZ above +3	0.092 P=123	0.002 [0.031] N=1,355	-0.004 [0.033] N=1,267	-0.007 [0.035] N=1,136	-0.015 [0.038] N=975	-0.020 [0.045] N=797
CIAF any stunting underweight or wasting	0.317 P=419	-0.032 [0.046] N=1,334	-0.035 [0.047] N=1,246	-0.008 [0.050] N=1,116	-0.028 [0.053] N=956	0.004 [0.059] N=780
Below-normal BMI-for-age BMIZ below -2	0.108 P=153	0.016 [0.036] N=1,363	0.021 [0.037] N=1,272	0.048 [0.041] N=1,136	0.047 [0.046] N=971	0.046 [0.050] N=798

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.093 [0.324] N=1,218	-0.077 [0.361] N=1,116	-0.041 [0.385] N=978	0.069 [0.423] N=806	-0.146 [0.599] N=621
WHO weight-for-age z score	0.092 [0.183] N=1,341	0.122 [0.196] N=1,225	-0.022 [0.217] N=1,077	0.020 [0.284] N=887	-0.178 [0.352] N=684
WHO weight-for-height z score	0.031 [0.282] N=1,148	-0.005 [0.301] N=1,060	-0.344 [0.355] N=929	-0.458 [0.438] N=768	-0.274 [0.547] N=590
WHO BMI-for-age z score	0.124 [0.289] N=1,158	0.154 [0.316] N=1,067	-0.080 [0.375] N=931	-0.320 [0.465] N=766	-0.078 [0.570] N=593
Stunted HAZ below -2	-0.004 [0.061] N=1,218	-0.021 [0.068] N=1,116	-0.023 [0.072] N=978	-0.061 [0.082] N=806	0.041 [0.109] N=621
Underweight WAZ below -2	**-0.052* [0.030] N=1,341**	**-0.060* [0.032] N=1,225**	-0.050 [0.038] N=1,077	-0.056 [0.041] N=887	-0.024 [0.057] N=684
Wasted WHZ below -2	0.013 [0.037] N=1,148	0.031 [0.036] N=1,060	**0.088* [0.045] N=929**	0.067 [0.057] N=768	0.036 [0.063] N=590
Overweight WHZ above +2	0.042 [0.060] N=1,148	0.039 [0.064] N=1,060	0.023 [0.072] N=929	-0.021 [0.086] N=768	0.021 [0.118] N=590
Obese WHZ above +3	0.027 [0.044] N=1,148	0.016 [0.048] N=1,060	0.017 [0.055] N=929	0.007 [0.068] N=768	0.022 [0.101] N=590
CIAF any stunting underweight or wasting	-0.000 [0.064] N=1,134	-0.002 [0.068] N=1,046	0.054 [0.079] N=916	0.015 [0.091] N=756	0.107 [0.114] N=580
Below-normal BMI-for-age BMIZ below -2	0.001 [0.038] N=1,158	0.006 [0.037] N=1,067	0.051 [0.045] N=931	0.049 [0.054] N=766	0.039 [0.065] N=593

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.495	0.074 [0.214] N=1,436	0.111 [0.223] N=1,334	0.117 [0.230] N=1,196	0.142 [0.244] N=1,024	0.069 [0.282] N=839
WHO weight-for-age z score	0.027	0.094 [0.129] N=1,579	0.115 [0.138] N=1,463	0.045 [0.142] N=1,315	0.070 [0.162] N=1,125	0.006 [0.168] N=922
WHO weight-for-height z score	0.369	-0.014 [0.210] N=1,355	-0.036 [0.216] N=1,267	-0.208 [0.232] N=1,136	-0.246 [0.259] N=975	-0.145 [0.280] N=797
WHO BMI-for-age z score	0.367	-0.033 [0.225] N=1,363	-0.033 [0.236] N=1,272	-0.181 [0.257] N=1,136	-0.288 [0.296] N=971	-0.207 [0.321] N=798
Stunted HAZ below -2	0.237 P=330	-0.019 [0.041] N=1,436	-0.030 [0.043] N=1,334	-0.027 [0.044] N=1,196	-0.037 [0.048] N=1,024	0.005 [0.053] N=839
Underweight WAZ below -2	0.063 P=100	-0.025 [0.022] N=1,579	-0.027 [0.023] N=1,463	-0.021 [0.024] N=1,315	-0.021 [0.025] N=1,125	-0.007 [0.028] N=922
Wasted WHZ below -2	0.106 P=150	0.004 [0.034] N=1,355	0.013 [0.035] N=1,267	0.039 [0.039] N=1,136	0.024 [0.045] N=975	0.002 [0.048] N=797
Overweight WHZ above +2	0.198 P=257	0.018 [0.046] N=1,355	0.015 [0.047] N=1,267	0.006 [0.051] N=1,136	-0.017 [0.055] N=975	-0.010 [0.063] N=797
Obese WHZ above +3	0.092 P=123	0.002 [0.031] N=1,355	-0.003 [0.032] N=1,267	-0.005 [0.034] N=1,136	-0.013 [0.038] N=975	-0.016 [0.044] N=797
CIAF any stunting underweight or wasting	0.317 P=419	-0.032 [0.045] N=1,334	-0.036 [0.047] N=1,246	-0.008 [0.050] N=1,116	-0.029 [0.053] N=956	-0.006 [0.059] N=780
Below-normal BMI-for-age BMIZ below -2	0.108 P=153	0.015 [0.036] N=1,363	0.021 [0.037] N=1,272	0.047 [0.041] N=1,136	0.045 [0.047] N=971	0.039 [0.051] N=798

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.035 [0.325] N=1,218	-0.001 [0.359] N=1,116	-0.013 [0.390] N=978	0.087 [0.446] N=806	-0.175 [0.634] N=621
WHO weight-for-age z score	0.106 [0.183] N=1,341	0.140 [0.198] N=1,225	0.003 [0.214] N=1,077	0.056 [0.274] N=887	-0.130 [0.338] N=684
WHO weight-for-height z score	0.027 [0.281] N=1,148	-0.002 [0.300] N=1,060	-0.340 [0.355] N=929	-0.452 [0.442] N=768	-0.264 [0.553] N=590
WHO BMI-for-age z score	0.124 [0.290] N=1,158	0.157 [0.317] N=1,067	-0.078 [0.373] N=931	-0.315 [0.470] N=766	-0.069 [0.578] N=593
Stunted HAZ below -2	-0.012 [0.061] N=1,218	-0.032 [0.068] N=1,116	-0.029 [0.072] N=978	-0.069 [0.084] N=806	0.035 [0.112] N=621
Underweight WAZ below -2	**-0.054* [0.030] N=1,341**	**-0.063* [0.033] N=1,225**	-0.056 [0.038] N=1,077	-0.064 [0.040] N=887	-0.035 [0.055] N=684
Wasted WHZ below -2	0.016 [0.037] N=1,148	0.033 [0.036] N=1,060	**0.089* [0.045] N=929**	0.068 [0.057] N=768	0.030 [0.062] N=590
Overweight WHZ above +2	0.042 [0.059] N=1,148	0.040 [0.063] N=1,060	0.024 [0.072] N=929	-0.018 [0.086] N=768	0.025 [0.116] N=590
Obese WHZ above +3	0.025 [0.044] N=1,148	0.015 [0.048] N=1,060	0.019 [0.055] N=929	0.008 [0.069] N=768	0.025 [0.099] N=590
CIAF any stunting underweight or wasting	-0.001 [0.064] N=1,134	-0.004 [0.069] N=1,046	0.052 [0.079] N=916	0.012 [0.091] N=756	0.095 [0.112] N=580
Below-normal BMI-for-age BMIZ below -2	0.002 [0.037] N=1,158	0.008 [0.037] N=1,067	0.052 [0.045] N=931	0.048 [0.055] N=766	0.032 [0.066] N=593

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.469	-0.092 [0.238] N=1,416	-0.093 [0.254] N=1,279	-0.092 [0.275] N=1,120	-0.173 [0.296] N=948	-0.191 [0.322] N=771
WHO weight-for-age z score	0.031	0.082 [0.136] N=1,550	0.158 [0.146] N=1,409	0.028 [0.147] N=1,227	0.005 [0.157] N=1,033	0.019 [0.168] N=829
WHO weight-for-height z score	0.383	0.037 [0.196] N=1,340	0.061 [0.201] N=1,213	0.027 [0.215] N=1,065	-0.004 [0.240] N=898	-0.022 [0.271] N=726
WHO BMI-for-age z score	0.350	-0.042 [0.225] N=1,343	-0.036 [0.237] N=1,215	-0.143 [0.254] N=1,067	-0.162 [0.276] N=899	-0.060 [0.305] N=732
Stunted HAZ below -2	0.235 P=333	0.041 [0.046] N=1,416	0.043 [0.049] N=1,279	0.042 [0.049] N=1,120	0.056 [0.053] N=948	0.047 [0.059] N=771
Underweight WAZ below -2	0.059 P=92	-0.021 [0.026] N=1,550	-0.038 [0.028] N=1,409	-0.018 [0.029] N=1,227	-0.023 [0.031] N=1,033	-0.023 [0.035] N=829
Wasted WHZ below -2	0.104 P=139	0.040 [0.033] N=1,340	0.027 [0.036] N=1,213	0.053 [0.037] N=1,065	0.036 [0.038] N=898	0.041 [0.044] N=726
Overweight WHZ above +2	0.192 P=261	0.025 [0.042] N=1,340	0.028 [0.043] N=1,213	0.047 [0.046] N=1,065	0.042 [0.051] N=898	0.049 [0.054] N=726
Obese WHZ above +3	0.094 P=131	0.020 [0.031] N=1,340	0.022 [0.031] N=1,213	0.017 [0.035] N=1,065	0.018 [0.041] N=898	0.006 [0.042] N=726
CIAF any stunting underweight or wasting	0.315 P=414	0.085 [0.054] N=1,317	0.059 [0.059] N=1,190	0.092 [0.060] N=1,043	0.086 [0.064] N=879	0.114 [0.071] N=712
Below-normal BMI-for-age BMIZ below -2	0.105 P=142	0.032 [0.038] N=1,343	0.026 [0.042] N=1,215	0.052 [0.044] N=1,067	0.039 [0.047] N=899	0.034 [0.052] N=732

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.174 [0.319] N=1,225	0.224 [0.350] N=1,088	0.324 [0.407] N=929	0.243 [0.507] N=757	0.539 [0.644] N=580
WHO weight-for-age z score	0.201 [0.202] N=1,346	0.349 [0.218] N=1,205	0.122 [0.231] N=1,023	0.129 [0.278] N=829	0.238 [0.359] N=625
WHO weight-for-height z score	-0.150 [0.291] N=1,165	-0.131 [0.297] N=1,038	-0.236 [0.337] N=890	-0.340 [0.420] N=723	-0.694 [0.587] N=551
WHO BMI-for-age z score	-0.358 [0.292] N=1,169	-0.391 [0.315] N=1,041	**-0.682* [0.357] N=893**	**-0.793* [0.449] N=725**	-0.943 [0.616] N=558
Stunted HAZ below -2	0.002 [0.070] N=1,225	-0.006 [0.078] N=1,088	-0.026 [0.085] N=929	-0.015 [0.107] N=757	-0.087 [0.134] N=580
Underweight WAZ below -2	-0.049 [0.036] N=1,346	**-0.080** [0.039] N=1,205**	-0.047 [0.043] N=1,023	-0.078 [0.048] N=829	**-0.115** [0.052] N=625**
Wasted WHZ below -2	**0.077* [0.041] N=1,165**	0.058 [0.047] N=1,038	**0.119** [0.050] N=890**	**0.096** [0.048] N=723**	**0.153** [0.061] N=551**
Overweight WHZ above +2	-0.004 [0.065] N=1,165	-0.005 [0.068] N=1,038	0.023 [0.077] N=890	0.000 [0.101] N=723	-0.049 [0.129] N=551
Obese WHZ above +3	-0.001 [0.046] N=1,165	-0.004 [0.048] N=1,038	-0.018 [0.055] N=890	-0.024 [0.070] N=723	-0.115 [0.087] N=551
CIAF any stunting underweight or wasting	0.059 [0.075] N=1,146	0.005 [0.084] N=1,019	0.054 [0.091] N=872	0.015 [0.107] N=708	0.045 [0.132] N=541
Below-normal BMI-for-age BMIZ below -2	**0.083** [0.039] N=1,169**	**0.082* [0.046] N=1,041**	**0.145*** [0.049] N=893**	**0.136*** [0.050] N=725**	**0.187*** [0.062] N=558**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	**Mean**	**8 years**	**7 years**	**6 years**	**5 years**	**4 years**
WHO height-for-age z score	-0.469	-0.095 [0.237] N=1,416	-0.095 [0.253] N=1,279	-0.093 [0.274] N=1,120	-0.172 [0.295] N=948	-0.167 [0.322] N=771
WHO weight-for-age z score	0.031	0.082 [0.136] N=1,550	0.158 [0.145] N=1,409	0.029 [0.147] N=1,227	0.007 [0.158] N=1,033	0.020 [0.170] N=829
WHO weight-for-height z score	0.383	0.040 [0.196] N=1,340	0.060 [0.202] N=1,213	0.027 [0.216] N=1,065	-0.009 [0.241] N=898	-0.055 [0.274] N=726
WHO BMI-for-age z score	0.350	-0.042 [0.226] N=1,343	-0.039 [0.238] N=1,215	-0.146 [0.255] N=1,067	-0.168 [0.278] N=899	-0.097 [0.309] N=732
Stunted HAZ below -2	0.235 P=333	0.041 [0.046] N=1,416	0.043 [0.049] N=1,279	0.042 [0.049] N=1,120	0.056 [0.053] N=948	0.047 [0.060] N=771
Underweight WAZ below -2	0.059 P=92	-0.022 [0.026] N=1,550	-0.038 [0.028] N=1,409	-0.017 [0.029] N=1,227	-0.022 [0.031] N=1,033	-0.021 [0.035] N=829
Wasted WHZ below -2	0.104 P=139	0.039 [0.033] N=1,340	0.027 [0.036] N=1,213	0.053 [0.037] N=1,065	0.038 [0.039] N=898	0.049 [0.044] N=726
Overweight WHZ above +2	0.192 P=261	0.024 [0.042] N=1,340	0.028 [0.043] N=1,213	0.047 [0.046] N=1,065	0.043 [0.051] N=898	0.050 [0.055] N=726
Obese WHZ above +3	0.094 P=131	0.020 [0.031] N=1,340	0.022 [0.031] N=1,213	0.017 [0.035] N=1,065	0.019 [0.041] N=898	0.009 [0.042] N=726
CIAF any stunting underweight or wasting	0.315 P=414	0.085 [0.055] N=1,317	0.059 [0.060] N=1,190	0.092 [0.061] N=1,043	0.088 [0.066] N=879	**0.120* [0.072] N=712**
Below-normal BMI-for-age BMIZ below -2	0.105 P=142	0.032 [0.038] N=1,343	0.027 [0.042] N=1,215	0.053 [0.045] N=1,067	0.041 [0.048] N=899	0.041 [0.053] N=732

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.160 [0.316] N=1,225	0.207 [0.345] N=1,088	0.316 [0.405] N=929	0.209 [0.499] N=757	0.511 [0.634] N=580
WHO weight-for-age z score	0.200 [0.201] N=1,346	0.348 [0.216] N=1,205	0.120 [0.229] N=1,023	0.127 [0.276] N=829	0.238 [0.358] N=625
WHO weight-for-height z score	-0.128 [0.291] N=1,165	-0.109 [0.297] N=1,038	-0.220 [0.335] N=890	-0.313 [0.419] N=723	-0.668 [0.594] N=551
WHO BMI-for-age z score	-0.334 [0.293] N=1,169	-0.367 [0.315] N=1,041	**-0.666* [0.356] N=893**	**-0.764* [0.450] N=725**	-0.924 [0.625] N=558
Stunted HAZ below -2	0.003 [0.070] N=1,225	-0.006 [0.078] N=1,088	-0.029 [0.086] N=929	-0.017 [0.107] N=757	-0.086 [0.134] N=580
Underweight WAZ below -2	-0.049 [0.036] N=1,346	**-0.080** [0.039] N=1,205**	-0.047 [0.043] N=1,023	-0.078 [0.048] N=829	**-0.115** [0.052] N=625**
Wasted WHZ below -2	**0.072* [0.041] N=1,165**	0.052 [0.047] N=1,038	**0.111** [0.050] N=890**	**0.086* [0.048] N=723**	**0.147** [0.060] N=551**
Overweight WHZ above +2	-0.004 [0.065] N=1,165	-0.005 [0.068] N=1,038	0.022 [0.077] N=890	-0.001 [0.101] N=723	-0.049 [0.129] N=551
Obese WHZ above +3	-0.004 [0.046] N=1,165	-0.007 [0.047] N=1,038	-0.024 [0.054] N=890	-0.029 [0.070] N=723	-0.118 [0.087] N=551
CIAF any stunting underweight or wasting	0.054 [0.076] N=1,146	-0.002 [0.085] N=1,019	0.043 [0.094] N=872	0.004 [0.110] N=708	0.039 [0.136] N=541
Below-normal BMI-for-age BMIZ below -2	**0.078** [0.039] N=1,169**	**0.077* [0.046] N=1,041**	**0.140*** [0.049] N=893**	**0.129** [0.050] N=725**	**0.184*** [0.062] N=558**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.469	-0.115 [0.235] N=1,416	-0.100 [0.250] N=1,279	-0.096 [0.274] N=1,120	-0.165 [0.293] N=948	-0.186 [0.310] N=771
WHO weight-for-age z score	0.031	0.070 [0.137] N=1,550	0.156 [0.147] N=1,409	0.029 [0.149] N=1,227	0.004 [0.159] N=1,033	0.018 [0.167] N=829
WHO weight-for-height z score	0.383	0.035 [0.200] N=1,340	0.058 [0.206] N=1,213	0.035 [0.223] N=1,065	0.002 [0.248] N=898	-0.019 [0.281] N=726
WHO BMI-for-age z score	0.350	-0.042 [0.226] N=1,343	-0.036 [0.239] N=1,215	-0.133 [0.258] N=1,067	-0.158 [0.282] N=899	-0.071 [0.312] N=732
Stunted HAZ below -2	0.235 P=333	0.046 [0.045] N=1,416	0.044 [0.048] N=1,279	0.042 [0.049] N=1,120	0.054 [0.053] N=948	0.049 [0.058] N=771
Underweight WAZ below -2	0.059 P=92	-0.020 [0.026] N=1,550	-0.038 [0.028] N=1,409	-0.018 [0.029] N=1,227	-0.022 [0.031] N=1,033	-0.020 [0.034] N=829
Wasted WHZ below -2	0.104 P=139	0.039 [0.033] N=1,340	0.025 [0.036] N=1,213	0.050 [0.038] N=1,065	0.033 [0.039] N=898	0.040 [0.043] N=726
Overweight WHZ above +2	0.192 P=261	0.024 [0.043] N=1,340	0.027 [0.044] N=1,213	0.047 [0.047] N=1,065	0.041 [0.051] N=898	0.051 [0.055] N=726
Obese WHZ above +3	0.094 P=131	0.020 [0.031] N=1,340	0.022 [0.032] N=1,213	0.018 [0.036] N=1,065	0.019 [0.041] N=898	0.012 [0.043] N=726
CIAF any stunting underweight or wasting	0.315 P=414	0.088 [0.055] N=1,317	0.058 [0.060] N=1,190	0.090 [0.061] N=1,043	0.080 [0.066] N=879	0.113 [0.071] N=712
Below-normal BMI-for-age BMIZ below -2	0.105 P=142	0.032 [0.038] N=1,343	0.026 [0.043] N=1,215	0.051 [0.045] N=1,067	0.038 [0.048] N=899	0.038 [0.053] N=732

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.184 [0.315] N=1,225	0.251 [0.345] N=1,088	0.399 [0.408] N=929	0.321 [0.503] N=757	0.667 [0.619] N=580
WHO weight-for-age z score	0.196 [0.200] N=1,346	**0.363* [0.218] N=1,205**	0.135 [0.235] N=1,023	0.121 [0.285] N=829	0.211 [0.368] N=625
WHO weight-for-height z score	-0.163 [0.298] N=1,165	-0.141 [0.305] N=1,038	-0.273 [0.347] N=890	-0.400 [0.442] N=723	-0.784 [0.605] N=551
WHO BMI-for-age z score	-0.353 [0.295] N=1,169	-0.384 [0.321] N=1,041	**-0.711* [0.365] N=893**	**-0.854* [0.468] N=725**	**-1.048* [0.629] N=558**
Stunted HAZ below -2	-0.004 [0.071] N=1,225	-0.018 [0.079] N=1,088	-0.045 [0.087] N=929	-0.039 [0.108] N=757	-0.121 [0.133] N=580
Underweight WAZ below -2	-0.054 [0.036] N=1,346	**-0.089** [0.040] N=1,205**	-0.059 [0.044] N=1,023	**-0.089* [0.049] N=829**	**-0.120** [0.052] N=625**
Wasted WHZ below -2	**0.071* [0.042] N=1,165**	0.049 [0.048] N=1,038	**0.110** [0.051] N=890**	0.083 [0.051] N=723	**0.141** [0.062] N=551**
Overweight WHZ above +2	-0.010 [0.067] N=1,165	-0.010 [0.071] N=1,038	0.011 [0.081] N=890	-0.022 [0.107] N=723	-0.073 [0.137] N=551
Obese WHZ above +3	-0.010 [0.046] N=1,165	-0.014 [0.048] N=1,038	-0.032 [0.056] N=890	-0.040 [0.073] N=723	-0.133 [0.087] N=551
CIAF any stunting underweight or wasting	0.048 [0.078] N=1,146	-0.012 [0.088] N=1,019	0.032 [0.098] N=872	-0.014 [0.114] N=708	0.014 [0.140] N=541
Below-normal BMI-for-age BMIZ below -2	**0.075* [0.039] N=1,169**	0.072 [0.047] N=1,041	**0.138*** [0.048] N=893**	**0.125** [0.051] N=725**	**0.182*** [0.062] N=558**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.469	-0.109 [0.229] N=1,416	-0.090 [0.245] N=1,279	-0.083 [0.269] N=1,120	-0.162 [0.288] N=948	-0.163 [0.309] N=771
WHO weight-for-age z score	0.031	0.081 [0.136] N=1,550	0.167 [0.146] N=1,409	0.040 [0.147] N=1,227	0.017 [0.157] N=1,033	0.020 [0.165] N=829
WHO weight-for-height z score	0.383	0.043 [0.196] N=1,340	0.064 [0.200] N=1,213	0.033 [0.214] N=1,065	-0.004 [0.238] N=898	-0.053 [0.276] N=726
WHO BMI-for-age z score	0.350	-0.029 [0.222] N=1,343	-0.026 [0.234] N=1,215	-0.137 [0.250] N=1,067	-0.160 [0.270] N=899	-0.091 [0.302] N=732
Stunted HAZ below -2	0.235 P=333	0.046 [0.045] N=1,416	0.043 [0.048] N=1,279	0.041 [0.049] N=1,120	0.054 [0.053] N=948	0.043 [0.058] N=771
Underweight WAZ below -2	0.059 P=92	-0.020 [0.026] N=1,550	-0.039 [0.028] N=1,409	-0.019 [0.029] N=1,227	-0.024 [0.031] N=1,033	-0.023 [0.035] N=829
Wasted WHZ below -2	0.104 P=139	0.039 [0.032] N=1,340	0.025 [0.036] N=1,213	0.052 [0.037] N=1,065	0.038 [0.038] N=898	0.047 [0.043] N=726
Overweight WHZ above +2	0.192 P=261	0.026 [0.042] N=1,340	0.030 [0.043] N=1,213	0.049 [0.046] N=1,065	0.046 [0.051] N=898	0.050 [0.056] N=726
Obese WHZ above +3	0.094 P=131	0.022 [0.031] N=1,340	0.024 [0.031] N=1,213	0.019 [0.035] N=1,065	0.022 [0.040] N=898	0.012 [0.043] N=726
CIAF any stunting underweight or wasting	0.315 P=414	0.089 [0.055] N=1,317	0.059 [0.060] N=1,190	0.092 [0.061] N=1,043	0.088 [0.066] N=879	0.115 [0.071] N=712
Below-normal BMI-for-age BMIZ below -2	0.105 P=142	0.029 [0.038] N=1,343	0.024 [0.042] N=1,215	0.051 [0.044] N=1,067	0.039 [0.047] N=899	0.035 [0.052] N=732

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.166 [0.301] N=1,225	0.247 [0.329] N=1,088	0.364 [0.395] N=929	0.280 [0.482] N=757	0.582 [0.577] N=580
WHO weight-for-age z score	0.202 [0.193] N=1,346	**0.364* [0.211] N=1,205**	0.141 [0.221] N=1,023	0.154 [0.264] N=829	0.215 [0.330] N=625
WHO weight-for-height z score	-0.142 [0.289] N=1,165	-0.125 [0.291] N=1,038	-0.241 [0.323] N=890	-0.349 [0.406] N=723	-0.726 [0.576] N=551
WHO BMI-for-age z score	-0.337 [0.288] N=1,169	-0.376 [0.311] N=1,041	**-0.684* [0.348] N=893**	**-0.802* [0.441] N=725**	-0.967 [0.604] N=558
Stunted HAZ below -2	0.003 [0.069] N=1,225	-0.014 [0.077] N=1,088	-0.039 [0.086] N=929	-0.031 [0.105] N=757	-0.104 [0.131] N=580
Underweight WAZ below -2	-0.049 [0.035] N=1,346	**-0.084** [0.039] N=1,205**	-0.052 [0.042] N=1,023	**-0.085* [0.046] N=829**	**-0.115** [0.050] N=625**
Wasted WHZ below -2	**0.072* [0.042] N=1,165**	0.050 [0.048] N=1,038	**0.110** [0.050] N=890**	**0.085* [0.049] N=723**	**0.143** [0.061] N=551**
Overweight WHZ above +2	-0.008 [0.064] N=1,165	-0.009 [0.068] N=1,038	0.016 [0.076] N=890	-0.010 [0.100] N=723	-0.066 [0.128] N=551
Obese WHZ above +3	-0.005 [0.046] N=1,165	-0.009 [0.047] N=1,038	-0.026 [0.054] N=890	-0.033 [0.069] N=723	-0.123 [0.084] N=551
CIAF any stunting underweight or wasting	0.057 [0.077] N=1,146	-0.007 [0.087] N=1,019	0.036 [0.097] N=872	-0.004 [0.111] N=708	0.025 [0.134] N=541
Below-normal BMI-for-age BMIZ below -2	**0.076* [0.039] N=1,169**	0.073 [0.047] N=1,041	**0.138*** [0.048] N=893**	**0.126** [0.051] N=725**	**0.176*** [0.063] N=558**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.482	-0.162 [0.239] N=1,337	-0.156 [0.257] N=1,204	-0.161 [0.280] N=1,054	-0.237 [0.293] N=891	-0.284 [0.315] N=726
WHO weight-for-age z score	0.018	0.123 [0.139] N=1,464	0.212 [0.148] N=1,327	0.071 [0.149] N=1,155	0.041 [0.162] N=972	0.016 [0.173] N=782
WHO weight-for-height z score	0.382	0.104 [0.206] N=1,267	0.147 [0.212] N=1,144	0.094 [0.227] N=1,004	0.020 [0.257] N=847	-0.071 [0.287] N=686
WHO BMI-for-age z score	0.353	0.026 [0.231] N=1,267	0.047 [0.244] N=1,143	-0.084 [0.259] N=1,003	-0.156 [0.279] N=845	-0.125 [0.311] N=690
Stunted HAZ below -2	0.238 P=316	0.051 [0.047] N=1,337	0.054 [0.049] N=1,204	0.054 [0.050] N=1,054	0.068 [0.053] N=891	0.070 [0.061] N=726
Underweight WAZ below -2	0.061 P=90	-0.024 [0.027] N=1,464	-0.042 [0.029] N=1,327	-0.023 [0.030] N=1,155	-0.026 [0.033] N=972	-0.021 [0.037] N=782
Wasted WHZ below -2	0.106 P=134	0.042 [0.034] N=1,267	0.027 [0.037] N=1,144	0.053 [0.039] N=1,004	0.047 [0.041] N=847	0.062 [0.046] N=686
Overweight WHZ above +2	0.193 P=246	0.033 [0.046] N=1,267	0.041 [0.047] N=1,144	0.058 [0.049] N=1,004	0.049 [0.054] N=847	0.053 [0.058] N=686
Obese WHZ above +3	0.097 P=127	0.022 [0.033] N=1,267	0.026 [0.033] N=1,144	0.021 [0.037] N=1,004	0.027 [0.043] N=847	0.023 [0.045] N=686
CIAF any stunting underweight or wasting	0.320 P=396	0.093 [0.058] N=1,245	0.067 [0.062] N=1,122	0.101 [0.064] N=983	0.100 [0.069] N=828	**0.150* [0.076] N=672**
Below-normal BMI-for-age BMIZ below -2	0.105 P=134	0.037 [0.040] N=1,267	0.030 [0.044] N=1,143	0.057 [0.046] N=1,003	0.061 [0.049] N=845	0.067 [0.054] N=690

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.153 [0.335] N=1,157	0.234 [0.372] N=1,024	0.354 [0.448] N=874	0.261 [0.529] N=711	0.386 [0.677] N=546
WHO weight-for-age z score	0.263 [0.204] N=1,272	**0.440** [0.218] N=1,135**	0.197 [0.229] N=963	0.207 [0.281] N=780	0.220 [0.376] N=590
WHO weight-for-height z score	0.005 [0.301] N=1,100	0.057 [0.304] N=977	-0.073 [0.346] N=837	-0.215 [0.431] N=680	-0.644 [0.623] N=519
WHO BMI-for-age z score	-0.215 [0.302] N=1,102	-0.221 [0.328] N=978	-0.545 [0.375] N=838	-0.713 [0.463] N=680	-0.909 [0.661] N=525
Stunted HAZ below -2	-0.005 [0.073] N=1,157	-0.016 [0.081] N=1,024	-0.045 [0.090] N=874	-0.044 [0.110] N=711	-0.088 [0.138] N=546
Underweight WAZ below -2	-0.056 [0.037] N=1,272	**-0.092** [0.041] N=1,135**	-0.065 [0.043] N=963	**-0.096* [0.048] N=780**	**-0.125** [0.050] N=590**
Wasted WHZ below -2	0.067 [0.043] N=1,100	0.040 [0.049] N=977	**0.097* [0.053] N=837**	**0.089* [0.053] N=680**	**0.153** [0.067] N=519**
Overweight WHZ above +2	0.011 [0.069] N=1,100	0.016 [0.073] N=977	0.039 [0.082] N=837	0.009 [0.108] N=680	-0.041 [0.140] N=519
Obese WHZ above +3	-0.004 [0.048] N=1,100	-0.008 [0.050] N=977	-0.027 [0.056] N=837	-0.028 [0.073] N=680	-0.105 [0.093] N=519
CIAF any stunting underweight or wasting	0.046 [0.080] N=1,082	-0.015 [0.089] N=959	0.027 [0.100] N=820	-0.016 [0.117] N=665	0.057 [0.146] N=509
Below-normal BMI-for-age BMIZ below -2	**0.074* [0.040] N=1,102**	0.067 [0.048] N=978	**0.129** [0.051] N=838**	**0.143*** [0.053] N=680**	**0.198*** [0.067] N=525**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.482	-0.162 [0.242] N=1,337	-0.157 [0.259] N=1,204	-0.158 [0.282] N=1,054	-0.236 [0.300] N=891	-0.298 [0.317] N=726
WHO weight-for-age z score	0.018	0.126 [0.140] N=1,464	0.218 [0.150] N=1,327	0.072 [0.151] N=1,155	0.053 [0.165] N=972	0.023 [0.177] N=782
WHO weight-for-height z score	0.382	0.110 [0.203] N=1,267	0.149 [0.210] N=1,144	0.091 [0.224] N=1,004	0.034 [0.251] N=847	-0.052 [0.286] N=686
WHO BMI-for-age z score	0.353	0.029 [0.229] N=1,267	0.050 [0.243] N=1,143	-0.087 [0.258] N=1,003	-0.146 [0.278] N=845	-0.110 [0.309] N=690
Stunted HAZ below -2	0.238 P=316	0.050 [0.047] N=1,337	0.054 [0.050] N=1,204	0.053 [0.051] N=1,054	0.068 [0.055] N=891	0.072 [0.061] N=726
Underweight WAZ below -2	0.061 P=90	-0.024 [0.027] N=1,464	-0.042 [0.029] N=1,327	-0.021 [0.030] N=1,155	-0.025 [0.032] N=972	-0.020 [0.036] N=782
Wasted WHZ below -2	0.106 P=134	0.041 [0.034] N=1,267	0.026 [0.037] N=1,144	0.054 [0.039] N=1,004	0.044 [0.040] N=847	0.059 [0.045] N=686
Overweight WHZ above +2	0.193 P=246	0.032 [0.045] N=1,267	0.039 [0.046] N=1,144	0.055 [0.048] N=1,004	0.050 [0.053] N=847	0.054 [0.058] N=686
Obese WHZ above +3	0.097 P=127	0.023 [0.032] N=1,267	0.026 [0.033] N=1,144	0.020 [0.037] N=1,004	0.027 [0.043] N=847	0.024 [0.045] N=686
CIAF any stunting underweight or wasting	0.320 P=396	0.090 [0.058] N=1,245	0.065 [0.063] N=1,122	0.098 [0.064] N=983	0.097 [0.069] N=828	**0.146* [0.075] N=672**
Below-normal BMI-for-age BMIZ below -2	0.105 P=134	0.036 [0.040] N=1,267	0.029 [0.044] N=1,143	0.058 [0.046] N=1,003	0.058 [0.049] N=845	0.063 [0.054] N=690

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.133 [0.340] N=1,157	0.198 [0.375] N=1,024	0.320 [0.449] N=874	0.203 [0.552] N=711	0.308 [0.688] N=546
WHO weight-for-age z score	0.270 [0.205] N=1,272	**0.450** [0.219] N=1,135**	0.197 [0.232] N=963	0.236 [0.286] N=780	0.260 [0.384] N=590
WHO weight-for-height z score	0.012 [0.300] N=1,100	0.070 [0.303] N=977	-0.078 [0.342] N=837	-0.187 [0.433] N=680	-0.623 [0.625] N=519
WHO BMI-for-age z score	-0.208 [0.301] N=1,102	-0.205 [0.326] N=978	-0.547 [0.366] N=838	-0.685 [0.465] N=680	-0.872 [0.659] N=525
Stunted HAZ below -2	-0.002 [0.073] N=1,157	-0.010 [0.081] N=1,024	-0.041 [0.090] N=874	-0.030 [0.111] N=711	-0.069 [0.136] N=546
Underweight WAZ below -2	-0.056 [0.036] N=1,272	**-0.091** [0.039] N=1,135**	-0.060 [0.042] N=963	**-0.091* [0.047] N=780**	**-0.119** [0.050] N=590**
Wasted WHZ below -2	0.067 [0.043] N=1,100	0.042 [0.049] N=977	**0.104** [0.052] N=837**	**0.086* [0.052] N=680**	**0.154** [0.065] N=519**
Overweight WHZ above +2	0.012 [0.068] N=1,100	0.017 [0.072] N=977	0.039 [0.081] N=837	0.015 [0.108] N=680	-0.036 [0.141] N=519
Obese WHZ above +3	-0.002 [0.048] N=1,100	-0.005 [0.050] N=977	-0.026 [0.057] N=837	-0.023 [0.074] N=680	-0.102 [0.093] N=519
CIAF any stunting underweight or wasting	0.049 [0.080] N=1,082	-0.009 [0.089] N=959	0.035 [0.099] N=820	-0.006 [0.116] N=665	0.066 [0.144] N=509
Below-normal BMI-for-age BMIZ below -2	**0.074* [0.040] N=1,102**	0.068 [0.048] N=978	**0.134*** [0.050] N=838**	**0.137** [0.053] N=680**	**0.197*** [0.066] N=525**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 24 to 59 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Neither aid control is combined with an income control. Aid specifications use only the six main questions; households with a blank or 99 on any question are excluded. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one earliest-born eligible child per household; all complete-DOB development respondents are included; raw ages below 24 map to 24 and above 59 map to 59. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.