Table 15 Reduced Form Effects on WHO Anthropometric Outcomes With Monthly Child Age Controls Using 12 November 2022 Child Age

Continuous z scores and prespecified adverse clinical classifications exclude WHO-flagged measurements. Child age is calculated on 12 November 2022 and capped at 24 and 59 months. Father RDD estimates are the primary analysis and mother RDD estimates are secondary.

Father RDD primary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.371	0.227 [0.214] N=1,517	0.247 [0.225] N=1,404	0.253 [0.236] N=1,262	0.236 [0.250] N=1,079	0.129 [0.287] N=887
WHO weight-for-age z score	0.130	0.089 [0.129] N=1,681	0.131 [0.137] N=1,552	0.081 [0.136] N=1,397	0.076 [0.162] N=1,191	-0.018 [0.163] N=979
WHO weight-for-height z score	0.369	-0.135 [0.194] N=1,441	-0.149 [0.196] N=1,343	-0.255 [0.216] N=1,207	-0.308 [0.247] N=1,031	-0.188 [0.262] N=846
WHO BMI-for-age z score	0.354	-0.163 [0.213] N=1,442	-0.167 [0.222] N=1,341	-0.304 [0.243] N=1,200	-0.379 [0.285] N=1,025	-0.282 [0.303] N=845
Stunted HAZ below -2	0.220 P=324	-0.033 [0.042] N=1,517	-0.044 [0.044] N=1,404	-0.036 [0.047] N=1,262	-0.052 [0.052] N=1,079	-0.004 [0.059] N=887
Underweight WAZ below -2	0.060 P=100	**-0.037* [0.021] N=1,681**	**-0.042* [0.021] N=1,552**	**-0.044** [0.022] N=1,397**	**-0.041* [0.024] N=1,191**	-0.020 [0.028] N=979
Wasted WHZ below -2	0.106 P=159	-0.017 [0.033] N=1,441	-0.006 [0.033] N=1,343	0.006 [0.037] N=1,207	0.000 [0.042] N=1,031	-0.009 [0.046] N=846
Overweight WHZ above +2	0.196 P=274	-0.013 [0.043] N=1,441	-0.012 [0.045] N=1,343	-0.015 [0.049] N=1,207	-0.023 [0.054] N=1,031	-0.010 [0.060] N=846
Obese WHZ above +3	0.090 P=129	-0.018 [0.031] N=1,441	-0.022 [0.032] N=1,343	-0.022 [0.034] N=1,207	-0.029 [0.038] N=1,031	-0.026 [0.044] N=846
CIAF any stunting underweight or wasting	0.298 P=419	**-0.083* [0.045] N=1,415**	**-0.083* [0.044] N=1,317**	-0.057 [0.047] N=1,183	-0.078 [0.051] N=1,010	-0.021 [0.057] N=827
Below-normal BMI-for-age BMIZ below -2	0.111 P=167	-0.000 [0.036] N=1,442	0.006 [0.036] N=1,341	0.022 [0.040] N=1,200	0.030 [0.045] N=1,025	0.039 [0.049] N=845

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	**8 years**	**7 years**	**6 years**	**5 years**	**4 years**
WHO height-for-age z score	0.173 [0.327] N=1,281	0.185 [0.365] N=1,168	0.217 [0.403] N=1,026	0.219 [0.464] N=843	-0.124 [0.666] N=651
WHO weight-for-age z score	0.116 [0.176] N=1,424	0.190 [0.188] N=1,295	0.118 [0.184] N=1,140	0.149 [0.259] N=934	-0.047 [0.297] N=722
WHO weight-for-height z score	-0.203 [0.274] N=1,216	-0.223 [0.292] N=1,118	-0.466 [0.352] N=982	-0.584 [0.448] N=806	-0.267 [0.545] N=621
WHO BMI-for-age z score	-0.127 [0.286] N=1,219	-0.103 [0.314] N=1,118	-0.364 [0.371] N=977	-0.511 [0.481] N=802	-0.213 [0.583] N=622
Stunted HAZ below -2	-0.014 [0.058] N=1,281	-0.033 [0.066] N=1,168	-0.016 [0.073] N=1,026	-0.054 [0.087] N=843	0.075 [0.113] N=651
Underweight WAZ below -2	**-0.063** [0.031] N=1,424**	**-0.075** [0.033] N=1,295**	**-0.090*** [0.034] N=1,140**	**-0.095** [0.041] N=934**	-0.053 [0.055] N=722
Wasted WHZ below -2	0.009 [0.038] N=1,216	0.030 [0.038] N=1,118	0.065 [0.045] N=982	0.061 [0.059] N=806	0.049 [0.067] N=621
Overweight WHZ above +2	-0.011 [0.056] N=1,216	-0.008 [0.060] N=1,118	-0.016 [0.067] N=982	-0.034 [0.082] N=806	0.030 [0.105] N=621
Obese WHZ above +3	0.000 [0.042] N=1,216	-0.005 [0.046] N=1,118	0.007 [0.053] N=982	0.002 [0.067] N=806	0.045 [0.094] N=621
CIAF any stunting underweight or wasting	-0.062 [0.063] N=1,198	-0.062 [0.065] N=1,100	-0.006 [0.072] N=966	-0.054 [0.086] N=793	0.105 [0.107] N=610
Below-normal BMI-for-age BMIZ below -2	-0.009 [0.041] N=1,219	-0.003 [0.041] N=1,118	0.026 [0.048] N=977	0.036 [0.060] N=802	0.050 [0.072] N=622

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.371	0.229 [0.214] N=1,517	0.250 [0.225] N=1,404	0.255 [0.235] N=1,262	0.238 [0.248] N=1,079	0.130 [0.285] N=887
WHO weight-for-age z score	0.130	0.088 [0.129] N=1,681	0.129 [0.137] N=1,552	0.081 [0.136] N=1,397	0.076 [0.162] N=1,191	-0.018 [0.163] N=979
WHO weight-for-height z score	0.369	-0.137 [0.194] N=1,441	-0.154 [0.197] N=1,343	-0.258 [0.217] N=1,207	-0.306 [0.247] N=1,031	-0.186 [0.262] N=846
WHO BMI-for-age z score	0.354	-0.166 [0.213] N=1,442	-0.171 [0.222] N=1,341	-0.306 [0.242] N=1,200	-0.376 [0.283] N=1,025	-0.278 [0.299] N=845
Stunted HAZ below -2	0.220 P=324	-0.033 [0.042] N=1,517	-0.044 [0.045] N=1,404	-0.036 [0.047] N=1,262	-0.052 [0.052] N=1,079	-0.005 [0.059] N=887
Underweight WAZ below -2	0.060 P=100	**-0.037* [0.021] N=1,681**	**-0.041* [0.021] N=1,552**	**-0.044** [0.022] N=1,397**	**-0.041* [0.024] N=1,191**	-0.020 [0.028] N=979
Wasted WHZ below -2	0.106 P=159	-0.015 [0.033] N=1,441	-0.004 [0.033] N=1,343	0.007 [0.036] N=1,207	-0.001 [0.041] N=1,031	-0.010 [0.044] N=846
Overweight WHZ above +2	0.196 P=274	-0.012 [0.043] N=1,441	-0.012 [0.045] N=1,343	-0.015 [0.049] N=1,207	-0.024 [0.054] N=1,031	-0.011 [0.060] N=846
Obese WHZ above +3	0.090 P=129	-0.018 [0.031] N=1,441	-0.022 [0.032] N=1,343	-0.021 [0.034] N=1,207	-0.029 [0.038] N=1,031	-0.026 [0.043] N=846
CIAF any stunting underweight or wasting	0.298 P=419	**-0.081* [0.044] N=1,415**	**-0.080* [0.044] N=1,317**	-0.055 [0.046] N=1,183	-0.078 [0.050] N=1,010	-0.021 [0.056] N=827
Below-normal BMI-for-age BMIZ below -2	0.111 P=167	0.001 [0.036] N=1,442	0.007 [0.036] N=1,341	0.022 [0.039] N=1,200	0.029 [0.044] N=1,025	0.038 [0.048] N=845

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.179 [0.326] N=1,281	0.194 [0.363] N=1,168	0.225 [0.401] N=1,026	0.233 [0.459] N=843	-0.098 [0.659] N=651
WHO weight-for-age z score	0.114 [0.177] N=1,424	0.187 [0.189] N=1,295	0.117 [0.185] N=1,140	0.149 [0.259] N=934	-0.048 [0.298] N=722
WHO weight-for-height z score	-0.212 [0.275] N=1,216	-0.238 [0.294] N=1,118	-0.481 [0.354] N=982	-0.593 [0.450] N=806	-0.297 [0.555] N=621
WHO BMI-for-age z score	-0.136 [0.287] N=1,219	-0.119 [0.317] N=1,118	-0.376 [0.373] N=977	-0.517 [0.484] N=802	-0.240 [0.592] N=622
Stunted HAZ below -2	-0.014 [0.058] N=1,281	-0.033 [0.066] N=1,168	-0.016 [0.073] N=1,026	-0.055 [0.087] N=843	0.073 [0.113] N=651
Underweight WAZ below -2	**-0.063** [0.031] N=1,424**	**-0.075** [0.033] N=1,295**	**-0.090*** [0.034] N=1,140**	**-0.095** [0.041] N=934**	-0.053 [0.054] N=722
Wasted WHZ below -2	0.012 [0.039] N=1,216	0.036 [0.039] N=1,118	0.070 [0.046] N=982	0.064 [0.060] N=806	0.057 [0.070] N=621
Overweight WHZ above +2	-0.010 [0.056] N=1,216	-0.007 [0.059] N=1,118	-0.015 [0.067] N=982	-0.033 [0.081] N=806	0.032 [0.104] N=621
Obese WHZ above +3	0.001 [0.042] N=1,216	-0.004 [0.045] N=1,118	0.007 [0.052] N=982	0.002 [0.067] N=806	0.046 [0.094] N=621
CIAF any stunting underweight or wasting	-0.059 [0.063] N=1,198	-0.056 [0.064] N=1,100	-0.001 [0.071] N=966	-0.051 [0.085] N=793	0.113 [0.107] N=610
Below-normal BMI-for-age BMIZ below -2	-0.007 [0.041] N=1,219	0.000 [0.042] N=1,118	0.029 [0.048] N=977	0.037 [0.061] N=802	0.055 [0.075] N=622

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.371	0.176 [0.210] N=1,517	0.177 [0.221] N=1,404	0.201 [0.232] N=1,262	0.164 [0.244] N=1,079	0.045 [0.280] N=887
WHO weight-for-age z score	0.130	0.032 [0.127] N=1,681	0.075 [0.136] N=1,552	0.028 [0.138] N=1,397	0.012 [0.162] N=1,191	-0.089 [0.167] N=979
WHO weight-for-height z score	0.369	-0.170 [0.192] N=1,441	-0.179 [0.196] N=1,343	-0.287 [0.215] N=1,207	-0.333 [0.244] N=1,031	-0.222 [0.259] N=846
WHO BMI-for-age z score	0.354	-0.182 [0.211] N=1,442	-0.179 [0.220] N=1,341	-0.319 [0.239] N=1,200	-0.387 [0.278] N=1,025	-0.289 [0.296] N=845
Stunted HAZ below -2	0.220 P=324	-0.021 [0.041] N=1,517	-0.027 [0.044] N=1,404	-0.024 [0.046] N=1,262	-0.038 [0.051] N=1,079	0.014 [0.058] N=887
Underweight WAZ below -2	0.060 P=100	-0.028 [0.020] N=1,681	-0.031 [0.021] N=1,552	-0.033 [0.022] N=1,397	-0.027 [0.024] N=1,191	-0.004 [0.027] N=979
Wasted WHZ below -2	0.106 P=159	-0.008 [0.033] N=1,441	0.005 [0.032] N=1,343	0.015 [0.035] N=1,207	0.007 [0.040] N=1,031	-0.001 [0.044] N=846
Overweight WHZ above +2	0.196 P=274	-0.013 [0.044] N=1,441	-0.011 [0.046] N=1,343	-0.015 [0.049] N=1,207	-0.025 [0.055] N=1,031	-0.014 [0.061] N=846
Obese WHZ above +3	0.090 P=129	-0.017 [0.030] N=1,441	-0.020 [0.031] N=1,343	-0.021 [0.034] N=1,207	-0.031 [0.037] N=1,031	-0.030 [0.044] N=846
CIAF any stunting underweight or wasting	0.298 P=419	-0.063 [0.043] N=1,415	-0.057 [0.042] N=1,317	-0.037 [0.043] N=1,183	-0.059 [0.048] N=1,010	0.005 [0.052] N=827
Below-normal BMI-for-age BMIZ below -2	0.111 P=167	0.006 [0.035] N=1,442	0.014 [0.036] N=1,341	0.029 [0.039] N=1,200	0.035 [0.043] N=1,025	0.046 [0.047] N=845

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.138 [0.321] N=1,281	0.122 [0.355] N=1,168	0.180 [0.397] N=1,026	0.148 [0.451] N=843	-0.219 [0.639] N=651
WHO weight-for-age z score	0.059 [0.172] N=1,424	0.129 [0.185] N=1,295	0.061 [0.183] N=1,140	0.063 [0.254] N=934	-0.159 [0.302] N=722
WHO weight-for-height z score	-0.235 [0.275] N=1,216	-0.249 [0.297] N=1,118	-0.495 [0.356] N=982	-0.614 [0.456] N=806	-0.340 [0.561] N=621
WHO BMI-for-age z score	-0.144 [0.287] N=1,219	-0.116 [0.317] N=1,118	-0.380 [0.373] N=977	-0.542 [0.491] N=802	-0.251 [0.608] N=622
Stunted HAZ below -2	-0.006 [0.057] N=1,281	-0.019 [0.064] N=1,168	-0.008 [0.071] N=1,026	-0.045 [0.086] N=843	0.090 [0.110] N=651
Underweight WAZ below -2	**-0.054* [0.031] N=1,424**	**-0.062* [0.033] N=1,295**	**-0.078** [0.034] N=1,140**	**-0.078* [0.040] N=934**	-0.030 [0.053] N=722
Wasted WHZ below -2	0.019 [0.039] N=1,216	0.044 [0.039] N=1,118	**0.077* [0.045] N=982**	0.073 [0.059] N=806	0.068 [0.071] N=621
Overweight WHZ above +2	-0.007 [0.057] N=1,216	-0.001 [0.061] N=1,118	-0.012 [0.068] N=982	-0.031 [0.083] N=806	0.034 [0.105] N=621
Obese WHZ above +3	0.003 [0.041] N=1,216	0.000 [0.045] N=1,118	0.009 [0.052] N=982	0.001 [0.066] N=806	0.035 [0.092] N=621
CIAF any stunting underweight or wasting	-0.044 [0.062] N=1,198	-0.034 [0.062] N=1,100	0.013 [0.068] N=966	-0.032 [0.083] N=793	0.145 [0.104] N=610
Below-normal BMI-for-age BMIZ below -2	-0.003 [0.041] N=1,219	0.006 [0.042] N=1,118	0.034 [0.047] N=977	0.047 [0.060] N=802	0.063 [0.075] N=622

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.371	0.121 [0.211] N=1,517	0.119 [0.222] N=1,404	0.145 [0.236] N=1,262	0.102 [0.248] N=1,079	-0.049 [0.283] N=887
WHO weight-for-age z score	0.130	0.036 [0.127] N=1,681	0.071 [0.135] N=1,552	0.026 [0.138] N=1,397	0.002 [0.163] N=1,191	-0.113 [0.163] N=979
WHO weight-for-height z score	0.369	-0.097 [0.194] N=1,441	-0.106 [0.197] N=1,343	-0.221 [0.217] N=1,207	-0.274 [0.250] N=1,031	-0.141 [0.265] N=846
WHO BMI-for-age z score	0.354	-0.121 [0.215] N=1,442	-0.118 [0.224] N=1,341	-0.262 [0.245] N=1,200	-0.335 [0.291] N=1,025	-0.225 [0.311] N=845
Stunted HAZ below -2	0.220 P=324	-0.012 [0.041] N=1,517	-0.019 [0.044] N=1,404	-0.014 [0.047] N=1,262	-0.024 [0.051] N=1,079	0.029 [0.057] N=887
Underweight WAZ below -2	0.060 P=100	-0.028 [0.021] N=1,681	-0.030 [0.021] N=1,552	-0.031 [0.023] N=1,397	-0.024 [0.025] N=1,191	-0.001 [0.028] N=979
Wasted WHZ below -2	0.106 P=159	-0.013 [0.033] N=1,441	-0.001 [0.033] N=1,343	0.012 [0.036] N=1,207	0.004 [0.042] N=1,031	-0.007 [0.045] N=846
Overweight WHZ above +2	0.196 P=274	-0.003 [0.044] N=1,441	-0.002 [0.045] N=1,343	-0.007 [0.049] N=1,207	-0.017 [0.055] N=1,031	-0.004 [0.061] N=846
Obese WHZ above +3	0.090 P=129	-0.010 [0.030] N=1,441	-0.013 [0.032] N=1,343	-0.013 [0.034] N=1,207	-0.022 [0.038] N=1,031	-0.017 [0.044] N=846
CIAF any stunting underweight or wasting	0.298 P=419	-0.058 [0.043] N=1,415	-0.054 [0.043] N=1,317	-0.031 [0.045] N=1,183	-0.047 [0.048] N=1,010	0.014 [0.053] N=827
Below-normal BMI-for-age BMIZ below -2	0.111 P=167	0.002 [0.036] N=1,442	0.009 [0.036] N=1,341	0.026 [0.040] N=1,200	0.033 [0.044] N=1,025	0.040 [0.049] N=845

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.119 [0.316] N=1,281	0.100 [0.349] N=1,168	0.167 [0.392] N=1,026	0.150 [0.439] N=843	-0.284 [0.614] N=651
WHO weight-for-age z score	0.064 [0.172] N=1,424	0.131 [0.185] N=1,295	0.062 [0.186] N=1,140	0.072 [0.259] N=934	-0.148 [0.305] N=722
WHO weight-for-height z score	-0.210 [0.276] N=1,216	-0.209 [0.295] N=1,118	-0.465 [0.357] N=982	-0.594 [0.457] N=806	-0.255 [0.556] N=621
WHO BMI-for-age z score	-0.132 [0.287] N=1,219	-0.089 [0.315] N=1,118	-0.361 [0.375] N=977	-0.525 [0.489] N=802	-0.192 [0.600] N=622
Stunted HAZ below -2	-0.001 [0.057] N=1,281	-0.014 [0.064] N=1,168	-0.006 [0.072] N=1,026	-0.043 [0.086] N=843	0.107 [0.111] N=651
Underweight WAZ below -2	**-0.051* [0.031] N=1,424**	**-0.061* [0.033] N=1,295**	**-0.075** [0.034] N=1,140**	**-0.073* [0.042] N=934**	-0.024 [0.055] N=722
Wasted WHZ below -2	0.020 [0.039] N=1,216	0.041 [0.039] N=1,118	0.075 [0.046] N=982	0.072 [0.059] N=806	0.061 [0.071] N=621
Overweight WHZ above +2	-0.003 [0.057] N=1,216	0.003 [0.060] N=1,118	-0.009 [0.068] N=982	-0.027 [0.084] N=806	0.045 [0.104] N=621
Obese WHZ above +3	0.006 [0.042] N=1,216	0.005 [0.046] N=1,118	0.014 [0.053] N=982	0.007 [0.067] N=806	0.052 [0.092] N=621
CIAF any stunting underweight or wasting	-0.036 [0.063] N=1,198	-0.031 [0.064] N=1,100	0.015 [0.070] N=966	-0.028 [0.083] N=793	0.159 [0.106] N=610
Below-normal BMI-for-age BMIZ below -2	-0.002 [0.041] N=1,219	0.002 [0.042] N=1,118	0.031 [0.048] N=977	0.043 [0.061] N=802	0.055 [0.075] N=622

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.380	0.180 [0.219] N=1,430	0.173 [0.229] N=1,329	0.177 [0.236] N=1,192	0.204 [0.240] N=1,022	0.103 [0.279] N=836
WHO weight-for-age z score	0.123	0.093 [0.134] N=1,583	0.132 [0.139] N=1,467	0.082 [0.142] N=1,319	0.069 [0.168] N=1,128	-0.002 [0.178] N=924
WHO weight-for-height z score	0.371	-0.047 [0.206] N=1,360	-0.068 [0.207] N=1,274	-0.167 [0.229] N=1,144	-0.227 [0.261] N=980	-0.112 [0.287] N=801
WHO BMI-for-age z score	0.363	-0.074 [0.220] N=1,358	-0.082 [0.227] N=1,269	-0.195 [0.249] N=1,135	-0.266 [0.290] N=972	-0.167 [0.314] N=798
Stunted HAZ below -2	0.221 P=306	-0.017 [0.044] N=1,430	-0.026 [0.047] N=1,329	-0.019 [0.049] N=1,192	-0.046 [0.052] N=1,022	0.007 [0.059] N=836
Underweight WAZ below -2	0.061 P=96	-0.034 [0.022] N=1,583	**-0.038* [0.022] N=1,467**	**-0.041* [0.024] N=1,319**	-0.038 [0.026] N=1,128	-0.021 [0.032] N=924
Wasted WHZ below -2	0.107 P=152	-0.014 [0.035] N=1,360	-0.006 [0.034] N=1,274	0.007 [0.037] N=1,144	-0.002 [0.043] N=980	-0.009 [0.048] N=801
Overweight WHZ above +2	0.197 P=258	-0.002 [0.046] N=1,360	-0.005 [0.048] N=1,274	-0.008 [0.052] N=1,144	-0.020 [0.058] N=980	-0.008 [0.067] N=801
Obese WHZ above +3	0.094 P=126	-0.015 [0.032] N=1,360	-0.020 [0.034] N=1,274	-0.020 [0.036] N=1,144	-0.031 [0.039] N=980	-0.029 [0.046] N=801
CIAF any stunting underweight or wasting	0.301 P=399	-0.062 [0.048] N=1,336	-0.065 [0.048] N=1,250	-0.043 [0.051] N=1,121	-0.076 [0.056] N=960	-0.012 [0.064] N=783
Below-normal BMI-for-age BMIZ below -2	0.111 P=157	-0.001 [0.036] N=1,358	0.003 [0.035] N=1,269	0.016 [0.039] N=1,135	0.021 [0.044] N=972	0.033 [0.048] N=798

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.135 [0.343] N=1,209	0.105 [0.380] N=1,108	0.138 [0.418] N=971	0.272 [0.465] N=801	-0.041 [0.698] N=615
WHO weight-for-age z score	0.110 [0.179] N=1,342	0.168 [0.182] N=1,226	0.077 [0.188] N=1,078	0.094 [0.261] N=887	-0.090 [0.321] N=683
WHO weight-for-height z score	-0.135 [0.282] N=1,147	-0.176 [0.294] N=1,061	-0.408 [0.350] N=931	-0.522 [0.448] N=767	-0.189 [0.558] N=588
WHO BMI-for-age z score	-0.089 [0.293] N=1,149	-0.089 [0.316] N=1,060	-0.302 [0.374] N=926	-0.440 [0.480] N=763	-0.129 [0.598] N=589
Stunted HAZ below -2	0.002 [0.061] N=1,209	-0.015 [0.069] N=1,108	-0.001 [0.075] N=971	-0.061 [0.087] N=801	0.077 [0.117] N=615
Underweight WAZ below -2	**-0.057* [0.031] N=1,342**	**-0.065** [0.031] N=1,226**	**-0.076** [0.034] N=1,078**	**-0.076* [0.040] N=887**	-0.034 [0.059] N=683
Wasted WHZ below -2	0.013 [0.040] N=1,147	0.031 [0.038] N=1,061	0.069 [0.044] N=931	0.055 [0.059] N=767	0.045 [0.069] N=588
Overweight WHZ above +2	-0.002 [0.058] N=1,147	-0.008 [0.060] N=1,061	-0.009 [0.067] N=931	-0.036 [0.083] N=767	0.039 [0.111] N=588
Obese WHZ above +3	0.007 [0.045] N=1,147	0.001 [0.049] N=1,061	0.011 [0.055] N=931	0.003 [0.070] N=767	0.056 [0.099] N=588
CIAF any stunting underweight or wasting	-0.036 [0.066] N=1,131	-0.040 [0.066] N=1,045	0.015 [0.074] N=916	-0.058 [0.084] N=755	0.114 [0.112] N=578
Below-normal BMI-for-age BMIZ below -2	-0.001 [0.042] N=1,149	0.003 [0.041] N=1,060	0.029 [0.047] N=926	0.033 [0.060] N=763	0.051 [0.073] N=589

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	**Mean**	**8 years**	**7 years**	**6 years**	**5 years**	**4 years**
WHO height-for-age z score	-0.380	0.181 [0.218] N=1,430	0.175 [0.228] N=1,329	0.150 [0.237] N=1,192	0.163 [0.245] N=1,022	0.037 [0.284] N=836
WHO weight-for-age z score	0.123	0.093 [0.136] N=1,583	0.131 [0.141] N=1,467	0.075 [0.142] N=1,319	0.068 [0.166] N=1,128	-0.012 [0.175] N=924
WHO weight-for-height z score	0.371	-0.046 [0.206] N=1,360	-0.065 [0.207] N=1,274	-0.165 [0.227] N=1,144	-0.212 [0.260] N=980	-0.087 [0.283] N=801
WHO BMI-for-age z score	0.363	-0.075 [0.220] N=1,358	-0.082 [0.227] N=1,269	-0.194 [0.249] N=1,135	-0.252 [0.292] N=972	-0.144 [0.314] N=798
Stunted HAZ below -2	0.221 P=306	-0.018 [0.044] N=1,430	-0.028 [0.047] N=1,329	-0.018 [0.049] N=1,192	-0.042 [0.053] N=1,022	0.011 [0.061] N=836
Underweight WAZ below -2	0.061 P=96	-0.036 [0.022] N=1,583	**-0.040* [0.023] N=1,467**	**-0.042* [0.024] N=1,319**	-0.039 [0.026] N=1,128	-0.021 [0.032] N=924
Wasted WHZ below -2	0.107 P=152	-0.015 [0.035] N=1,360	-0.007 [0.035] N=1,274	0.005 [0.038] N=1,144	-0.006 [0.044] N=980	-0.018 [0.049] N=801
Overweight WHZ above +2	0.197 P=258	-0.002 [0.046] N=1,360	-0.004 [0.047] N=1,274	-0.006 [0.051] N=1,144	-0.017 [0.057] N=980	-0.002 [0.066] N=801
Obese WHZ above +3	0.094 P=126	-0.015 [0.032] N=1,360	-0.020 [0.033] N=1,274	-0.019 [0.035] N=1,144	-0.029 [0.039] N=980	-0.024 [0.045] N=801
CIAF any stunting underweight or wasting	0.301 P=399	-0.063 [0.048] N=1,336	-0.066 [0.048] N=1,250	-0.042 [0.051] N=1,121	-0.075 [0.055] N=960	-0.017 [0.063] N=783
Below-normal BMI-for-age BMIZ below -2	0.111 P=157	-0.001 [0.036] N=1,358	0.003 [0.036] N=1,269	0.015 [0.039] N=1,135	0.018 [0.045] N=972	0.026 [0.049] N=798

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.200 [0.345] N=1,209	0.182 [0.380] N=1,108	0.165 [0.420] N=971	0.288 [0.488] N=801	-0.053 [0.724] N=615
WHO weight-for-age z score	0.130 [0.180] N=1,342	0.193 [0.185] N=1,226	0.109 [0.185] N=1,078	0.143 [0.252] N=887	-0.029 [0.306] N=683
WHO weight-for-height z score	-0.123 [0.281] N=1,147	-0.154 [0.294] N=1,061	-0.390 [0.350] N=931	-0.498 [0.454] N=767	-0.168 [0.565] N=588
WHO BMI-for-age z score	-0.081 [0.292] N=1,149	-0.074 [0.316] N=1,060	-0.295 [0.371] N=926	-0.418 [0.486] N=763	-0.106 [0.610] N=589
Stunted HAZ below -2	-0.006 [0.061] N=1,209	-0.024 [0.069] N=1,108	-0.005 [0.075] N=971	-0.070 [0.089] N=801	0.069 [0.117] N=615
Underweight WAZ below -2	**-0.061* [0.031] N=1,342**	**-0.071** [0.033] N=1,226**	**-0.086** [0.034] N=1,078**	**-0.088** [0.039] N=887**	-0.052 [0.056] N=683
Wasted WHZ below -2	0.015 [0.039] N=1,147	0.032 [0.038] N=1,061	0.069 [0.044] N=931	0.054 [0.059] N=767	0.038 [0.069] N=588
Overweight WHZ above +2	0.001 [0.058] N=1,147	-0.001 [0.060] N=1,061	-0.005 [0.067] N=931	-0.030 [0.083] N=767	0.047 [0.111] N=588
Obese WHZ above +3	0.006 [0.044] N=1,147	-0.000 [0.049] N=1,061	0.013 [0.055] N=931	0.004 [0.070] N=767	0.060 [0.098] N=588
CIAF any stunting underweight or wasting	-0.038 [0.065] N=1,131	-0.043 [0.067] N=1,045	0.011 [0.074] N=916	-0.069 [0.085] N=755	0.094 [0.109] N=578
Below-normal BMI-for-age BMIZ below -2	-0.002 [0.041] N=1,149	0.002 [0.040] N=1,060	0.028 [0.047] N=926	0.028 [0.060] N=763	0.040 [0.073] N=589

Mother RDD secondary analysis

Specification 1 Monthly child-age fixed effects No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.326	-0.103 [0.241] N=1,411	-0.146 [0.256] N=1,275	-0.148 [0.280] N=1,116	-0.190 [0.312] N=943	-0.242 [0.319] N=765
WHO weight-for-age z score	0.133	0.042 [0.137] N=1,552	0.118 [0.146] N=1,412	-0.036 [0.145] N=1,230	-0.067 [0.157] N=1,036	-0.021 [0.170] N=832
WHO weight-for-height z score	0.387	0.079 [0.196] N=1,343	0.122 [0.202] N=1,215	0.068 [0.215] N=1,067	0.034 [0.241] N=900	-0.015 [0.269] N=729
WHO BMI-for-age z score	0.339	-0.035 [0.224] N=1,339	-0.005 [0.237] N=1,213	-0.133 [0.253] N=1,065	-0.161 [0.271] N=897	-0.130 [0.289] N=732
Stunted HAZ below -2	0.219 P=308	0.014 [0.045] N=1,411	0.025 [0.047] N=1,275	0.027 [0.049] N=1,116	0.033 [0.053] N=943	0.047 [0.058] N=765
Underweight WAZ below -2	0.056 P=89	-0.011 [0.024] N=1,552	-0.026 [0.026] N=1,412	-0.003 [0.027] N=1,230	-0.014 [0.029] N=1,036	-0.034 [0.031] N=832
Wasted WHZ below -2	0.105 P=142	0.051 [0.032] N=1,343	0.040 [0.034] N=1,215	**0.065* [0.037] N=1,067**	0.045 [0.037] N=900	0.053 [0.041] N=729
Overweight WHZ above +2	0.194 P=263	0.047 [0.041] N=1,343	0.052 [0.042] N=1,215	0.058 [0.046] N=1,067	0.057 [0.051] N=900	0.064 [0.055] N=729
Obese WHZ above +3	0.097 P=134	0.023 [0.031] N=1,343	0.034 [0.031] N=1,215	0.021 [0.036] N=1,067	0.031 [0.041] N=900	0.015 [0.044] N=729
CIAF any stunting underweight or wasting	0.297 P=392	0.075 [0.053] N=1,319	0.057 [0.056] N=1,191	0.091 [0.060] N=1,044	0.076 [0.064] N=880	**0.114* [0.066] N=713**
Below-normal BMI-for-age BMIZ below -2	0.108 P=148	0.050 [0.037] N=1,339	0.040 [0.041] N=1,213	0.064 [0.044] N=1,065	0.055 [0.045] N=897	0.061 [0.047] N=732

Specification 1 Monthly child-age fixed effects 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.315 [0.294] N=1,221	0.299 [0.328] N=1,085	0.413 [0.380] N=926	0.471 [0.492] N=753	0.719 [0.571] N=575
WHO weight-for-age z score	0.167 [0.213] N=1,347	0.312 [0.229] N=1,207	0.037 [0.248] N=1,025	0.017 [0.310] N=831	0.180 [0.370] N=627
WHO weight-for-height z score	-0.122 [0.287] N=1,168	-0.064 [0.297] N=1,040	-0.196 [0.329] N=892	-0.325 [0.422] N=725	-0.769 [0.573] N=554
WHO BMI-for-age z score	-0.344 [0.289] N=1,165	-0.328 [0.318] N=1,039	**-0.631* [0.355] N=891**	-0.758 [0.459] N=723	**-1.026* [0.584] N=558**
Stunted HAZ below -2	-0.027 [0.067] N=1,221	-0.014 [0.075] N=1,085	-0.025 [0.082] N=926	-0.035 [0.100] N=753	-0.047 [0.130] N=575
Underweight WAZ below -2	-0.007 [0.033] N=1,347	-0.028 [0.037] N=1,207	0.018 [0.042] N=1,025	-0.008 [0.050] N=831	-0.064 [0.049] N=627
Wasted WHZ below -2	**0.093** [0.040] N=1,168**	**0.076* [0.045] N=1,040**	**0.135** [0.053] N=892**	**0.105** [0.052] N=725**	**0.170*** [0.058] N=554**
Overweight WHZ above +2	0.024 [0.063] N=1,168	0.026 [0.066] N=1,040	0.034 [0.076] N=892	0.009 [0.095] N=725	-0.031 [0.123] N=554
Obese WHZ above +3	0.007 [0.046] N=1,168	0.020 [0.048] N=1,040	-0.006 [0.055] N=892	-0.002 [0.071] N=725	-0.100 [0.089] N=554
CIAF any stunting underweight or wasting	0.053 [0.074] N=1,148	0.017 [0.082] N=1,020	0.073 [0.096] N=873	0.015 [0.112] N=709	0.074 [0.128] N=542
Below-normal BMI-for-age BMIZ below -2	**0.103*** [0.037] N=1,165**	**0.093** [0.045] N=1,039**	**0.150*** [0.053] N=891**	**0.144*** [0.054] N=723**	**0.203*** [0.056] N=558**

Specification 2 Monthly fixed effects and female No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.326	-0.109 [0.241] N=1,411	-0.155 [0.256] N=1,275	-0.153 [0.280] N=1,116	-0.193 [0.312] N=943	-0.221 [0.321] N=765
WHO weight-for-age z score	0.133	0.043 [0.137] N=1,552	0.118 [0.146] N=1,412	-0.036 [0.145] N=1,230	-0.067 [0.157] N=1,036	-0.024 [0.171] N=832
WHO weight-for-height z score	0.387	0.082 [0.197] N=1,343	0.126 [0.202] N=1,215	0.071 [0.215] N=1,067	0.032 [0.242] N=900	-0.040 [0.271] N=729
WHO BMI-for-age z score	0.339	-0.033 [0.224] N=1,339	-0.003 [0.238] N=1,213	-0.133 [0.253] N=1,065	-0.164 [0.272] N=897	-0.158 [0.292] N=732
Stunted HAZ below -2	0.219 P=308	0.015 [0.045] N=1,411	0.026 [0.047] N=1,275	0.027 [0.049] N=1,116	0.033 [0.053] N=943	0.046 [0.059] N=765
Underweight WAZ below -2	0.056 P=89	-0.012 [0.024] N=1,552	-0.026 [0.026] N=1,412	-0.003 [0.027] N=1,230	-0.013 [0.029] N=1,036	-0.032 [0.031] N=832
Wasted WHZ below -2	0.105 P=142	0.050 [0.032] N=1,343	0.038 [0.035] N=1,215	**0.064* [0.037] N=1,067**	0.046 [0.037] N=900	0.061 [0.040] N=729
Overweight WHZ above +2	0.194 P=263	0.047 [0.041] N=1,343	0.052 [0.042] N=1,215	0.058 [0.046] N=1,067	0.058 [0.051] N=900	0.065 [0.055] N=729
Obese WHZ above +3	0.097 P=134	0.023 [0.031] N=1,343	0.033 [0.031] N=1,215	0.020 [0.036] N=1,067	0.031 [0.041] N=900	0.018 [0.045] N=729
CIAF any stunting underweight or wasting	0.297 P=392	0.074 [0.053] N=1,319	0.056 [0.057] N=1,191	0.089 [0.061] N=1,044	0.077 [0.065] N=880	**0.120* [0.067] N=713**
Below-normal BMI-for-age BMIZ below -2	0.108 P=148	0.049 [0.037] N=1,339	0.040 [0.041] N=1,213	0.063 [0.044] N=1,065	0.056 [0.045] N=897	0.067 [0.047] N=732

Specification 2 Monthly fixed effects and female 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.298 [0.293] N=1,221	0.276 [0.328] N=1,085	0.402 [0.382] N=926	0.434 [0.488] N=753	0.695 [0.571] N=575
WHO weight-for-age z score	0.171 [0.212] N=1,347	0.316 [0.228] N=1,207	0.041 [0.246] N=1,025	0.021 [0.307] N=831	0.182 [0.374] N=627
WHO weight-for-height z score	-0.101 [0.288] N=1,168	-0.038 [0.297] N=1,040	-0.175 [0.327] N=892	-0.293 [0.422] N=725	-0.741 [0.588] N=554
WHO BMI-for-age z score	-0.323 [0.291] N=1,165	-0.302 [0.320] N=1,039	**-0.614* [0.355] N=891**	-0.729 [0.461] N=723	**-1.009* [0.601] N=558**
Stunted HAZ below -2	-0.025 [0.067] N=1,221	-0.012 [0.075] N=1,085	-0.028 [0.083] N=926	-0.034 [0.100] N=753	-0.047 [0.130] N=575
Underweight WAZ below -2	-0.009 [0.033] N=1,347	-0.030 [0.037] N=1,207	0.016 [0.042] N=1,025	-0.010 [0.049] N=831	-0.065 [0.050] N=627
Wasted WHZ below -2	**0.086** [0.040] N=1,168**	0.064 [0.046] N=1,040	**0.122** [0.054] N=892**	**0.092* [0.054] N=725**	**0.162*** [0.061] N=554**
Overweight WHZ above +2	0.023 [0.062] N=1,168	0.025 [0.066] N=1,040	0.030 [0.075] N=892	0.007 [0.095] N=725	-0.031 [0.123] N=554
Obese WHZ above +3	0.003 [0.046] N=1,168	0.014 [0.048] N=1,040	-0.015 [0.055] N=892	-0.009 [0.070] N=725	-0.103 [0.088] N=554
CIAF any stunting underweight or wasting	0.047 [0.075] N=1,148	0.006 [0.084] N=1,020	0.056 [0.099] N=873	0.001 [0.116] N=709	0.066 [0.134] N=542
Below-normal BMI-for-age BMIZ below -2	**0.098*** [0.037] N=1,165**	**0.085* [0.046] N=1,039**	**0.141*** [0.053] N=891**	**0.135** [0.054] N=723**	**0.200*** [0.058] N=558**

Specification 3 Monthly fixed effects female and household income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.326	-0.128 [0.239] N=1,411	-0.156 [0.254] N=1,275	-0.152 [0.279] N=1,116	-0.182 [0.309] N=943	-0.231 [0.309] N=765
WHO weight-for-age z score	0.133	0.034 [0.138] N=1,552	0.117 [0.147] N=1,412	-0.036 [0.147] N=1,230	-0.074 [0.158] N=1,036	-0.030 [0.169] N=832
WHO weight-for-height z score	0.387	0.081 [0.201] N=1,343	0.124 [0.207] N=1,215	0.081 [0.222] N=1,067	0.038 [0.248] N=900	-0.019 [0.277] N=729
WHO BMI-for-age z score	0.339	-0.030 [0.225] N=1,339	0.000 [0.239] N=1,213	-0.117 [0.257] N=1,065	-0.158 [0.277] N=897	-0.143 [0.297] N=732
Stunted HAZ below -2	0.219 P=308	0.020 [0.044] N=1,411	0.027 [0.047] N=1,275	0.027 [0.049] N=1,116	0.031 [0.052] N=943	0.046 [0.056] N=765
Underweight WAZ below -2	0.056 P=89	-0.010 [0.024] N=1,552	-0.025 [0.026] N=1,412	-0.003 [0.027] N=1,230	-0.012 [0.029] N=1,036	-0.030 [0.031] N=832
Wasted WHZ below -2	0.105 P=142	0.049 [0.031] N=1,343	0.037 [0.034] N=1,215	0.061 [0.037] N=1,067	0.042 [0.036] N=900	0.054 [0.039] N=729
Overweight WHZ above +2	0.194 P=263	0.047 [0.041] N=1,343	0.051 [0.043] N=1,215	0.058 [0.046] N=1,067	0.055 [0.051] N=900	0.063 [0.055] N=729
Obese WHZ above +3	0.097 P=134	0.024 [0.031] N=1,343	0.033 [0.031] N=1,215	0.021 [0.036] N=1,067	0.032 [0.041] N=900	0.020 [0.045] N=729
CIAF any stunting underweight or wasting	0.297 P=392	0.078 [0.053] N=1,319	0.054 [0.057] N=1,191	0.087 [0.061] N=1,044	0.070 [0.066] N=880	0.111 [0.068] N=713
Below-normal BMI-for-age BMIZ below -2	0.108 P=148	0.049 [0.037] N=1,339	0.039 [0.041] N=1,213	0.061 [0.044] N=1,065	0.054 [0.045] N=897	0.064 [0.046] N=732

Specification 3 Monthly fixed effects female and household income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.334 [0.294] N=1,221	0.339 [0.329] N=1,085	0.506 [0.383] N=926	0.584 [0.494] N=753	**0.937* [0.562] N=575**
WHO weight-for-age z score	0.170 [0.210] N=1,347	0.333 [0.227] N=1,207	0.062 [0.249] N=1,025	0.027 [0.314] N=831	0.172 [0.384] N=627
WHO weight-for-height z score	-0.128 [0.293] N=1,168	-0.059 [0.305] N=1,040	-0.204 [0.339] N=892	-0.369 [0.443] N=725	-0.869 [0.603] N=554
WHO BMI-for-age z score	-0.335 [0.293] N=1,165	-0.312 [0.325] N=1,039	**-0.643* [0.365] N=891**	**-0.817* [0.481] N=723**	**-1.152* [0.612] N=558**
Stunted HAZ below -2	-0.032 [0.068] N=1,221	-0.024 [0.076] N=1,085	-0.045 [0.084] N=926	-0.063 [0.102] N=753	-0.092 [0.130] N=575
Underweight WAZ below -2	-0.013 [0.032] N=1,347	-0.038 [0.037] N=1,207	0.004 [0.041] N=1,025	-0.021 [0.049] N=831	-0.072 [0.048] N=627
Wasted WHZ below -2	**0.084** [0.040] N=1,168**	0.060 [0.047] N=1,040	**0.119** [0.055] N=892**	0.089 [0.056] N=725	**0.157** [0.061] N=554**
Overweight WHZ above +2	0.021 [0.064] N=1,168	0.023 [0.068] N=1,040	0.025 [0.079] N=892	-0.009 [0.100] N=725	-0.055 [0.130] N=554
Obese WHZ above +3	-0.001 [0.047] N=1,168	0.009 [0.048] N=1,040	-0.019 [0.056] N=892	-0.016 [0.072] N=725	-0.118 [0.088] N=554
CIAF any stunting underweight or wasting	0.043 [0.076] N=1,148	-0.002 [0.086] N=1,020	0.046 [0.103] N=873	-0.018 [0.120] N=709	0.038 [0.138] N=542
Below-normal BMI-for-age BMIZ below -2	**0.094** [0.037] N=1,165**	**0.079* [0.046] N=1,039**	**0.138** [0.053] N=891**	**0.132** [0.055] N=723**	**0.199*** [0.056] N=558**

Specification 4 Monthly fixed effects female and per-person income groups No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.326	-0.108 [0.235] N=1,411	-0.135 [0.251] N=1,275	-0.126 [0.278] N=1,116	-0.156 [0.308] N=943	-0.184 [0.310] N=765
WHO weight-for-age z score	0.133	0.046 [0.137] N=1,552	0.129 [0.146] N=1,412	-0.021 [0.146] N=1,230	-0.050 [0.158] N=1,036	-0.017 [0.170] N=832
WHO weight-for-height z score	0.387	0.081 [0.196] N=1,343	0.127 [0.200] N=1,215	0.077 [0.212] N=1,067	0.034 [0.238] N=900	-0.049 [0.271] N=729
WHO BMI-for-age z score	0.339	-0.025 [0.220] N=1,339	0.007 [0.232] N=1,213	-0.123 [0.248] N=1,065	-0.157 [0.265] N=897	-0.158 [0.284] N=732
Stunted HAZ below -2	0.219 P=308	0.017 [0.044] N=1,411	0.023 [0.047] N=1,275	0.024 [0.049] N=1,116	0.027 [0.053] N=943	0.039 [0.057] N=765
Underweight WAZ below -2	0.056 P=89	-0.011 [0.024] N=1,552	-0.027 [0.026] N=1,412	-0.005 [0.027] N=1,230	-0.016 [0.030] N=1,036	-0.035 [0.032] N=832
Wasted WHZ below -2	0.105 P=142	0.050 [0.031] N=1,343	0.037 [0.034] N=1,215	**0.063* [0.037] N=1,067**	0.047 [0.036] N=900	0.059 [0.039] N=729
Overweight WHZ above +2	0.194 P=263	0.049 [0.041] N=1,343	0.054 [0.042] N=1,215	0.061 [0.046] N=1,067	0.060 [0.051] N=900	0.062 [0.055] N=729
Obese WHZ above +3	0.097 P=134	0.025 [0.031] N=1,343	0.035 [0.031] N=1,215	0.022 [0.035] N=1,067	0.034 [0.040] N=900	0.019 [0.045] N=729
CIAF any stunting underweight or wasting	0.297 P=392	0.078 [0.053] N=1,319	0.055 [0.058] N=1,191	0.088 [0.062] N=1,044	0.076 [0.066] N=880	**0.114* [0.068] N=713**
Below-normal BMI-for-age BMIZ below -2	0.108 P=148	0.047 [0.036] N=1,339	0.036 [0.040] N=1,213	0.061 [0.043] N=1,065	0.054 [0.044] N=897	0.060 [0.045] N=732

Specification 4 Monthly fixed effects female and per-person income groups 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.311 [0.284] N=1,221	0.322 [0.319] N=1,085	0.464 [0.375] N=926	0.541 [0.479] N=753	0.836 [0.537] N=575
WHO weight-for-age z score	0.175 [0.203] N=1,347	0.333 [0.221] N=1,207	0.070 [0.237] N=1,025	0.062 [0.298] N=831	0.185 [0.351] N=627
WHO weight-for-height z score	-0.114 [0.286] N=1,168	-0.051 [0.290] N=1,040	-0.189 [0.315] N=892	-0.328 [0.410] N=725	-0.815 [0.571] N=554
WHO BMI-for-age z score	-0.330 [0.285] N=1,165	-0.312 [0.313] N=1,039	**-0.631* [0.346] N=891**	**-0.770* [0.452] N=723**	**-1.064* [0.583] N=558**
Stunted HAZ below -2	-0.025 [0.067] N=1,221	-0.021 [0.075] N=1,085	-0.041 [0.084] N=926	-0.058 [0.101] N=753	-0.077 [0.130] N=575
Underweight WAZ below -2	-0.009 [0.032] N=1,347	-0.033 [0.036] N=1,207	0.010 [0.041] N=1,025	-0.019 [0.048] N=831	-0.072 [0.048] N=627
Wasted WHZ below -2	**0.086** [0.040] N=1,168**	0.062 [0.046] N=1,040	**0.120** [0.054] N=892**	**0.092* [0.055] N=725**	**0.158** [0.060] N=554**
Overweight WHZ above +2	0.022 [0.062] N=1,168	0.022 [0.065] N=1,040	0.027 [0.074] N=892	-0.001 [0.093] N=725	-0.049 [0.121] N=554
Obese WHZ above +3	0.003 [0.046] N=1,168	0.012 [0.047] N=1,040	-0.018 [0.055] N=892	-0.013 [0.069] N=725	-0.112 [0.087] N=554
CIAF any stunting underweight or wasting	0.051 [0.076] N=1,148	0.001 [0.086] N=1,020	0.048 [0.102] N=873	-0.011 [0.118] N=709	0.045 [0.135] N=542
Below-normal BMI-for-age BMIZ below -2	**0.096*** [0.037] N=1,165**	**0.082* [0.046] N=1,039**	**0.140*** [0.052] N=891**	**0.134** [0.055] N=723**	**0.193*** [0.057] N=558**

Specification 5 Monthly fixed effects female and number of aid types (0-6) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.343	-0.169 [0.236] N=1,333	-0.217 [0.253] N=1,201	-0.225 [0.278] N=1,051	-0.249 [0.301] N=887	-0.309 [0.315] N=721
WHO weight-for-age z score	0.122	0.085 [0.140] N=1,466	0.169 [0.147] N=1,330	0.003 [0.147] N=1,158	-0.037 [0.162] N=975	-0.030 [0.177] N=785
WHO weight-for-height z score	0.386	0.140 [0.206] N=1,270	0.198 [0.213] N=1,146	0.122 [0.229] N=1,006	0.047 [0.257] N=849	-0.074 [0.290] N=689
WHO BMI-for-age z score	0.340	0.020 [0.228] N=1,263	0.066 [0.243] N=1,141	-0.086 [0.259] N=1,001	-0.163 [0.275] N=843	-0.200 [0.301] N=690
Stunted HAZ below -2	0.221 P=293	0.030 [0.045] N=1,333	0.044 [0.047] N=1,201	0.044 [0.049] N=1,051	0.049 [0.051] N=887	0.070 [0.058] N=721
Underweight WAZ below -2	0.057 P=86	-0.015 [0.025] N=1,466	-0.030 [0.026] N=1,330	-0.010 [0.027] N=1,158	-0.017 [0.031] N=975	-0.032 [0.033] N=785
Wasted WHZ below -2	0.107 P=137	0.054 [0.033] N=1,270	0.039 [0.036] N=1,146	0.063 [0.039] N=1,006	0.054 [0.039] N=849	**0.072* [0.042] N=689**
Overweight WHZ above +2	0.194 P=248	0.055 [0.043] N=1,270	0.062 [0.044] N=1,146	0.063 [0.047] N=1,006	0.057 [0.052] N=849	0.062 [0.058] N=689
Obese WHZ above +3	0.101 P=131	0.026 [0.033] N=1,270	0.037 [0.033] N=1,146	0.021 [0.038] N=1,006	0.034 [0.043] N=849	0.025 [0.048] N=689
CIAF any stunting underweight or wasting	0.302 P=376	0.087 [0.055] N=1,248	0.069 [0.059] N=1,124	0.099 [0.064] N=985	0.089 [0.066] N=830	**0.144** [0.069] N=674**
Below-normal BMI-for-age BMIZ below -2	0.109 P=140	0.056 [0.038] N=1,263	0.043 [0.042] N=1,141	0.066 [0.046] N=1,001	0.075 [0.046] N=843	**0.090* [0.048] N=690**

Specification 5 Monthly fixed effects female and number of aid types (0-6) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.281 [0.298] N=1,154	0.280 [0.337] N=1,022	0.396 [0.403] N=872	0.475 [0.472] N=708	0.627 [0.560] N=542
WHO weight-for-age z score	0.240 [0.216] N=1,273	**0.407* [0.229] N=1,137**	0.109 [0.246] N=965	0.094 [0.314] N=782	0.193 [0.392] N=592
WHO weight-for-height z score	0.049 [0.298] N=1,103	0.134 [0.307] N=979	-0.042 [0.345] N=839	-0.200 [0.430] N=682	-0.684 [0.610] N=522
WHO BMI-for-age z score	-0.190 [0.297] N=1,098	-0.150 [0.332] N=976	-0.497 [0.374] N=836	-0.650 [0.463] N=678	-0.940 [0.623] N=525
Stunted HAZ below -2	-0.025 [0.071] N=1,154	-0.012 [0.079] N=1,022	-0.036 [0.087] N=872	-0.057 [0.099] N=708	-0.045 [0.131] N=542
Underweight WAZ below -2	-0.013 [0.034] N=1,273	-0.037 [0.038] N=1,137	0.002 [0.041] N=965	-0.021 [0.052] N=782	-0.070 [0.049] N=592
Wasted WHZ below -2	**0.083** [0.042] N=1,103**	0.055 [0.048] N=979	**0.108* [0.056] N=839**	0.092 [0.058] N=682	**0.159** [0.066] N=522**
Overweight WHZ above +2	0.042 [0.064] N=1,103	0.045 [0.068] N=979	0.039 [0.076] N=839	0.004 [0.097] N=682	-0.021 [0.130] N=522
Obese WHZ above +3	0.005 [0.048] N=1,103	0.015 [0.050] N=979	-0.022 [0.057] N=839	-0.016 [0.073] N=682	-0.100 [0.094] N=522
CIAF any stunting underweight or wasting	0.048 [0.079] N=1,085	0.005 [0.089] N=961	0.043 [0.105] N=822	-0.023 [0.121] N=667	0.072 [0.138] N=511
Below-normal BMI-for-age BMIZ below -2	**0.094** [0.038] N=1,098**	0.075 [0.047] N=976	**0.126** [0.055] N=836**	**0.144** [0.057] N=678**	**0.206*** [0.060] N=525**

Specification 6 Monthly fixed effects female and any aid (0/1) No donut hole

Outcome	Mean	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	-0.343	-0.172 [0.239] N=1,333	-0.216 [0.255] N=1,201	-0.218 [0.279] N=1,051	-0.254 [0.306] N=887	-0.328 [0.315] N=721
WHO weight-for-age z score	0.122	0.087 [0.142] N=1,466	0.173 [0.149] N=1,330	0.001 [0.149] N=1,158	-0.028 [0.165] N=975	-0.023 [0.181] N=785
WHO weight-for-height z score	0.386	0.144 [0.204] N=1,270	0.199 [0.211] N=1,146	0.116 [0.225] N=1,006	0.059 [0.252] N=849	-0.060 [0.288] N=689
WHO BMI-for-age z score	0.340	0.025 [0.226] N=1,263	0.068 [0.242] N=1,141	-0.089 [0.257] N=1,001	-0.156 [0.272] N=843	-0.188 [0.297] N=690
Stunted HAZ below -2	0.221 P=293	0.030 [0.046] N=1,333	0.044 [0.048] N=1,201	0.043 [0.049] N=1,051	0.049 [0.052] N=887	0.071 [0.058] N=721
Underweight WAZ below -2	0.057 P=86	-0.015 [0.025] N=1,466	-0.030 [0.026] N=1,330	-0.008 [0.027] N=1,158	-0.018 [0.030] N=975	-0.032 [0.032] N=785
Wasted WHZ below -2	0.107 P=137	0.052 [0.033] N=1,270	0.038 [0.036] N=1,146	0.064 [0.039] N=1,006	0.052 [0.039] N=849	0.070 [0.042] N=689
Overweight WHZ above +2	0.194 P=248	0.055 [0.042] N=1,270	0.061 [0.043] N=1,146	0.060 [0.047] N=1,006	0.058 [0.052] N=849	0.062 [0.058] N=689
Obese WHZ above +3	0.101 P=131	0.026 [0.032] N=1,270	0.037 [0.033] N=1,146	0.020 [0.037] N=1,006	0.034 [0.043] N=849	0.024 [0.048] N=689
CIAF any stunting underweight or wasting	0.302 P=376	0.085 [0.055] N=1,248	0.068 [0.060] N=1,124	0.098 [0.064] N=985	0.088 [0.067] N=830	**0.142** [0.069] N=674**
Below-normal BMI-for-age BMIZ below -2	0.109 P=140	0.055 [0.038] N=1,263	0.043 [0.043] N=1,141	0.068 [0.046] N=1,001	0.073 [0.046] N=843	**0.087* [0.048] N=690**

Specification 6 Monthly fixed effects female and any aid (0/1) 12 month donut hole

Outcome	8 years	7 years	6 years	5 years	4 years
WHO height-for-age z score	0.266 [0.305] N=1,154	0.262 [0.339] N=1,022	0.392 [0.400] N=872	0.437 [0.506] N=708	0.572 [0.580] N=542
WHO weight-for-age z score	0.243 [0.217] N=1,273	**0.411* [0.231] N=1,137**	0.103 [0.249] N=965	0.117 [0.318] N=782	0.213 [0.400] N=592
WHO weight-for-height z score	0.053 [0.296] N=1,103	0.140 [0.305] N=979	-0.052 [0.339] N=839	-0.171 [0.438] N=682	-0.682 [0.619] N=522
WHO BMI-for-age z score	-0.184 [0.296] N=1,098	-0.143 [0.328] N=976	-0.506 [0.366] N=836	-0.638 [0.474] N=678	-0.934 [0.629] N=525
Stunted HAZ below -2	-0.022 [0.071] N=1,154	-0.006 [0.079] N=1,022	-0.031 [0.087] N=872	-0.044 [0.102] N=708	-0.029 [0.132] N=542
Underweight WAZ below -2	-0.013 [0.033] N=1,273	-0.036 [0.037] N=1,137	0.007 [0.042] N=965	-0.019 [0.051] N=782	-0.066 [0.049] N=592
Wasted WHZ below -2	**0.083** [0.042] N=1,103**	0.057 [0.048] N=979	**0.114** [0.056] N=839**	0.091 [0.058] N=682	**0.165** [0.066] N=522**
Overweight WHZ above +2	0.042 [0.064] N=1,103	0.046 [0.067] N=979	0.039 [0.076] N=839	0.012 [0.098] N=682	-0.018 [0.131] N=522
Obese WHZ above +3	0.007 [0.048] N=1,103	0.018 [0.050] N=979	-0.020 [0.057] N=839	-0.008 [0.073] N=682	-0.094 [0.094] N=522
CIAF any stunting underweight or wasting	0.051 [0.078] N=1,085	0.013 [0.089] N=961	0.055 [0.105] N=822	-0.010 [0.121] N=667	0.086 [0.139] N=511
Below-normal BMI-for-age BMIZ below -2	**0.094** [0.037] N=1,098**	0.078 [0.047] N=976	**0.134** [0.054] N=836**	**0.140** [0.057] N=678**	**0.207*** [0.060] N=525**

Notes: Cells report the OLS policy coefficient, birth-month-clustered standard error in brackets, and N. Exact child-month fixed effects for analysis months 24 to 59 are included but not displayed. Specification 2 adds female; specifications 3 and 4 separately add five household-income and five per-person-income groups; specifications 5 and 6 separately add the categorical number of aid types (0 as reference) and any aid. Neither aid control is combined with an income control. Aid specifications use only the six main questions; households with a blank or 99 on any question are excluded. Columns use symmetric 8- to 4-year bandwidths around the January 1987 cutoff. Every regression includes reform eligibility and split linear trends. Donut estimates exclude running-variable months -12 through +11 and show only the policy estimate, standard error, and N. Regressions use FAKTOR_FERT child survey weights. The sample retains one latest-born eligible child per household; all complete-DOB development respondents are included; raw ages below 24 map to 24 and above 59 map to 59. Significant policy coefficients are bold. P in the mean column is the unweighted number of positive outcomes in the no-donut 8-year sample. *** p<0.01, ** p<0.05, * p<0.10.