

A Certified Bivariate Entropy Inequality Related to Liu’s Entropy Bound

Kåre

Abstract

We prove, with certified computation, a bivariate entropy inequality that is suggested by Liu’s conditionally independent coupling method for the union-closed sets conjecture. The proof combines a Boppana-style Rolle method on the diagonal, a degree-27 Bernstein positivity certificate, and a finite Taylor-model interval-arithmetic verification of the off-diagonal critical-point structure. This version records the verified bivariate inequality and its computational certificates. It does not claim the union-closed corollary: the previously proposed reduction from the pointwise bivariate inequality to Liu’s full conditionally i.i.d. variational problem implicitly used an independence assumption that is not available in the conditionally i.i.d. coupling. The remaining task is to connect the certified bivariate inequality to the correct distributional optimization problem, or to replace it by a stronger distributional certificate. The code implementing all certificates is available at the repository listed in the appendix.

Status note. An earlier draft claimed that the bivariate inequality proved here implied the constant $0.382709087918735\dots$ for Frankl’s union-closed sets conjecture. Jingbo Liu pointed out that the entropy argument in that draft implicitly identified two different distributions of conditional probabilities. In Liu’s conditionally i.i.d. framework, the independent entropy term and the conditionally independent entropy term are integrated against different measures. The present version therefore withdraws the union-closed corollary and keeps only the certified bivariate inequality and the precise statement of the remaining distributional gap.

1 Introduction

A finite family $\mathcal{F}$ of sets is *union-closed* if

$$A, B \in \mathcal{F} \implies A \cup B \in \mathcal{F}. \tag{1.1}$$

Frankl’s union-closed sets conjecture, formulated in 1979, asserts that if $\mathcal{F}$ is finite, union-closed, and not equal to $\{\emptyset\}$, then some element belongs to at least half of the sets in $\mathcal{F}$ [1, 2]. Despite its elementary formulation, the conjecture has remained open for more than four decades. It is one of the central extremal problems concerning finite set systems.

A major breakthrough was made by Gilmer [3], who introduced an entropy method and proved the first absolute constant lower bound. In Gilmer’s argument, one samples

independent random sets A, B from $\mathcal{F}$, compares $H(A \cup B)$ with $H(A)$, and uses union-closedness to force a popular element. Gilmer's initial constant was 0.01. This was the first proof that there is an absolute $c > 0$ such that every finite nonempty union-closed family different from $\{\emptyset\}$ has an element appearing in at least a c -fraction of its members.

Shortly afterward, the constant was improved to

$$\frac{3 - \sqrt{5}}{2} = 0.3819660112 \dots \quad (1.2)$$

by several works, including Alweiss–Huang–Sellke [4], Chase–Lovett [5], Sawin [6], Cambie [8], Yu [9], and Pebody [10]. A central analytic ingredient in this line of work is the one-variable entropy inequality

$$h(x^2) \geq \varphi x h(x), \quad 0 \leq x \leq 1, \quad \varphi = \frac{1 + \sqrt{5}}{2}, \quad (1.3)$$

which goes back to Boppana's work on amplification of probabilistic Boolean formulas [11]. Chase and Lovett also showed that the constant $(3 - \sqrt{5})/2$ is optimal for an approximate union-closed formulation, thereby identifying a barrier for the plain independent two-sample entropy method.

Liu [7] introduced a further refinement based on conditionally independent couplings. Instead of taking two independent samples, one allows the two samples to be independent only after conditioning on an auxiliary random variable. This produces a richer class of coordinatewise couplings and leads to the numerical constant

$$0.3827090879 \dots, \quad (1.4)$$

slightly above the golden-ratio threshold. Liu proved analytically a strict improvement over the previous 0.38234-level constants, but the explicit value 0.38271... in his paper depends on additional numerically verified hypotheses. More precisely, Liu's explicit value requires two extra certificates: first, a positive-semidefinite condition for an infinite kernel matrix associated with the conditionally independent protocol; second, a global optimizer-structure hypothesis for a nine-dimensional variational problem. The optimizer is observed numerically to have a two-point boundary structure, but Liu does not prove this structure analytically.

The present paper studies the bivariate inequality that one obtains by taking the apparent two-point boundary optimizer suggested by Liu's computation and writing down the corresponding pointwise entropy certificate. Let

$$I(x, y) = xy(1 + (1 - x)(1 - y)). \quad (1.5)$$

We prove that, for explicit constants m_* and β_* ,

$$m_*((1 - \beta_*)h(xy) + \beta_*h(I(x, y))) \geq \frac{xh(y) + yh(x)}{2} \quad (0 \leq x, y \leq 1). \quad (1.6)$$

Here

$$m_* = 0.617290912081265 \dots, \quad 1 - m_* = 0.382709087918735 \dots \quad (1.7)$$

The inequality (1.6) is a certified bivariate entropy inequality related to Liu’s proposed constant. It is not, by itself, a proof of the corresponding union-closed lower bound. The reason is distributional: in Liu’s variational problem the independent term and the conditionally independent term are averaged with respect to different measures, whereas a pointwise inequality such as (1.6) controls both terms on the same pair (x, y) . A further distributional argument would be needed to recover the full union-closed consequence.

The proof of (1.6) has three parts. On the boundary of the square $[0, 1]^2$, the inequality is immediate from $m_* > 1/2$. On the diagonal $x = y$, we write $M(x) = G(x, x)$ and prove $M(x) \geq 0$ by a fifth-derivative sign argument: a symbolic computation reduces the sign of $M^{(5)}$ to the positivity of a degree-27 polynomial P , and we certify $P > 0$ on $[0, 1]$ by verifying the positivity of all 28 of its Bernstein coefficients. Off the diagonal, any negative interior minimum would be an off-diagonal critical point. Passing to variables

$$u = \frac{x + y}{2}, \quad v = \frac{x - y}{2},$$

we analyze the branch $F = \Gamma_v/v = 0$. Along this branch, the derivative of Γ_u is controlled by the Schur-complement identity

$$\frac{d}{du}\Gamma_u(u, v(u)) = \frac{\det D^2\Gamma(u, v(u))}{\Gamma_{vv}(u, v(u))}.$$

A Taylor-model interval computation certifies the required determinant signs on the two branch arms, while a separate interval-Newton and topological argument certifies that there are no additional components of $F = 0$. This rules out all off-diagonal critical points.

Computer-assisted inequalities have already played a role in the recent entropy approach to the union-closed conjecture. In particular, Alweiss–Huang–Sellke used computer verification in their treatment of the one-variable inequality underlying the $(3 - \sqrt{5})/2$ bound [4]. The present paper uses certified computation in the same spirit: all numerical claims are reduced to finite interval statements with explicit margins. The analytic part of the proof identifies the correct bivariate inequality and reduces the off-diagonal problem to branch isolation and determinant sign checks; the appendix records the finite certificates needed to make those reductions rigorous.

2 Definitions and constants

Throughout the paper we use natural logarithms. Define the binary entropy function by

$$h(t) = -t \log t - (1 - t) \log(1 - t), \quad 0 \leq t \leq 1, \quad (2.1)$$

with the convention $0 \log 0 = 0$. Let

$$L(t) = h'(t) = \log \frac{1 - t}{t}. \quad (2.2)$$

Define

$$I(x, y) = xy(1 + (1 - x)(1 - y)) = xy(2 - x - y + xy), \quad (2.3)$$

and

$$D(x) = I(x, x) = x^2(2 - 2x + x^2). \quad (2.4)$$

Let $x_* \in (0, 1)$ be the unique root of

$$x^4 - 2x^3 + 3x^2 - 1 = 0. \quad (2.5)$$

Equivalently,

$$x_*^2(2 + (1 - x_*)^2) = 1. \quad (2.6)$$

Set

$$p_* = \frac{h(x_*)}{h(x_*^2)}, \quad (2.7)$$

$$m_* = p_* x_*, \quad c' = 1 - m_*. \quad (2.8)$$

Numerically,

$$x_* = 0.690787593924988 \dots, \quad m_* = 0.617290912081265 \dots, \quad c' = 0.382709087918735 \dots. \quad (2.9)$$

The parameter β_* is defined by the stationarity condition

$$\frac{d}{dx} [m_*((1 - \beta_*)h(x^2) + \beta_*h(D(x))) - xh(x)]_{x=x_*} = 0. \quad (2.10)$$

Equivalently,

$$\beta_* = \frac{\frac{h(x_*^2)}{h(x_*)} \left(h'(x_*) + \frac{h(x_*)}{x_*} \right) - 2x_*h'(x_*^2)}{D'(x_*)h'(D(x_*)) - 2x_*h'(x_*^2)}. \quad (2.11)$$

Numerically,

$$\beta_* = 0.100052559862893 \dots. \quad (2.12)$$

We write

$$\alpha_* = 1 - \beta_*. \quad (2.13)$$

Define

$$G(x, y) = m_*(\alpha_*h(xy) + \beta_*h(I(x, y))) - \frac{xh(y) + yh(x)}{2}. \quad (2.14)$$

The main analytic result of the paper is the inequality

$$G(x, y) \geq 0 \quad (0 \leq x, y \leq 1). \quad (2.15)$$

We record the equality identities at (x_*, x_*) . From (2.5) and (2.4),

$$D(x_*) = 1 - x_*^2. \quad (2.16)$$

By entropy symmetry,

$$h(D(x_*)) = h(1 - x_*^2) = h(x_*^2). \quad (2.17)$$

Therefore

$$G(x_*, x_*) = m_*h(x_*^2) - x_*h(x_*) = 0, \quad (2.18)$$

where the last equality is (2.7) and (2.8). Finally, (2.10) says

$$\left. \frac{d}{dx} G(x, x) \right|_{x=x_*} = 0. \quad (2.19)$$

Since G is symmetric, (2.19) implies

$$\nabla G(x_*, x_*) = 0. \quad (2.20)$$

3 Boundary cases

We first prove nonnegativity on the boundary of the square. If $x = 0$, then $xy = 0$, $I(0, y) = 0$, and $xh(y) = 0$, hence

$$G(0, y) = 0. \quad (3.1)$$

Similarly,

$$G(x, 0) = 0. \quad (3.2)$$

If $x = 1$, then $xy = y$ and $I(1, y) = y$, so

$$G(1, y) = \left(m_* - \frac{1}{2} \right) h(y) \geq 0, \quad (3.3)$$

because $m_* > 1/2$. By symmetry,

$$G(x, 1) = \left(m_* - \frac{1}{2} \right) h(x) \geq 0. \quad (3.4)$$

Thus $G \geq 0$ on $\partial[0, 1]^2$, with equality on the axes and at the corner $(1, 1)$.

4 The diagonal inequality

Define

$$M(x) = G(x, x). \quad (4.1)$$

Using $I(x, x) = D(x)$, we have

$$M(x) = m_*(\alpha_* h(x^2) + \beta_* h(D(x))) - xh(x). \quad (4.2)$$

Lemma 1 (Diagonal nonnegativity). *For every $x \in [0, 1]$,*

$$M(x) \geq 0. \quad (4.3)$$

On $[m_, 1]$, equality occurs only at $x = x_*$ and $x = 1$. On $[0, 1]$, equality occurs at $x = 0$, $x = x_*$, and $x = 1$.*

Proof. A symbolic differentiation gives

$$M^{(5)}(x) = -\frac{2P(x)}{x^3(1-x)^4(1+x)^4(x^2-2x+2)^4(x^3-x^2+x+1)^4}. \quad (4.4)$$

The denominator is strictly positive on $(0, 1)$. The polynomial P is

$$P(x) = P_0(x) + m_*P_1(x) + m_*\beta_*P_2(x). \quad (4.5)$$

The three polynomials are

$$\begin{aligned} P_0(x) = & 3x^{27} - 36x^{26} + 200x^{25} - 666x^{24} + 1364x^{23} - 1360x^{22} - 1096x^{21} + 6288x^{20} \\ & - 9854x^{19} + 4312x^{18} + 11216x^{17} - 23492x^{16} + 15180x^{15} + 11376x^{14} - 29832x^{13} \\ & + 18608x^{12} + 9251x^{11} - 22292x^{10} + 9976x^9 + 6670x^8 - 9176x^7 + 1408x^6 \\ & + 2784x^5 - 1488x^4 - 400x^3 + 320x^2 - 32, \end{aligned} \quad (4.6)$$

$$\begin{aligned} P_1(x) = & 24x^{26} - 288x^{25} + 1860x^{24} - 8208x^{23} + 27296x^{22} - 71472x^{21} + 150320x^{20} \\ & - 254064x^{19} + 338288x^{18} - 333680x^{17} + 197272x^{16} + 27248x^{15} - 209888x^{14} \\ & + 233136x^{13} - 104048x^{12} - 45904x^{11} + 96120x^{10} - 46640x^9 - 12924x^8 \\ & + 25312x^7 - 6336x^6 - 5312x^5 + 3360x^4 + 640x^3 - 640x^2 + 64, \end{aligned} \quad (4.7)$$

$$\begin{aligned} P_2(x) = & 24x^{26} - 168x^{25} + 60x^{24} + 2832x^{23} - 14144x^{22} + 41712x^{21} - 99488x^{20} \\ & + 213456x^{19} - 389072x^{18} + 532880x^{17} - 458888x^{16} + 88048x^{15} + 372160x^{14} \\ & - 587200x^{13} + 435712x^{12} - 83408x^{11} - 223240x^{10} + 324792x^9 - 210324x^8 \\ & + 22720x^7 + 77632x^6 - 70096x^5 + 29760x^4 - 5760x^3 - 256x^2 + 192x + 64. \end{aligned} \quad (4.8)$$

The degree-27 Bernstein coefficients b_i of P on $[0, 1]$ have certified lower bounds shown in Table 1. Since the Bernstein basis is nonnegative on $[0, 1]$, the table implies

$$P(x) > 0 \quad (0 \leq x \leq 1). \quad (4.9)$$

Consequently,

$$M^{(5)}(x) < 0 \quad (0 < x < 1). \quad (4.10)$$

We first prove the claim on $[m_*, 1]$. The following signs are certified by interval evaluation:

$$\begin{aligned} M(m_*) &> 0, & M'(m_*) &< 0, & M''(m_*) &> 0, \\ M^{(3)}(m_*) &> 0, & M^{(4)}(m_*) &< 0. \end{aligned} \quad (4.11)$$

Numerically these are

$$\begin{aligned} M(m_*) &= 0.000777106330304900\dots, \\ M'(m_*) &= -0.0199407396705102\dots, \\ M''(m_*) &= 0.219173887997792\dots, \end{aligned}$$

Table 1: Certified lower bounds for the Bernstein coefficients of P .

i	lower bound	i	lower bound
0	11.45935667296576	14	4.610838565116360
1	11.898549817384106	15	4.964896105425193
2	12.078833739776622	16	5.668547978061419
3	11.876898948434980	17	6.798303977430921
4	11.307562491478990	18	8.465673305310538
5	10.464009676814568	19	10.840728674923076
6	9.465777139811940	20	14.191030026624640
7	8.423319403297192	21	18.945974671912115
8	7.421560841110654	22	25.804885986190340
9	6.518660405060565	23	35.924168951745940
10	5.753064074484635	24	51.256141180419560
11	5.152687277174934	25	75.205102681605100
12	4.742718854790539	26	114.019777429866760
13	4.550899799454681	27	180.158840956815370

$$\begin{aligned} M^{(3)}(m_*) &= 1.55739338128487\dots, \\ M^{(4)}(m_*) &= -4.41104974811261\dots \end{aligned} \quad (4.12)$$

Moreover,

$$M(x_*) = 0, \quad M'(x_*) = 0, \quad (4.13)$$

and interval evaluation gives

$$M''(x_*) = 0.317062949668787\dots > 0. \quad (4.14)$$

We also need the behavior at 1. Put $s = 1 - x$. As $s \downarrow 0$,

$$\begin{aligned} h(x^2) &= 2s \log \frac{1}{s} + O(s), \\ h(D(x)) &= 2s \log \frac{1}{s} + O(s), \\ xh(x) &= s \log \frac{1}{s} + O(s). \end{aligned} \quad (4.15)$$

Therefore

$$M(x) = (2m_* - 1)s \log \frac{1}{s} + O(s). \quad (4.16)$$

Since $2m_* - 1 > 0$, differentiating with respect to $x = 1 - s$ gives

$$\begin{aligned} M'(1^-) &= -\infty, \\ M''(1^-) &= -\infty, \\ M^{(3)}(1^-) &= -\infty, \\ M^{(4)}(1^-) &= -\infty. \end{aligned} \quad (4.17)$$

By (4.10), $M^{(4)}$ is strictly decreasing. Since $M^{(4)}(m_*) < 0$, we have

$$M^{(4)}(x) < 0 \quad (m_* \leq x < 1). \quad (4.18)$$

Hence $M^{(3)}$ is strictly decreasing. From $M^{(3)}(m_*) > 0$ and $M^{(3)}(1^-) = -\infty$, there is a unique

$$a \in (m_*, 1) \quad (4.19)$$

with $M^{(3)}(a) = 0$. Thus M'' increases on (m_*, a) and decreases on $(a, 1)$. Since $M''(m_*) > 0$ and $M''(1^-) = -\infty$, there is a unique

$$b \in (a, 1) \quad (4.20)$$

with $M''(b) = 0$.

Consequently M' increases on (m_*, b) and decreases on $(b, 1)$. We have $M'(m_*) < 0$, $M'(x_*) = 0$, and $M''(x_*) > 0$, so $x_* \in (m_*, b)$ and is the unique zero of M' in (m_*, b) . Also $M'(b) > 0$, while $M'(1^-) = -\infty$; hence there is a unique

$$d \in (b, 1) \quad (4.21)$$

with $M'(d) = 0$. The sign pattern is

$$M' < 0 \text{ on } (m_*, x_*), \quad M' > 0 \text{ on } (x_*, d), \quad M' < 0 \text{ on } (d, 1). \quad (4.22)$$

Since $M(m_*) > 0$, $M(x_*) = 0$, and $M(1) = 0$, it follows that

$$M(x) \geq 0 \quad (m_* \leq x \leq 1). \quad (4.23)$$

It remains to treat $[0, m_*]$. As $x \downarrow 0$,

$$\begin{aligned} h(x^2) &= 2x^2 \log \frac{1}{x} + O(x^2), \\ h(D(x)) &= 4x^2 \log \frac{1}{x} + O(x^2), \\ xh(x) &= x^2 \log \frac{1}{x} + O(x^2). \end{aligned} \quad (4.24)$$

Hence

$$M(x) = \gamma x^2 \log \frac{1}{x} + O(x^2), \quad \gamma = 2m_*(1 + \beta_*) - 1 > 0. \quad (4.25)$$

Thus $M(x) > 0$ for all sufficiently small positive x , $M'(0^+) = 0$, $M''(0^+) = +\infty$, $M^{(3)}(0^+) = -\infty$, and $M^{(4)}(0^+) = +\infty$.

Since $M^{(5)} < 0$, $M^{(4)}$ is strictly decreasing on $(0, m_*)$. From $M^{(4)}(0^+) = +\infty$ and $M^{(4)}(m_*) < 0$, there is a unique zero $c \in (0, m_*)$ of $M^{(4)}$. Thus $M^{(3)}$ increases on $(0, c)$ and decreases on (c, m_*) . Since $M^{(3)}(0^+) = -\infty$ and $M^{(3)}(m_*) > 0$, $M^{(3)}$ has exactly one zero $a_0 \in (0, c)$. Therefore M'' decreases on $(0, a_0)$ and increases on (a_0, m_*) .

The interval evaluation

$$M''(0.1) < 0 \quad (4.26)$$

combined with $M''(0^+) = +\infty$ and $M''(m_*) > 0$ implies that M'' has exactly two zeros $b_1 < b_2$ in $(0, m_*)$. Thus M' increases, then decreases, then increases. Since $M'(0^+) = 0$ and $M'(x) > 0$ for small $x > 0$, while $M'(m_*) < 0$, the function M' has exactly one zero $r \in (0, m_*)$. Hence M increases on $(0, r)$ and decreases on (r, m_*) . Since $M(0) = 0$ and $M(m_*) > 0$, we get

$$M(x) \geq 0 \quad (0 \leq x \leq m_*). \quad (4.27)$$

This completes the proof. $\square$

5 Off-diagonal critical points

Introduce

$$u = \frac{x+y}{2}, \quad v = \frac{x-y}{2}. \quad (5.1)$$

Thus

$$x = u + v, \quad y = u - v. \quad (5.2)$$

The feasible half-triangle is

$$0 < u < 1, \quad 0 < v < \min(u, 1-u). \quad (5.3)$$

Define

$$\Gamma(u, v) = G(u+v, u-v). \quad (5.4)$$

Since G is symmetric, Γ is even in v . For $v > 0$, define

$$F(u, v) = \frac{\Gamma_v(u, v)}{v}. \quad (5.5)$$

An off-diagonal critical point satisfies

$$F(u, v) = 0, \quad \Gamma_u(u, v) = 0, \quad v > 0. \quad (5.6)$$

Set

$$a = u^2 - v^2, \quad b = 2 - 2u + u^2 - v^2, \quad J = ab. \quad (5.7)$$

Thus $a = xy$, $b = 1 + (1-x)(1-y)$, and $J = I(x, y)$. Let

$$\Delta(u, v) = \det D^2\Gamma(u, v) = \Gamma_{uu}\Gamma_{vv} - \Gamma_{uv}^2. \quad (5.8)$$

Along a regular branch $v = v(u)$ of $F = 0$, one has $\Gamma_v(u, v(u)) = 0$. Differentiating gives

$$v'(u) = -\frac{\Gamma_{uv}}{\Gamma_{vv}}. \quad (5.9)$$

For

$$\Phi(u) = \Gamma_u(u, v(u)), \quad (5.10)$$

we therefore get

$$\Phi'(u) = \Gamma_{uu} - \frac{\Gamma_{uv}^2}{\Gamma_{vv}} = \frac{\Delta}{\Gamma_{vv}}. \quad (5.11)$$

This identity is the monotonicity mechanism for the off-diagonal analysis.

5.1 Endpoint positivity on the upper arc

Put

$$x = 1 - \varepsilon, \quad y = s. \quad (5.12)$$

Let

$$q = 2m_* - 1 = 0.234581824162530 \dots \quad (5.13)$$

On the upper branch, $F = 0$ is equivalent to $G_x = G_y$, hence

$$\Gamma_u = G_x + G_y = 2G_y. \quad (5.14)$$

Since

$$G_y = m_*(\alpha_* x L(xy) + \beta_* I_y(x, y) L(I(x, y))) - \frac{h(x) + xL(y)}{2}, \quad (5.15)$$

we have the exact identity

$$\Gamma_u = 2m_*(\alpha_* x L(xy) + \beta_* I_y(x, y) L(I(x, y))) - h(x) - xL(y). \quad (5.16)$$

Here

$$I_y(x, y) = x(2 - x - 2y + 2xy). \quad (5.17)$$

Lemma 2 (Endpoint positivity). *For every point of the upper arc with $0 < s = y \leq s_1 = 0.04$,*

$$\Gamma_u \geq \frac{2m_* - 1}{2} \log \frac{1}{s} > 0. \quad (5.18)$$

Proof. The endpoint branch satisfies

$$0 < \varepsilon(s) \leq s^2 \quad (0 < s \leq 0.04). \quad (5.19)$$

Indeed, from the explicit formula for $F = (G_x - G_y)/v$, one obtains

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} F(\varepsilon, s) &= +\infty, \\ F(s^2, s) &\leq -\frac{q}{2} \log \frac{1}{s} + 0.003 < 0, \end{aligned} \quad (5.20)$$

and

$$\partial_\varepsilon F(\varepsilon, s) \leq -\frac{1}{4s} < 0 \quad (0 < \varepsilon \leq s^2, 0 < s \leq 0.04). \quad (5.21)$$

Thus the unique endpoint solution satisfies (5.19).

Assume $0 < s \leq s_1$ and $0 < \varepsilon \leq s^2$. Then $x = 1 - \varepsilon \geq 1 - s^2$. Since $xy \leq s$ and L is decreasing,

$$L(xy) \geq L(s). \quad (5.22)$$

Moreover,

$$I_y = (1 - \varepsilon)(1 + \varepsilon - 2\varepsilon s) \geq 1 - s^2, \quad (5.23)$$

and

$$I(x, y) \leq s(1 + s^2). \quad (5.24)$$

Therefore

$$L(I(x, y)) \geq L(s(1 + s^2)). \quad (5.25)$$

Also $h(x) = h(\varepsilon) \leq h(s^2)$ and $-xL(y) \geq -L(s)$. Substituting these bounds into (5.16) gives

$$\Gamma_u \geq 2m_*(\alpha_*(1 - s^2)L(s) + \beta_*(1 - s^2)L(s(1 + s^2))) - h(s^2) - L(s). \quad (5.26)$$

Let

$$\delta(s) = L(s) - L(s(1 + s^2)). \quad (5.27)$$

Then

$$\Gamma_u \geq (2m_*(1 - s^2) - 1)L(s) - 2m_*\beta_*(1 - s^2)\delta(s) - h(s^2). \quad (5.28)$$

For $s \leq 1/25$,

$$L(s) \geq \log \frac{1}{s} - \frac{1}{24}, \quad (5.29)$$

while

$$\delta(s) \leq 2s^2, \quad (5.30)$$

and

$$h(s^2) \leq h(s_1^2). \quad (5.31)$$

Set

$$A_0 = 2m_*(1 - s_1^2) - 1, \quad C_0 = \frac{A_0}{24} + 4m_*\beta_*s_1^2 + h(s_1^2). \quad (5.32)$$

Numerically,

$$A_0 = 0.232606493243870\dots, \quad C_0 = 0.0219863330048360\dots \quad (5.33)$$

Thus

$$\Gamma_u \geq A_0 \log \frac{1}{s} - C_0. \quad (5.34)$$

Finally,

$$\left(A_0 - \frac{q}{2}\right) \log 25 - C_0 = 0.349200203430100\dots > 0. \quad (5.35)$$

Since $\log(1/s) \geq \log 25$, (5.18) follows. $\square$

Proposition 3 (Certified branch structure). *In the feasible region (5.3), the zero set $F(u, v) = 0$ consists of exactly two smooth arcs meeting at a single fold point. More precisely:*

(i) *The lower arc begins at the bifurcation point $(x_0, 0)$, with*

$$x_0 = 0.220715379651680\dots, \quad (5.36)$$

and ends at the fold

$$(u_f, v_f) = (0.552276058360800\dots, 0.434687834510229\dots). \quad (5.37)$$

Along this arc,

$$\Gamma_{vv} < 0, \quad \Delta < 0. \quad (5.38)$$

(ii) The upper arc begins at the boundary point $(x, y) = (1, 0)$ and ends at (u_f, v_f) . On the compact part $y \geq 0.04$,

$$\Gamma_{vv} > 0, \quad \Delta < 0. \quad (5.39)$$

For $0 < y \leq 0.04$, Lemma 2 gives $\Gamma_u > 0$.

(iii) The endpoint values satisfy

$$\Gamma_u(x_0, 0) > 0, \quad \Gamma_u(u_f, v_f) > 0.41. \quad (5.40)$$

Proof. The proposition is the certified interval statement proved in Appendix A. The proof uses one-variable interval Newton for the bifurcation point, interval Newton for the fold, Taylor-model interval arithmetic for (5.38) and (5.39), and a uniform band certificate for the approach to $(1, 0)$. $\square$

Lemma 4 (No off-diagonal critical points). *There is no point $(x, y) \in (0, 1)^2$ with $x \neq y$ such that*

$$G_x(x, y) = G_y(x, y) = 0. \quad (5.41)$$

Proof. By symmetry, assume $v > 0$. Any such critical point must satisfy $F = 0$ and $\Gamma_u = 0$. On the lower arc, Proposition 3 gives $\Gamma_{vv} < 0$ and $\Delta < 0$. Hence by (5.11), Γ_u is strictly increasing along the lower arc. Since $\Gamma_u(x_0, 0) > 0$, $\Gamma_u > 0$ throughout the lower arc.

On the upper arc with $y \leq 0.04$, Lemma 2 gives $\Gamma_u > 0$. On the compact part of the upper arc, Proposition 3 gives $\Gamma_{vv} > 0$ and $\Delta < 0$. Hence by (5.11), Γ_u decreases toward the fold. Since $\Gamma_u(u_f, v_f) > 0.41$, it remains positive on this compact part. Therefore no point on $F = 0$, $v > 0$, can satisfy $\Gamma_u = 0$. $\square$

6 Proof of the main inequality

Theorem 5 (Bivariate entropy inequality). *For every $x, y \in [0, 1]$,*

$$m_*((1 - \beta_*)h(xy) + \beta_*h(I(x, y))) \geq \frac{xh(y) + yh(x)}{2}. \quad (6.1)$$

Equality occurs on the axes, at (x_, x_*) , and at $(1, 1)$.*

Proof. This is exactly the assertion $G(x, y) \geq 0$. Since G is continuous on $[0, 1]^2$, it attains a minimum. If a minimizer lies on the boundary, Section 3 gives nonnegativity. If it lies in the interior and on the diagonal, Lemma 1 gives nonnegativity. If it lies in the interior and off the diagonal, it would be an off-diagonal critical point, contradicting Lemma 4. Thus $G \geq 0$ everywhere. The equality cases are the boundary equalities from Section 3 and the diagonal equality (x_*, x_*) from Lemma 1. $\square$

7 The remaining distributional gap

We record explicitly why Theorem 5 does not, by itself, prove the union-closed consequence suggested by Liu’s constant.

Let A be uniform on a union-closed family $\mathcal{F}$, and reveal coordinates in order. Let X_i denote the conditional probability that coordinate i is absent, given the previous coordinates. The chain rule gives

$$H(A) = \sum_i \mathbb{E}h(X_i). \quad (7.1)$$

For the independent two-sample coupling, if two histories give conditional absence probabilities x and y , the absence probability in the union is xy . Thus the independent entropy contribution involves an expectation of $h(XY)$ with respect to a product distribution of conditional probabilities.

For a conditionally i.i.d. coupling, however, the entropy term involving

$$I(x, y) = xy(1 + (1 - x)(1 - y)) \quad (7.2)$$

is not generally averaged with respect to the same product distribution. In Liu’s notation, the relevant optimization problem contains two numerator terms: one averaged with respect to a product measure of the marginal mixture, and one averaged with respect to a mixture of product measures. Schematically, if

$$P = (1 - q)P_0 + qP_1, \quad (7.3)$$

then the two terms have the form

$$\mathbb{E}_{P^{\otimes 2}}h(XY) \quad \text{and} \quad \mathbb{E}_{(1-q)P_0^{\otimes 2} + qP_1^{\otimes 2}}h(I(X, Y)). \quad (7.4)$$

These are generally expectations with respect to different distributions on (X, Y) .

The proof of Theorem 5 establishes the pointwise inequality

$$m_*((1 - \beta_*)h(xy) + \beta_*h(I(x, y))) \geq \frac{xh(y) + yh(x)}{2}. \quad (7.5)$$

Averaging (7.5) over a single joint distribution of (X, Y) is legitimate, but it does not imply a bound for the two-measure quantity in (7.4). In particular, the factorization

$$\mathbb{E} \frac{Xh(Y) + Yh(X)}{2} = \mathbb{E}X \mathbb{E}h(X) \quad (7.6)$$

requires an independence relation that is not available for the conditionally i.i.d. term.

Consequently, the present paper should be read as a certified proof of the bivariate inequality (7.5), together with a precise identification of the remaining obstacle. To obtain Liu’s full 0.382709... union-closed bound from this approach, one would need either

1. a distributional inequality matching the two expectations in (7.4), or
2. an independent proof that the worst case reduces to the same measure for both terms, such as a proof of the relevant optimizer-structure statement.

We do not prove either statement here.

A Computational certificates

This appendix records the finite certificates used in Sections 4 and 5. All interval computations use outward rounding and 50 decimal digits of precision. The implementation is available at

$$\text{https://github.com/kaare85-eng/liu-entropy-bound-certificate.} \quad (\text{A.1})$$

Proposition 6 (Constant enclosures). *The constants x_* , m_* , β_* , x_0 , u_f , and v_f are enclosed in intervals of radius less than 10^{-14} around the values displayed in (2.9), (2.12), (5.36), and (5.37). The enclosure for x_* certifies the unique root of (2.5) in $(0, 1)$, and the other constants are evaluated from their defining formulas.*

Proposition 7 (Bernstein positivity). *For*

$$P = P_0 + m_*P_1 + m_*\beta_*P_2,$$

all 28 degree-27 Bernstein coefficients of P on $[0, 1]$ are positive. The minimum certified lower bound is

$$\min_i b_i > 4.550899799454681. \quad (\text{A.2})$$

Consequently,

$$P(x) > 0 \quad (0 \leq x \leq 1).$$

Proposition 8 (Diagonal signs). *The interval evaluation certifies*

$$\begin{aligned} M(m_*) &> 0, & M'(m_*) &< 0, & M''(m_*) &> 0, & M^{(3)}(m_*) &> 0, & M^{(4)}(m_*) &< 0, \\ M''(x_*) &> 0, & M''(0.1) &< 0. \end{aligned} \quad (\text{A.3})$$

The numerical margins are those displayed in (4.12), (4.14), and (4.26).

Proposition 9 (Unique bifurcation). *Let*

$$H(u) = \frac{1}{2}\Gamma_{vv}(u, 0). \quad (\text{A.4})$$

Then

$$H(u) = L(u) + \frac{1}{2(1-u)} - m_*\alpha_*L(u^2) - 2m_*\beta_*(1-u+u^2)L(D(u)). \quad (\text{A.5})$$

The interval proof certifies

$$\begin{aligned} H(u) &< 0 & (0 < u < x_0), \\ H(u) &> 0 & (x_0 < u < 1), \end{aligned}$$

and a unique zero at x_0 . On the root box, $H'(u) > 0$.

Proposition 10 (Fold uniqueness and value). *The system*

$$F(u, v) = 0, \quad \Gamma_{vv}(u, v) = 0 \quad (\text{A.6})$$

has a unique solution in the fold box centered at (u_f, v_f) . At this solution,

$$\Gamma_u(u_f, v_f) > 0.41. \quad (\text{A.7})$$

Moreover, the interval evaluation gives

$$\Delta(u_f, v_f) < -31.17. \quad (\text{A.8})$$

Proof of branch isolation. We now explain how the preceding certificates are assembled into the branch isolation statement. Consider the zero set

$$\{(u, v) : F(u, v) = 0, v > 0\}$$

inside the feasible triangle

$$0 < u < 1, \quad 0 < v < \min(u, 1 - u).$$

Any connected component of this zero set must fall into one of the following topological alternatives.

First, it may be a closed loop in the interior. Such a loop must have at least two extrema in the u -projection, and hence at least two fold points, that is, at least two solutions of

$$F(u, v) = 0, \quad F_v(u, v) = 0.$$

On $F = 0$ this is equivalent to

$$F(u, v) = 0, \quad \Gamma_{vv}(u, v) = 0.$$

Proposition 10 certifies that there is only one such fold. Thus closed loops are excluded.

Second, a component may meet the diagonal $v = 0$. Since Γ is even in v , branches can bifurcate from the diagonal only where

$$\Gamma_{vv}(u, 0) = 0.$$

Proposition 9 certifies that the only such point is $u = x_0$. Thus the only possible diagonal birth is the already traced lower branch.

Third, a component may approach the boundary of the feasible triangle away from the corner $(x, y) = (1, 0)$. The side $y = 0$, away from $x = 1$, is excluded analytically: for fixed $0 < x < 1$,

$$F(x, y) \rightarrow -\infty \quad (y \rightarrow 0^+).$$

The side $x = 1$, away from $y = 0$, is also excluded analytically: for fixed $0 < y < 1$,

$$F(x, y) \rightarrow +\infty \quad (x \rightarrow 1^-).$$

The reflected sides $x = 0$ and $y = 1$ do not occur as distinct sides in the half-triangle $v > 0$; they are obtained by symmetry.

Fourth, a component may approach the corner $(x, y) = (1, 0)$. The upper arc has exactly this endpoint. The uniqueness of this approach is certified by a one-dimensional interval Newton band computation in the variable $s = y$. For $s \in [0.04, 0.08]$, the computation certifies a unique solution of $F(\varepsilon, s) = 0$, where $\varepsilon = 1 - x$, in the upper-branch tube. The eight s -bands have width 0.005; the maximal Newton contraction factor is 0.6296. For $0 < s < 0.04$, Lemma 2 gives the analytic endpoint estimate and identifies the same unique branch approaching the corner.

The two known arcs account for the unique diagonal bifurcation, the unique fold, and the unique corner approach. The alternatives above exhaust all possible connected components. Therefore there are no additional components of $F = 0$, $v > 0$, in the feasible triangle.

Proposition 11 (Branch isolation). *The zero set*

$$F(u, v) = 0, \quad v > 0,$$

in the feasible triangle (5.3) consists of exactly the two arcs stated in Proposition 3. The lower arc runs from the diagonal bifurcation point $(x_0, 0)$ to the fold (u_f, v_f) . The upper arc runs from the endpoint $(x, y) = (1, 0)$ to the same fold.

Proposition 12 (Lower arm determinant sign). *On the lower arc of $F = 0$, the Taylor-model interval certificate verifies*

$$\Gamma_{vv} < 0, \quad \Delta < 0. \quad (\text{A.9})$$

The bifurcation zone uses (u, w) -coordinates with $w = v^2$. The Taylor-model cover contains 23 boxes in the near-bifurcation zone and 56 boxes in the compact lower zone. The largest certified upper bound for Δ is

$$-0.00116054425855. \quad (\text{A.10})$$

Proposition 13 (Upper arm determinant sign). *On the compact upper arc with $y \in [0.08, y_f]$, Taylor-model interval arithmetic in the coordinates*

$$(\varepsilon, s) = (1 - x, y)$$

verifies

$$\Gamma_{vv} > 0, \quad \Delta < 0. \quad (\text{A.11})$$

The cover uses 42 adaptive boxes with maximum step size at most 0.001. The largest certified upper bound for Δ is

$$-31.1897707236, \quad (\text{A.12})$$

and the smallest certified lower bound for Γ_{vv} is

$$0.000394450122601. \quad (\text{A.13})$$

Proposition 14 (Endpoint positivity). *For the endpoint region $0 < y \leq 0.04$, Lemma 2 gives the analytic lower bound*

$$\Gamma_u \geq \frac{2m_* - 1}{2} \log \frac{1}{y} > 0. \quad (\text{A.14})$$

At the transition point $y = 0.04$, the computed value is

$$\Gamma_u = 0.745513582988373 \dots, \quad \frac{2m_* - 1}{2} \log 25 = 0.377544881375125 \dots \quad (\text{A.15})$$

Acknowledgements

This paper was developed with substantial AI assistance. The mathematical strategy, proof structure, and all certificate computations were carried out through extended interaction with large language models.

References

- [1] P. Frankl. Union-closed sets conjecture. Problem statement, 1979. See also D. B. West’s *Open Problems in Graph Theory and Combinatorics*.
- [2] H. Bruhn and O. Schaudt. The journey of the union-closed sets conjecture. *Graphs and Combinatorics*, 31(6):2043–2074, 2015.
- [3] J. Gilmer. A constant lower bound for the union-closed sets conjecture. arXiv:2211.09055, 2022.
- [4] R. Alweiss, B. Huang, and M. Sellke. Improved lower bound for Frankl’s union-closed sets conjecture. *The Electronic Journal of Combinatorics*, 31(3):P3.35, 2024.
- [5] Z. Chase and S. Lovett. Approximate union closed conjecture. arXiv:2211.11689, 2022.
- [6] W. Sawin. An improved lower bound for the union-closed sets conjecture. arXiv:2211.11504, 2022.
- [7] J. Liu. Improving the lower bound for the union-closed sets conjecture via conditionally IID coupling. arXiv:2306.08824, 2023.
- [8] S. Cambie. Better bounds for the union-closed sets conjecture using the entropy approach. arXiv:2212.12500, 2022.
- [9] L. Yu. Dimension-free bounds for the union-closed sets conjecture. *Entropy*, 25(5):767, 2023.
- [10] L. Pebody. Extension of a method of Gilmer. arXiv:2211.13139, 2022.
- [11] R. B. Boppana. Amplification of probabilistic Boolean formulas. In *26th Annual Symposium on Foundations of Computer Science (SFCS 1985)*, pages 20–29. IEEE, 1985.
- [12] J. Pulaj and K. Wood. Local configurations in union-closed families. *Experimental Mathematics*, to appear; arXiv:2301.01331, 2024.