Subexponential online mistake bounds for narrow isotropic Bayesian tanh networks

Dmitry Vaintrob Kaarel Hänni ChatGPT-6 Astra

November 2026

Abstract

We study Bayesian prediction by a scalar tanh network on the Boolean cube, with independent Gaussian weights, equal variances among the first-layer coordinates, an absolute output margin at every input, and exact conditioning on revealed labels. In the strong-prior limit, the total number of mistakes under an arbitrary nonanticipating input sequence is controlled by the number of exact Gaussian symmetry orbits below the target's minimum affine margin cost. If that cost is K, we prove an upper bound $\exp(O(K^{1/3}\log(K+1)))[1+K\log(d+2)]$ on cumulative posterior error probabilities. It applies to posterior sampling, posterior majority, and a fixed confidence criterion, at every fixed scalar depth. For one-neuron targets with all weights bounded by one this is $\exp(O(d^{1/3}\log d))$; for the AND of all d inputs it is $\exp(O(d^{2/3}\log d))$. Thus neither target class admits $\exp(\Omega(d))$ expected sampled mistakes, or that many majority or confidence failures, in this scalar architecture with the exact isotropic prior. A second argument covers every fixed learner width n at one hidden layer, for depth-two targets with a width-one implementation whose weights are at most one in absolute value. Its budget is $\exp(O_n(d^{n/(n+2)}\log^4(d+2)))$. These bounds are not polynomial or uniform in growing learner width. The proof uses coefficient separation, low-dimensional packing, and exact equality of function masses within symmetry orbits; it does not assume integer minimum energies or uniformity across a whole energy level.

1 The learner and the question

A polynomial sample bound for independent training examples does not itself bound all the mistakes made when examples arrive in an arbitrary order. The latter question is especially delicate when a strong prior successively favors different minimum-energy functions. A large number of nearly tied but distinct energies can matter, and equal energy alone does not imply equal Bayesian mass.

We examine this question for an exact isotropic Gaussian prior in the scalar architecture. Our conclusion is an upper bound: the target classes specified below cannot incur exponentially many expected sampled mistakes, majority errors, or confidence failures as a function of input dimension. A polynomial upper bound, or a superpolynomial lower bound below our upper bound, remains open for these target classes. Section 6 separately extends the result to every fixed hidden width at depth two, by encoding higher moments of networks averaged over exact input symmetries.

Fix an input dimension $d \geq 1$ and a depth $L \geq 2$. There are $L-1$ hidden tanh neurons, one at each hidden layer, no separate internal biases, and one linear output. Write

$$h_1(x) = \tanh(a \cdot (1, x)), \tag{1}$$

$$h_j(x) = \tanh(v_j h_{j-1}(x)) \quad (2 \leq j \leq L-1), \tag{2}$$

$$z(x) = b h_{L-1}(x), \qquad x \in X = \{-1, 1\}^d. \tag{3}$$

For $L = 2$ the internal weights v_j are absent. Independently sample

$$a_i \sim N(0, \sigma^2/d) \quad (0 \le i \le d), \qquad v_j, b \sim N(0, \sigma^2).$$

Before training, condition on the global event $|z(x)| > 1/2$ for every $x \in X$. After each revealed label, condition on agreement with that label. All conditioning events considered below have positive probability because a strict-margin target implementation exists. The *strong-prior limit* sends σ to zero at fixed d, L, after conditioning on each history.

The target $f : X \to \{-1, 1\}$ is fixed. At round t, choose x_t before the fresh posterior draw, make a prediction, and then reveal $f(x_t)$. The query may depend on all past queries, labels, and predictions. Let

$$q_t = \mathbb{P}\{\operatorname{sign} z(x_t) \neq f(x_t) \mid \text{revealed history}\}.$$

Posterior sampling draws a fresh classifier each round. Posterior majority chooses the more probable label, with any fixed tie convention. A strict-2/3 confidence failure means that the correct label has probability at most 2/3. These are different prediction criteria, but the same bound on $\sum_t q_t$ controls all three.

Every represented Boolean function is an affine threshold function. Its full-cube affine margin cost is

$$Q(g) = \min\left\{\|a\|_2^2 : g(x)a \cdot (1, x) \ge 1 \text{ for every } x \in X\right\}. \tag{4}$$

The minimum exists and is unique. It is at least one: averaging the squared scores of a unit-margin representative over the cube gives $\|a\|^2 \ge 1$.

2 Main bound

Fix $K \ge Q(f)$ and define

$$p = d + 1, \qquad B = \lceil K \rceil, \qquad R = \min\{p, \lfloor (3K)^{1/3} \rfloor\}, \tag{5}$$

$$J = \sum_{r=1}^{R} [B(1 + K)]^r, \qquad T = J[1 + B \log(2p)]. \tag{6}$$

All logarithms are natural.

Theorem 1. *For every $\beta > 0$ there is $\sigma_0 > 0$ such that, for every $0 < \sigma < \sigma_0$ and every nonanticipating input process,*

$$\sum_{t \ge 1} q_t \le T + \beta 2^d \tag{7}$$

pathwise. This single scale threshold works for all histories, input orders, and time horizons. Consequently, with $A = T + \beta 2^d$,

1. *the expected total posterior-sampling mistakes are at most A;*
2. *the total posterior-majority mistakes are at most $2A$;*
3. *the total strict-2/3 confidence failures are at most $3A$;*
4. *for every $\delta \in (0, 1)$, posterior-sampling mistakes M satisfy*

$$\mathbb{P}\{M > (e - 1)A + \log(1/\delta)\} \le \delta.$$

The same conclusions with $\beta = 0$ hold for every simultaneous subsequential strong-prior limit of the posteriors at all realizable label histories. In particular,

$$T \le \exp(O(K^{1/3}\log(K+1)))[1 + K\log(d+2)]. \tag{8}$$

The constants in this last inequality are absolute. Depth affects the required scale σ_0, not this budget.

The theorem does not assert that the posterior limit is unique. There are finitely many partial label assignments, so every sequence of positive scales has a subsequence along which all their posteriors converge simultaneously. The stated bound applies to each resulting prediction system.

Corollary 2 (Bounded-weight scalar targets)**.** *Suppose f has a scalar implementation at the specified depth with every weight of absolute value at most one and signed output margin greater than $1/2$ on the full cube. Then $Q(f) \le d+1$, and the strong-prior bounds in Theorem 1 are*

$$\exp(O(d^{1/3}\log(d+2))).$$

For sufficiently small positive scale the same bound, plus one, follows by choosing $\beta = 2^{-d}$.

This includes odd majority at $L = 2$, with implementation $z(x) = \tanh(\sum_i x_i)$ and teacher width one. At large scalar depth the bounded-weight premise may be impossible because repeated tanh contraction reduces the output; the corollary is conditional on the stated implementation, rather than an assertion that such targets exist at every scalar depth.

Corollary 3 (AND)**.** *Let $f(x) = +1$ exactly when every input coordinate is $+1$. Then*

$$Q(f) = d + (d-1)^2 = d^2 - d + 1.$$

At every fixed scalar depth, the exact isotropic strong-prior learner has cumulative posterior error at most

$$\exp(O(d^{2/3}\log(d+2)))$$

on every nonanticipating input sequence, with the corresponding sampling, majority, and confidence consequences. This rules out $\exp(\Omega(d))$ expected sampled mistakes, majority errors, or confidence failures for this scalar AND learner. It does not assert that scalar width meets a separate bounded-weight teacher-width condition for AND.

3 Neural energy and affine margin cost

The Gaussian energy is

$$\mathcal{E}(a, v, b) = d\|a\|^2 + \sum_{j=2}^{L-1} v_j^2 + b^2.$$

Let U_g be the event $g(x)z(x) > 1/2$ for every cube input. Absorb all scalar signs into a, using oddness of tanh, so that $v_j, b > 0$ without changing energy. For $u > 0$, let $H_v(u)$ be the output of the hidden scalar chain at first-layer preactivation u. It is positive and strictly increasing in u.

Lemma 4. *The infimum energy of U_g is*

$$E_L(g) = \Phi_{d,L}(Q(g)), \qquad \Phi_{d,L}(Q) = \min_{u>0,\, v_j>0} \left\{ dQu^2 + \sum_{j=2}^{L-1} v_j^2 + \frac{1}{4H_v(u)^2} \right\}. \tag{9}$$

The minimum is attained and $\Phi_{d,L}$ is strictly increasing on $Q \geq 1$. For $L = 2$, this specializes to

$$\Phi_{d,2}(Q) = \min_{u>0}\{dQu^2 + \tfrac{1}{4}\coth^2 u\}.$$

Moreover,

$$\lim_{\sigma\downarrow 0} \sigma^2 \log \mathbb{P}_\sigma(U_g) = -E_L(g)/2. \tag{10}$$

Proof. An affine representative with minimum signed preactivation u has squared norm at least $u^2Q(g)$. Conversely, scale the optimizer in (4) by u and choose $b = 1/(2H_v(u))$ to achieve weak output margin; increasing b slightly achieves strict margin. This proves the infimum formula.

On a bounded objective sublevel, u, every v_j, and $b = 1/(2H_v(u))$ are bounded above. No u or v_j can approach zero on that sublevel: continuity of the finite hidden chain would then send $H_v(u)$ to zero, contradicting the bound on b. Thus a minimizing sequence has an interior convergent subsequence, and the minimum is attained. At a minimizer for $Q_2 > Q_1$, the difference between the two objective values is $d(Q_2 - Q_1)u^2 > 0$. Evaluating the Q_1 minimum at the same point proves strict monotonicity.

Finally, the Gaussian density has exponent $-\mathcal{E}/(2\sigma^2)$. The standard finite-dimensional Gaussian small-noise upper bound for a closed set and lower bound for an open set give (10). Their energy infima agree here: the closure of U_g is exactly its weak-margin version, since multiplying its readout by $1 + \epsilon$ approaches every weak-margin parameter from within U_g. □

Let $\mathcal{A} = \{g : Q(g) \leq Q(f)\}$. Finiteness of the represented Boolean class and strict monotonicity imply

$$\frac{\mathbb{P}_\sigma(U_{\mathcal{A}^c})}{\mathbb{P}_\sigma(U_f)} \longrightarrow 0. \tag{11}$$

After any history consistent with f, the posterior denominator is at least the raw mass of U_f. Thus, at one sufficiently small scale, posterior mass outside $\mathcal{A}$ is at most β uniformly over all histories. Global-margin normalization cancels in this ratio.

4 Coefficient separation and a small orbit count

Extend g to the homogeneous odd threshold function on $\{-1, 1\}^p$ given by

$$F(y) = y_0 g(y_1/y_0, \ldots, y_d/y_0).$$

Its unit-margin constraints are exactly the original affine constraints and their antipodes.

Lemma 5 (Separated optimal magnitudes)**.** *In the unique minimum-norm unit-margin representative of F, each nonzero absolute coefficient is at least one, and any two distinct nonzero absolute coefficients differ by at least one. If their distinct values are $\lambda_1 < \cdots < \lambda_r$, with multiplicities n_j, then*

$$\lambda_j \geq j, \qquad Q(g) = \sum_j n_j \lambda_j^2 \geq r^3/3, \qquad \sum_j n_j \leq Q(g).$$

Proof. An inessential coordinate can be removed by averaging the two constraints differing only in that coordinate. Its optimal coefficient is therefore zero. An essential coordinate has a cube edge with oppositely labeled endpoints. Their unit-margin constraints force the corresponding coefficient to have absolute value at least one.

Orient two nonzero coefficients positively by coordinate sign flips. If their distinct values differ by less than one, swapping their input coordinates changes every score by less than two. Opposite

labels with unit margin would require a score difference at least two, so the function is invariant under that swap. The swapped coefficient vector is another feasible representative. Averaging the two strictly decreases squared norm, contradicting optimality. This proves separation. Now $\lambda_j \geq j$, and $\sum_{j=1}^{r} j^2 \geq r^3/3$ proves the remaining assertions. □

A symmetry orbit means an orbit of functions under signed permutations of the p augmented coordinates. This is larger than ordinary input permutation symmetry, because the augmented constant coordinate can be exchanged with an input coordinate.

Proposition 6 (Orbit count by packing)**.** *The functions of cost $Q(g) \leq K$ occupy at most J symmetry orbits, with J from (6). Each such orbit has at most $(2p)^B$ functions.*

Proof. Orient the unique minimum representative positively and sort its distinct coefficient magnitudes. By Lemma 5, it has some $r \leq R$ groups, with positive multiplicities $n_1, \ldots, n_r \leq B$ and support size $s = \sum_j n_j \leq K$. Fix this multiplicity list. Its function orbits have canonical representatives with positive magnitudes $\lambda_1 < \cdots < \lambda_r$, each repeated according to the list, and the remaining coordinates zero. Distinct orbits give distinct canonical Boolean functions.

Associate to a canonical representative the vector

$$u = (\sqrt{n_1}\lambda_1, \ldots, \sqrt{n_r}\lambda_r) \in \mathbb{R}^r, \qquad \|u\| \leq \sqrt{K}.$$

For two distinct canonical functions there is an augmented cube point where their labels differ. If c_j is the sum of its input signs in group j, then $|c_j| \leq n_j$, and the two unit-margin scores differ by at least two. For the two vectors u, u',

$$2 \leq \left| \sum_j \frac{c_j}{\sqrt{n_j}} (u_j - u'_j) \right| \leq \sqrt{s}\, \|u - u'\| \leq \sqrt{K}\, \|u - u'\|.$$

These vectors are therefore $2/\sqrt{K}$ separated in the radius-$\sqrt{K}$ ball. Disjoint open balls of radius $1/\sqrt{K}$ about them fit into the radius-$(\sqrt{K} + 1/\sqrt{K})$ ball. Comparing volumes bounds their number by $(1 + K)^r$.

There are at most B^r multiplicity lists of length r. Sum the product $B^r(1 + K)^r$ over $1 \leq r \leq R$ to obtain J. Finally a coefficient multiset with support s has at most $(2p)^s$ signed placements in p positions. Since $s \leq B$, this bounds each orbit size. □

An alternative elementary count avoids volume comparison. Round every optimal magnitude to a multiple of $1/(4B)$. The worst score perturbation is at most $s/(8B) \leq 1/8$, so all signs are preserved. There are at most $\lceil 4B^{3/2} \rceil$ rounded magnitudes and B possible multiplicities per group. The rounded list determines the function orbit, giving the slightly weaker count $\sum_{r \leq R} [B \lceil 4B^{3/2} \rceil]^r$ with the same asymptotic exponent. Neither count assumes integer minimum energies or enumerates the active constraints of the margin quadratic program.

5 Exact Gaussian symmetry and online elimination

Let S be a signed permutation matrix on $\mathbb{R}^p$. For each $x \in X$, write

$$S^\top(1, x) = \epsilon_S(x)(1, \tau_S(x)), \qquad \epsilon_S(x) \in \{-1, 1\}.$$

The map τ_S is a cube bijection: S permutes antipodal pairs of augmented cube vertices. Oddness of every hidden layer gives

$$z_{Sa,v,b}(x) = \epsilon_S(x) z_{a,v,b}(\tau_S(x)).$$

The transformation $a \mapsto Sa$ preserves the raw isotropic Gaussian law and the global absolute-margin event. It maps whole function events bijectively to their transformed events. Consequently all functions in one symmetry orbit have exactly equal prior mass at every positive scale. After exact label conditioning, surviving members of that orbit remain equiprobable. No assumption is made about the relative masses of different orbits, even at equal energy.

Lemma 7 (Shrinking uniform blocks)**.** *Partition a finite hypothesis class into fixed blocks. Suppose a consistent prediction distribution is uniform inside the surviving portion of each block. Its weights on different blocks may change arbitrarily with the history. A block of initial size m contributes at most $1 + \log m$ to the sum of posterior error probabilities on any input sequence.*

Proof. If the block has $n > 0$ survivors and the revealed label eliminates a of them, its error contribution is at most a/n. If $a < n$, then

$$\frac{a}{n} \le \log n - \log(n - a).$$

These terms telescope until the block is exhausted or the sequence ends. Its final exhausting deletion contributes at most one. The estimate is pathwise and does not depend on the block's varying total posterior mass. □

Proof of Theorem 1. By (11), choose a scale threshold so that the posterior outside $\mathcal{A}$ has mass at most β after every history. The class $\mathcal{A}$ is a union of at most J exact Gaussian symmetry orbits, each of size at most $(2p)^B$. Lemma 7 bounds their combined cumulative contribution by T. Hypotheses outside $\mathcal{A}$ contribute at most β per first appearance. They contribute zero on repeats because all surviving functions agree with the previously revealed label. There are at most 2^d first appearances, proving (7).

For sampled mistake indicators X_t, conditional on the past and current query,

$$\mathbb{E}[e^{X_t} \mid \text{past}, x_t] = 1 + (e - 1)q_t \le e^{(e-1)q_t}.$$

The resulting exponential supermartingale gives the stated tail at each finite horizon, hence for total mistakes by monotone convergence. Tonelli's theorem gives the expected-mistake bound. A majority mistake requires $q_t \ge 1/2$, while a strict-2/3 confidence failure requires $q_t \ge 1/3$.

In a simultaneous posterior limit at all realizable histories, the outside mass is zero and equality among surviving orbit members persists. Apply the same pathwise block argument with $\beta = 0$. Finally $J \le R[B(1 + K)]^R$, $R = O(K^{1/3})$, and $B(1 + K) = O((K + 1)^2)$, proving (8). □

6 Every fixed width at one hidden layer

The affine reduction above does not extend to several hidden neurons. Nevertheless, exact input-permutation symmetry combined with a higher-order approximation gives a subexponential result at every fixed hidden width. In this section there is one hidden layer of width n, and

$$z(y) = b^\top \tanh(Ay), \qquad y \in \{-1, 1\}^p, \quad p = d + 1.$$

The original cube is the section $y_0 = 1$. The canonical Gaussian energy and variances are

$$\mathcal{E}_n(A, b) = \frac{d}{n}\|A\|_F^2 + n\|b\|^2, \quad A_{ji} \sim N(0, n\sigma^2/d), \quad b_j \sim N(0, \sigma^2/n).$$

As before, condition first on absolute output margin greater than $1/2$ everywhere and then on exact training labels. Write $E_n(g)$ for the energy infimum of a function event.

Theorem 8 (Fixed-width shallow networks)**.** *Fix $n \geq 1$. Suppose the target has a depth-two, width-one implementation in which every weight has absolute value at most one and the signed output margin is greater than $1/2$ everywhere. At learner width n, the strong-prior posterior has, on every nonanticipating input sequence,*

$$\sum_t q_t \leq \exp\Big(O_n\Big(d^{\frac{n}{n+2}} \log^4(d+2)\Big)\Big). \tag{12}$$

The constant may depend on the fixed learner width n. The same bound, plus one, holds for all sufficiently small positive scales. Expected posterior-sampling mistakes, majority mistakes, confidence failures, and high-probability sampled mistakes obey the consequences of Theorem 1.

More generally, for fixed n and fixed integer $q \geq 1$, a target energy bound $E_n(f) \leq K$ gives a strong-prior budget with logarithm at most

$$O_{n,q}\left(\left[1 + \left(\frac{K^2}{d} + K^{1+1/(q+1)}\right)^{\frac{n}{n+2}}\right] \log(d+K+2)\right). \tag{13}$$

For the targets of the first assertion, whose energy is $K = O(d)$, each fixed $\epsilon > 0$ and a sufficiently large fixed Taylor order give the alternative exponent

$$O_{n,\epsilon}(d^{n/(n+2)+\epsilon} \log(d+2)).$$

The fixed-width hypothesis matters. This is not a bound uniform over widths that grow with d. For widths $n = 2$ and $n = 3$, the exponents of d in (12) are respectively $1/2$ and $3/5$, up to the displayed logarithms. For $n = 1$, the scalar argument already gives the sharper logarithmic factor in Corollary 2.

6.1 Close input columns force an exact Boolean symmetry

Every function with $E_n(g) \leq K$ has a weak-margin representative of that energy. Indeed, the energy is coercive and its weak-margin cell is closed; increasing the readout slightly shows that weak and strict infima agree. Define

$$B = \sqrt{K/n}, \qquad E_0 = nK/d.$$

The representative satisfies $\|b\| \leq B$ and $\sum_i \|A_i\|^2 \leq E_0$, where A_i denotes its ith column. The map $s \mapsto b^\top \tanh(s)$ is B-Lipschitz.

Choose a radius $\rho \leq 1/(8B)$, select a maximal ρ-separated subset of the columns, and assign every column to a center within distance ρ. Thus each group lies in a radius-ρ ball. Swapping two coordinates from the same group changes the preactivation by at most 4ρ and the output by at most $4B\rho \leq 1/2$. Two opposite labels with absolute margin at least $1/2$ would require an output difference at least one. Hence the Boolean function is invariant under this swap, and therefore under every permutation within each group.

Let there be r groups. The centers are original columns, so their squared norms sum to at most E_0. At least $r/2$ lie in the radius-$\sqrt{2E_0/r}$ ball. Packing radius-$\rho/2$ balls about these centers gives

$$\frac{r}{2} \leq \left(1 + \frac{2\sqrt{2E_0/r}}{\rho}\right)^n.$$

If $r > 2^{n+1}$, subtracting one from the nth root of the left side loses at most a factor two. Rearranging, and including the complementary case, gives

$$r \leq C_n \left[1 + (E_0\rho^{-2})^{n/(n+2)}\right]. \tag{14}$$

Lemma 9 (Permutation moments and Taylor approximation)**.** *Let $A \in \mathbb{R}^{n\times p}$ have its columns partitioned into groups, with every group contained in a Euclidean ball of radius ρ. For a fixed $y \in \{-1,1\}^p$, independently permute its coordinates within each group to obtain Y. Write $S = AY$, $\mu = \mathbb{E}S$, and $U = S - \mu$. Then, for every row j, every real t, and every integer $s \geq 1$,*

$$\mathbb{E}e^{tU_j} \leq \exp(8p\rho^2 t^2), \tag{15}$$

$$\mathbb{E}|U_j|^s \leq (8\sqrt{s}\sqrt{p}\,\rho)^s. \tag{16}$$

Consequently, if $\|b\|_2 \leq B$ and $q \geq 1$ is an integer, the function

$$P(y) = \sum_{j=1}^{n} b_j \sum_{a=0}^{q} \frac{\tanh^{(a)}(\mu_j)}{a!}\mathbb{E}U_j^a$$

satisfies

$$\left|\mathbb{E}[b^T \tanh(AY)] - P(y)\right| \leq \sqrt{n}B(16\sqrt{q+1}\sqrt{p}\,\rho)^{q+1}. \tag{17}$$

Proof. Within a group of size m, center the scalar entries in row j as $z_1, \ldots, z_m$, so $\sum_i z_i = 0$. Their range has length at most 2ρ, since the original group lies in a radius-ρ ball. The centered contribution to U_j is twice the sum of a uniformly selected k-subset of these entries, where k is the number of positive coordinates of y in the group. Its moment generating function is

$$\frac{e_k(e^{2tz_1}, \ldots, e^{2tz_m})}{\binom{m}{k}} \leq \left(\frac{1}{m}\sum_{i=1}^{m} e^{2tz_i}\right)^k.$$

Here e_k is the elementary symmetric polynomial. The inequality follows by repeatedly averaging two of its nonnegative arguments while preserving their sum: the coefficient of their product is nonnegative, so this operation cannot decrease e_k, and its maximum at a fixed sum occurs when all arguments are equal. The cases $k = 0, m$ cause no difficulty.

A uniform draw Z from the centered entries has mean zero and lies in an interval of length at most 2ρ. The elementary bounded-variable Hoeffding lemma gives $\mathbb{E}e^{2tZ} \leq \exp(2\rho^2 t^2)$; even the looser range bound 4ρ gives $\exp(8\rho^2 t^2)$, which we use here. Thus the group contribution has moment generating function at most $\exp(8m\rho^2 t^2)$. Independence across groups proves (15).

Chernoff's argument gives $\mathbb{P}(|U_j| > u) \leq 2\exp(-u^2/(32p\rho^2))$. For an integer $\ell \geq 1$, integration yields

$$\mathbb{E}|U_j|^{2\ell} \leq 2(32p\rho^2)^\ell \ell! \leq 2(32\ell p\rho^2)^\ell.$$

Take $\ell = \lceil s/2 \rceil \leq s$ and use monotonicity of L^r norms. The resulting bound is at most $(8\sqrt{s}\sqrt{p}\rho)^s$, proving (16).

For completeness, $|\tanh^{(s)}(u)| \leq 2^s s!$ for all $u \in \mathbb{R}$: write the sth derivative as $P_s(\tanh u)$, start from $P_0(v) = v$, and use $P_{s+1}(v) = (1-v^2)P_s'(v)$. Its coefficient ℓ^1 norm is at most $2^s s!$ by induction, since $\deg P_s = s+1$. Taylor's theorem therefore bounds the order-q remainder by $2^{q+1}|U_j|^{q+1}$. Apply (16), sum with weights $|b_j|$, and use $\|b\|_1 \leq \sqrt{n}B$ to obtain (17). □

Choice of radius. Set $s = q+1$ and

$$\rho = \min\left\{\frac{1}{8B}, \frac{(8\sqrt{n}B)^{-1/s}}{16\sqrt{s}\sqrt{p}}\right\}.$$

Then swaps within a group change the original output by at most $4B\rho \leq 1/2$, giving exact Boolean invariance under all such swaps. The Taylor error in (17) is at most $1/8$. If the original output has

signed margin at least $1/2$ on the full cube, then the permutation average has the same margin, and the approximating function P has signed margin at least $3/8$ on every input. Moreover

$$\rho^{-2} \le C_{n,q}\left(B^2 + pB^{2/(q+1)}\right),$$

where $C_{n,q}$ is a constant depending only on the two fixed integers.

6.2 A finite description of the symmetrized Taylor polynomial

Fix the multiplicities $m_1, \dots, m_r$ of the column groups. For row j and group g, record the power sums

$$P_{g,j,\ell} = \sum_{i \in G_g} A_{ji}^{\ell}, \qquad 1 \le \ell \le q.$$

Together with the n readout weights, these are $nqr + n$ real descriptors. Their number is linear in r when n, q are fixed.

We verify that they determine every moment used in the Taylor approximation. For one group, write its scalar entries as $a_1, \dots, a_m$ and suppose the queried sign vector has k positive coordinates in that group. A uniform permutation is equivalent to choosing a uniform k-subset, with indicators ξ_i. Put $V = \sum_i a_i \xi_i$ and $P_\ell = \sum_i a_i^\ell$. For $h \le q$,

$$\mathbb{E}V^h = \sum_{\pi \in \mathcal{P}([h])} \frac{(k)_{|\pi|}}{(m)_{|\pi|}} \sum_{\substack{i_B \in [m] \ (B \in \pi) \\ \text{all distinct}}} \prod_{B \in \pi} a_{i_B}^{|B|}. \tag{18}$$

Here $(t)_v$ is a falling factorial, and terms with $|\pi| > m$ are zero. The formula partitions ordered index tuples according to their coincidence pattern. Each coefficient $(k)_v/(m)_v$ lies in $[0,1]$.

If a partition has block sizes $s_1, \dots, s_v$, its distinct-index sum is a polynomial in $P_1, \dots, P_h$. Explicitly, inclusion–exclusion on coincidence patterns gives

$$\sum_{i_1,\dots,i_v \text{ distinct}} \prod_{t=1}^{v} a_{i_t}^{s_t} = \sum_{\tau \in \mathcal{P}([v])} \left[\prod_{C \in \tau} (-1)^{|C|-1}(|C|-1)!\right] \prod_{C \in \tau} P_{\sum_{t \in C} s_t}. \tag{19}$$

This is the usual partition-lattice inclusion–exclusion; it also follows inductively by subtracting all nontrivial coincidence patterns from the unrestricted product of power sums. Thus (18) has degree at most h in the descriptors, and its coefficient magnitudes and number of terms are bounded by constants depending only on h.

The group's centered preactivation is

$$U_g = 2\left(V - \frac{k}{m} P_1\right).$$

Its moments through order q are likewise polynomials of degree at most their order, with coefficients uniformly bounded in m, k. Different groups are permuted independently. Expanding the moments of $U = \sum_g U_g$ therefore gives polynomials of degree at most h in all group descriptors, with at most $C_q r^q$ terms and coefficients bounded by C_q, for $h \le q$. The mean is

$$\mu_j = \sum_g \left(\frac{2k_g}{m_g} - 1\right) P_{g,j,1}.$$

Consequently the function

$$T(y) = \sum_{j=1}^{n} b_j \sum_{h=0}^{q} \frac{\tanh^{(h)}(\mu_j)}{h!} \mathbb{E} U_j^h \tag{20}$$

is completely determined by the group sizes, these descriptors, and the number k_g of positive signs in each queried group. It is not required to be a neural network: its role is to encode the Boolean function, which is already determined by its uniformly separated sign.

6.3 Uniform quantization of the descriptors

Let $A_* = \sqrt{E_0}$ and

$$H = p \max\{1, A_*\}^q.$$

Every power-sum descriptor has absolute value at most H, and every readout weight has absolute value at most B. Quantize all of them on a common grid with mesh $\eta \le 1$. We give the elementary uniform continuity estimate needed for this step, including its dependence on the growing dimension p.

For fixed n, q, the displayed moment polynomials have degree at most q, at most $C_q p^q$ monomials, and coefficients bounded by C_q. On the descriptor box of radius $H + 1$, changing each coordinate by at most η changes any such polynomial by at most

$$C_q p^q (H + 1)^q \eta.$$

This follows by telescoping the factors of each monomial; the constants absorb its degree and number of terms. The mean μ_j changes by at most $p\eta$. Every derivative of tanh of any fixed order is bounded on the real line: inductively it is a polynomial in tanh, using $(\tanh)' = 1 - \tanh^2$. Applying the mean value theorem to the coefficients $\tanh^{(h)}(\mu_j)$ and then changing the readout weights therefore gives

$$|T(y) - \widetilde{T}(y)| \le C_{n,q}(1 + B) p^{q+1} (H + 1)^q \eta \tag{21}$$

for every full-cube input y, with one constant independent of all group sizes and sign counts. This deliberately loose bound also controls the magnitudes of the moment polynomials appearing when the readout is changed.

Choose

$$\eta = \min\left\{1, [16 C_{n,q}(1 + B) p^{q+1} (H + 1)^q]^{-1}\right\}.$$

The output perturbation is at most $1/16$. If the Taylor approximation has signed margin at least $3/8$, its quantized version retains margin at least $5/16$ and exactly the same Boolean function.

There are at most

$$M = 3 + \left\lceil \frac{2 \max\{H, B\}}{\eta} \right\rceil$$

grid values per descriptor. Since $E_0 = nK/d$, $B = \sqrt{K/n}$ and $p = d + 1$, for fixed n, q this obeys

$$\log M = O_{n,q}(\log(d + K + 2)).$$

One can put every group consecutively by an input permutation. The group sizes have at most p^r possibilities, and the quantized descriptors have at most M^{nqr+n} possibilities. Choose one grouping and one quantized description deterministically for each function orbit. A shared description would

define the same canonical Boolean function by (20); hence two distinct orbits cannot share it. The number of exact signed-augmented-coordinate symmetry orbits below energy K is therefore at most

$$J_{n,q}(d,K) \le \sum_{r=1}^{\min\{p,\lceil R_{n,q}\rceil\}} p^r M^{nqr+n},$$

where the group count $R_{n,q}$ comes from the energy packing estimate. This count permits invalid descriptor lists and repeated encodings, both of which only overcount.

6.4 Orbit count, Gaussian posterior, and the growing Taylor order

The radius above and (14) imply

$$r \le C_{n,q}\left[1+\left(\frac{K^2}{d}+K^{1+1/(q+1)}\right)^{n/(n+2)}\right].$$

Here $p/d \le 2$ has been absorbed into the constant. The finite descriptor count therefore gives

$$\log J_{n,q}(d,K) \le O_{n,q}\left(\left[1+\left(\frac{K^2}{d}+K^{1+1/(q+1)}\right)^{n/(n+2)}\right]\log(d+K+2)\right).$$

Every exact signed-augmented-coordinate orbit has at most $2^p p!$ functions, and all its functions have equal Gaussian mass at every positive scale. The proof is the same parameter isometry used in the scalar case, applied simultaneously to the first-layer columns. Gaussian small-noise rates give a posterior tail at most β outside $E_n(g) \le E_n(f)$ after every target-consistent history. Lemma 7 consequently yields

$$\sum_t q_t \le J_{n,q}(d,K)[1+\log(2^p p!)] + \beta 2^d.$$

The polynomial orbit-size factor is absorbed in (13); the tail vanishes in a strong-prior limit. This proves the fixed-q assertion with all posterior prediction criteria.

For clarity we now track the constants needed when the Taylor order grows. The number of partitions of a set of size q is at most q^q, and every factorial occurring in (19) is at most $q!$. Expanding (18), the centering binomial, and the sum over independent groups therefore bounds the relevant coefficients and the constants in (21) by

$$\exp(O(q\log(q+1)))$$

apart from a factor depending on the fixed n. One way to see that combining groups does not produce an extra exponential in q^2 is to expand the h ordered factors directly: there are at most r^h assignments of their group indices, and the product of their groupwise partition counts and factorial bounds is at most $\exp(O(h\log(h+1)))$, since the group degrees sum to $h \le q$. The tanh derivative bound in Lemma 9 has the same factorial size.

Such a target, with every weight bounded by one, is an augmented signed majority with affine optimum a_* satisfying $\|a_*\|^2 \le p$. Set $a = a_*/\sqrt{d}$ and $b_* = 1/[2\tanh(1/\sqrt{d})]$. Since $\tanh(t) \ge t/2$ for $0 \le t \le 1$, its weak-margin scalar energy is

$$d\|a\|^2 + b_*^2 \le p + d = 2d+1.$$

Cloning it n times with readouts b_*/n preserves $\mathcal{E}_n$ exactly. Thus we may take $K = 2d+1$, so $E_0 = O_n(1)$ and $B = O_n(\sqrt{p})$.

Choose $q = \lceil \log(p+1) \rceil$ and $s = q + 1$. The explicit radius gives

$$\rho^{-2} = \max\{64B^2,\, 256sp(8\sqrt{n}B)^{2/s}\} = O_n(p \log(p+1)).$$

Consequently

$$r = O_n\Big((p \log(p+1))^{n/(n+2)}\Big).$$

Since $H = p \max\{1, \sqrt{E_0}\}^q$, we have $\log(H+1) = O_n(\log(p+1))$. The explicit quantization mesh and the coefficient bounds just established give

$$\log M = O_n(\log^2(p+1)).$$

There are $nqr + n = O_n(r \log(p+1))$ descriptors and at most p^r group-size lists. Hence

$$\log J_{n,q}(d, 2d+1) = O_n\Big(p^{n/(n+2)} \log^{3+n/(n+2)}(p+1)\Big).$$

This implies the slightly simpler bound (12), completing the theorem. No growing-order approximation is needed for the separately stated fixed-q conclusion.

7 Target costs and limitations

Proof of Corollary 2. With all scalar weights bounded by one, the absolute final output is no larger than $|\tanh(a \cdot (1, x))|$. Thus every initial affine score has absolute value greater than $u = \operatorname{arctanh}(1/2) > 1/2$. Extend to the augmented homogeneous function and delete inessential coordinates by averaging signed-margin constraints. Each remaining coefficient has magnitude greater than u, from an edge with opposite labels, and at most one.

After orienting coefficients positively, any two differ by less than $1 - u < u$. Swapping their coordinates changes any score by less than $2u$, and so cannot change a label. The Boolean function on its active augmented coordinates is therefore symmetric, monotone, and odd. It is unweighted majority on an odd number s of coordinates: symmetry and monotonicity make it a cardinality threshold, and oddness places the threshold at the midpoint and excludes even s. Unit coefficients give margin one, and essentiality gives the reverse lower bound, so $Q(f) = s \le p$. □

Proof of Corollary 3. By permutation symmetry and uniqueness, the minimizing input coefficients of AND are all equal, say t. Let the bias be c. Comparing the all-positive input with an input having exactly one negative coordinate gives $t \ge 1$. The latter input imposes $c \le -1 - (d-2)t$, while the former imposes $c \ge 1 - dt$.

For $d \ge 2$, the minimum permitted absolute bias at a fixed $t \ge 1$ is $1 + (d-2)t$. Hence

$$Q(f) \ge dt^2 + [1 + (d-2)t]^2 \ge d + (d-1)^2.$$

Equality is attained by $t = 1, c = 1 - d$. For $d = 1$ the target is a dictator and has cost one, agreeing with the formula. Substitute this cost into Theorem 1. □

These results concern the actual posterior with exact coordinate-variance symmetry. Breaking that symmetry can split a large equal-mass family into distinct energy levels; the present proof then no longer applies. The orbit-count estimates proved here are subexponential but not polynomial. Exact symmetry alone has not supplied a polynomial mistake bound.

The architectural restrictions remain substantive. The scalar theorem allows arbitrary fixed depth; the wider-network theorem allows every fixed width but only one hidden layer and a specified simple target class. Multiple hidden layers and widths growing with dimension introduce additional issues. An arbitrary-width polynomial upper bound, or a bounded-resource exponential counterexample in that regime, remains unresolved by this paper.

Verification status. The results have conventional proofs. They are not Lean formalized. The quantitative orbit count strengthens an earlier research argument based on active-constraint enumeration; all ingredients needed for the present proof are included above.