A certified margin-repair bound for cubic parity predictors

One arbitrary bit, twenty-six symmetric bits, and parity-block lifts

Kaarel Hänni Dmitry Vaintrob OpenAI Codex

16 October 2026

Abstract

We prove an explicit lower bound for repairing the errors of a globally gapped parity predictor in a cusp variogram model. On twenty-seven bits, the predictor may be any polynomial of degree at most three that is symmetric in twenty-six coordinates, with arbitrary dependence on the remaining bit. If its parity error is at most one third, its mean repair energy is at least $77247413969/13589544960 > 5.68$. The result allows unbounded amplitudes and arbitrary intermediate radial sign changes. A second bound, exceeding 4.69, holds for error at most 0.34. The proof combines an analytic repair inequality with exhaustive, exactly verified rational certificates for the relevant sign-cone orbits. Equal parity-block substitution extends the result to a structured family in arbitrarily large dimension, including nonsymmetric amplitude representatives of degree fraction at most $1/9$. This is a theorem about the stated repair model and sign-pattern family, not a general posterior or neural-network sample-complexity theorem.

1 Model and result

Write $\Omega_K = \{-1,1\}^K$, with uniform measure, Hamming distance d_H, and parity $f(u) = \prod_{i=1}^K u_i$. A *head* is a real function P on Ω_K satisfying $|P(u)| \geq 1$. Put $g = \operatorname{sgn} P$, $B = |P|$, and $E = \{u : g(u) \neq f(u)\}$. Degree always means degree of the multilinear Walsh expansion.

Fix $0 < \beta \leq 1/2$. A cell has points $(u,h) \in \Omega_K \times \mathbb{Z}_{\geq 0}$. For a finitely supported real charge q with $\sum q = 0$, define its squared dual norm by

$$V_\beta(q) = -\sum_{u,h,v,j} q_{u,h} q_{v,j} \big(d_H(u,v) + h + j\big)^\beta. \tag{1}$$

This is nonnegative. Indeed, the integral representation of s^β reduces it to a positive integral of quadratic forms with kernel $e^{-td_H(u,v)}e^{-th}e^{-tj}$, which is positive semidefinite. The zero-total-charge condition cancels the constant term.

A repair $a(u,h)$ has squared energy $\mathcal{E}$ if

$$\Big|\sum q_{u,h} a(u,h)\Big|^2 \leq \mathcal{E} V_\beta(q) \quad \text{for every such } q. \tag{2}$$

Equivalently, its least possible energy is the supremum of these ratios; zero-variance contrasts have the usual limiting interpretation. At a wrong center $u_0 \in E$, an admissible repair satisfies

$$\begin{aligned} |P(u) + a(u,h)| &\geq 1 && \text{for every } (u,h), \\ \operatorname{sgn}(P(u_0) + a(u_0,0)) &= f(u_0), \\ \operatorname{sgn}(P(u) + a(u,h)) &= g(u) && \text{for every } u \text{ and all sufficiently large } h. \end{aligned}$$

The last condition is eventual anchoring to the head signs. There is no restriction on intermediate signs. Repairs at different wrong centers may be chosen independently. Their population cost is $\mathbb{E}[\mathbf{1}_E(u)\mathcal{E}_u]$; it is not normalized by the error probability. Define

$$C_0(\beta) = \frac{4}{2 - 2^\beta} \le 4 + 2\sqrt{2}.$$

Theorem 1. *Let $K = 27$. Suppose P has degree at most three, is invariant under permutations of coordinates $2, \dots, 27$, has global margin $|P| \ge 1$, and has parity error at most $1/3$. Every collection of admissible repairs satisfies*

$$\mathbb{E}[\mathbf{1}_E\mathcal{E}_u] \;\ge\; L_* := \frac{77247413969}{13589544960} = 5.684326752394806\ldots > \frac{11}{2} > \frac{C_0(\beta)}{2}.$$

The conclusion holds uniformly for $0 < \beta \le 1/2$, without an amplitude bound. If the error assumption is relaxed to $R \le 17/50 = 0.34$, the conclusion becomes

$$\mathbb{E}[\mathbf{1}_E\mathcal{E}_u] \ge L_{.34} := \frac{1785556483667}{380507258880} = 4.692568780219008\ldots > \frac{9}{2} > \frac{C_0(\beta)}{2}.$$

The proof below is computer assisted only in a finite rational inequality. A standard-library Python checker and the rational certificate data accompany this paper. The checker does not call an optimization package.

2 A repair inequality valid through all radial sign changes

Lemma 1. *Let u be wrong and have $n \ge 12$ correct active neighbors. Set*

$$L_u = 2 + B(u) - \frac{1}{n} \sum_{v \sim u,\ v \notin E} B(v), \qquad T = \frac{11}{3}.$$

Every admissible repair at u satisfies

$$\mathcal{E}_u \ge \Phi_u(P) := \max\left\{4,\ 2T\left[B(u) - \frac{1}{n}\sum_{v\sim u,\ v\notin E} B(v)\right] + 4T - T^2\left(1 + \frac{3}{5n}\right)\right\}. \tag{3}$$

When $n < 12$, the bound $\Phi_u(P) = 4$ remains valid.

Proof. The variance of the contrast between (v, h) and $(v, h+1)$ is

$$v_h = 2(2h+1)^\beta - (2h)^\beta - (2h+2)^\beta.$$

It is positive and decreasing. For $h \ge 1$, the integral formula

$$v_h = \beta(1-\beta)\int_0^1 \int_0^1 (2h + s + t)^{\beta - 2}\, ds\, dt$$

gives $v_h \le [4(2h)^{3/2}]^{-1}$. Also v_1 increases with β on the stated interval, since both factors in the integrand increase, so

$$v_h \le v_1 \le 2\sqrt{3} - \sqrt{2} - 2 < \frac{1}{20}.$$

A radial sign change requires a contrast of magnitude at least two. Thus every such change costs at least $4/v_h$, and every change at $h \geq 1$ costs more than 80. Since a repaired wrong center must eventually regain its original sign, some radial change occurs. Consequently

$$\mathcal{E}_u \geq 4/v_0 = C_0(\beta) \geq 4. \tag{4}$$

If all correct neighbors have their head signs at shell $H \geq 1$, their field signs are opposite to the repaired center sign: an active edge reverses parity, and u is wrong while its chosen neighbors are correct. The center-to-uniform-neighbor contrast of the repair therefore has magnitude at least $(L_u)_+$. Its exact variance is

$$F_H = (2 - 2^\beta)(H+1)^\beta + \frac{(2H+2)^\beta - (2H)^\beta}{n}.$$

At $H = 1$, writing $x = 2^\beta$,

$$F_1 = x(2-x) + \frac{x^2 - x}{n} \leq 1 + \frac{3}{5n},$$

because $1 \leq x \leq \sqrt{2}$ and $x^2 - x \leq 2 - \sqrt{2} < 3/5$.

If $\mathcal{E}_u < 80$, eventual anchoring and the radial-change bound force all active points to have their head signs already at shell one. Cauchy–Schwarz in (2) gives $(L_u)_+^2 \leq \mathcal{E}_u F_1$, hence $\mathcal{E}_u \geq 2tL_u - t^2 F_1$ for every $t \geq 0$.

If $\mathcal{E}_u \geq 80$, put

$$H = \left\lfloor \frac{1}{2}(\mathcal{E}_u/16)^{2/3} \right\rfloor + 1.$$

A change at any shell $h \geq H$ would cost at least $16(2h)^{3/2} > \mathcal{E}_u$. Thus shell H is anchored. Since $n \geq 12$,

$$\begin{aligned} F_H &\leq \sqrt{(H+1)/2} + 1/12 \\ &\leq \sqrt{1 + \tfrac{1}{4}(\mathcal{E}_u/16)^{2/3}} + 1/12 \leq \frac{\mathcal{E}_u}{4T^2}. \end{aligned}$$

For the last inequality divide by $\mathcal{E}_u$: both terms decrease, so it is enough to check $\mathcal{E}_u = 80$. There the left side before division is less than $17/12$, whereas $80/(4T^2) = 180/121 > 17/12$. The contrast bound now gives $(L_u)_+ \leq \mathcal{E}_u/(2T)$, and therefore $\mathcal{E}_u \geq 2TL_u \geq 2TL_u - T^2 F_1$. Combining the two energy regimes with (4) proves (3). □

For a fixed sign pattern, Φ_u is a maximum of affine functions of the head amplitudes, hence convex. This fact will also permit the block extension.

3 An exhaustive finite proof

Write $x \in \{0, 1\}$ for the first coordinate's negative-bit indicator and $h \in \{0, \ldots, 26\}$ for the remaining Hamming weight. A symmetric degree-three head has precisely the form

$$P(x, h) = A_3(h) + (-1)^x B_2(h),$$

where the subscripts bound ordinary polynomial degrees. Each of its two fiber sign sequences has at most three transitions. There are exactly

$$2 \sum_{j=0}^{3} \binom{26}{j} = 5904$$

such sequences. For one sequence define

$$d(g) = 2^{-26} \sum_{h=0}^{26} \binom{26}{h} (-1)^h g(h).$$

The full parity advantage is $(d(g_0) - d(g_1))/2$. Integer binomial arithmetic enumerates exactly 1460 ordered pairs with advantage at least $1/3$. The relaxed error cutoff $17/50$ requires advantage at least $8/25$ and gives 2256 pairs.

Lemma 2. *A pair of sign sequences with at most three transitions each is realizable by a globally gapped head of the displayed form if and only if it is not the case that both fibers have three transitions and opposite right-end signs.*

Proof. A fiber with three transitions has three real roots inside $(0, 26)$ and degree exactly three. Its leading-coefficient sign is therefore its sign at 26. The two fiber polynomials must have equal cubic coefficients, proving necessity. For sufficiency, put one root in each required transition interval. A fiber with fewer than three transitions can be completed to degree three by roots outside $[0, 26]$. Choosing the parity of the number placed to the right permits either leading-coefficient sign while retaining the desired signs on the integer interval. Scale both fibers to the same sufficiently large leading coefficient. Their difference has degree at most two, and their integer-node margins are at least one. □

Exactly 762 of the accurate pairs satisfy this criterion. The transformations $h \mapsto 26 - h$ and $(g_0, g_1) \mapsto (-g_1, -g_0)$ preserve the problem. They reduce the feasible pairs to 193 orbits. For the relaxed cutoff there are 1176 feasible pairs and 298 orbits; the feasibility proof is unchanged.

For each representative we use the seven integer-valued basis functions

$$(-1)^{ix} \binom{h}{j}, \qquad i \in \{0, 1\}, \quad i + j \le 3.$$

The polynomial coefficients and one cost variable at every wrong orbit are the linear-program variables. The constraints are the 54 signed margin inequalities, cost at least four, and the affine bound in (3) where the correct-neighbor multiplicity is at least twelve. The objective weights are $\binom{26}{h}/2^{27}$. Every coefficient is rational.

Proposition 1. *For every representative at either error cutoff, the companion data contain an exactly nonnegative rational dual vector. It annihilates the seven head coefficients and reproduces the negative objective coefficients on the cost variables. Its exact objective is at least L_* for error at most $1/3$, and at least $L_{.34}$ for error at most $17/50$.*

Proof. This is a finite rational certificate, checked by the accompanying program. For clarity, if the constraints are $Az \le b$ and the objective is $c^T z$, the checker verifies $y \ge 0$ and $A^T y = -c$. It then obtains $c^T z \ge -b^T y$. All identities and bounds are checked over $\mathbb{Q}$. The smallest certified bounds are exactly $77247413969/13589544960$ and $1785556483667/380507258880$, respectively. The checker separately re-enumerates all sign pairs and checks orbit coverage, so no floating optimization output is used as proof. □

Theorem 1 follows from Lemma 1, exhaustive coverage, and Proposition 1.

4 Parity-block lifts and scope

Theorem 2. *Partition* $27b$ *active coordinates into* 27 *blocks of size* $b \geq 1$. *Suppose a head* P *has degree at most* $3b$, *global margin one, and parity error at most* $1/3$. *Assume only its* sign *depends on the* 27 *block parities and is invariant under permutations of the last* 26 *blocks. Its amplitudes may depend arbitrarily on all physical coordinates. Then the conclusion of Theorem 1 holds with the same constant* L_*. *The error-*$17/50$ *conclusion also holds with constant* $L_{.34}$.

Proof. Average P over even coordinate flips within each block and over permutations of the last 26 blocks. These transformations preserve its sign pattern and the signed margin constraints. The resulting head is a function of block parities. An invariant Walsh monomial uses either every coordinate of a block or none; thus the degree bound $3b$ makes the averaged head an outer polynomial of degree at most three.

An outer wrong point with n correct neighbors corresponds to nb correct physical neighbors. The uniform-neighbor contrast has variance at most $1 + 3/(5nb) \leq 1 + 3/(5n)$. Hence the same coarse rational loss used in the certificate is a valid physical loss whenever $n \geq 12$. It is convex within the fixed sign cone and invariant under the averaging group. Jensen's inequality therefore lower-bounds the original mean loss by the loss of the averaged head. The relevant finite certificate applies to that head and yields L_* or $L_{.34}$. □

The block theorem has active degree fraction at most $1/9$, but its sign-pattern hypothesis is essential. Averaging arbitrary sign patterns need not preserve accuracy or margin. Neither theorem transfers the cell model to an adaptive finite-dimensional posterior; such a transfer, and a result for unrestricted low-degree sign patterns, are separate questions.

5 Reproducibility

The companion directory contains:

- `one_bit_symmetric_k27r3_universal_exact_certificates.json`, the rational data for error at most $1/3$;
- `one_bit_symmetric_k27r3_risk034_exact_certificates.json`, the data for error at most $17/50$;
- `verify_one_bit_symmetric_certificates.py`, the independent standard-library checker;
- `certify_one_bit_symmetric_phi_universal.py`, the optional SciPy/SymPy certificate generator.

Place the certificate data and checker together and run

```
python verify_one_bit_symmetric_certificates.py
python verify_one_bit_symmetric_certificates.py --risk034
```

The checker verifies the fiber count 5904, the pair/feasibility/orbit counts at both cutoffs, all rational dual identities, and the stated lower bounds. The generator is not required to verify the result.