


B
Seeds (n = 1,176)
PCoA 2 (8.4%)
0.50
0.25
0.00
−0.25
R² = 8.3%, p = 0.001
−0.50
−0.25
0.00
0.25
PCoA 1 (17.7%)
C
PCoA 2 (8.4%)
0.50
0.25
0.00
−0.25
R² = 6.3%, p = 0.001
−0.50
−0.25
0.00
0.25
PCoA 1 (17.7%)
D
Seedlings (n = 551)
PCoA 2 (10.0%)
0.25
0.00
−0.25
R² = 16.2%, p = 0.001
−0.25
0.00
0.25
PCoA 1 (11.9%)
E
PCoA 2 (10.0%)
0.25
0.00
−0.25
R² = 7.3%, p = 0.001
−0.25
0.00
0.25
PCoA 1 (11.9%)


C
Seeds (n = 1,176)
PCoA 2 (8.4%)
0.50
0.25
0.00
−0.25
R² = 2.6%, p = 0.001
−0.50
−0.25
0.00
0.25
PCoA 1 (17.7%)
D
Seedlings (n = 551)
PCoA 2 (10.0%)
0.25
0.00
−0.25
R² = 19.2%, p = 0.001
−0.25
0.00
0.25
PCoA 1 (11.9%)
Study
Chesneau et al. 2022 (PRJEB45079)
metaSEED (PRJEB6729)
Chandel et al. 2022 (PRJNA810761)
Kim et al. 2022 (PRJNA735565)
Cui et al. 2023 (PRJNA784432)
Nunes et al. 2022 (PRJNA649549)
Chandel et al. 2022 (PRJNA811248)
Acuña et al. 2023 (PRJNA926007)


E
F
G
H
I
J
Seeds (n = 1,277)
PCoA 2 (7.6%)
Seedlings (n = 602)
PCoA 2 (5.9%)
PCoA 1 (20.5%)
PCoA 1 (10.6%)
R² = 2.5%, p = 0.001
R² = 8.0%, p = 0.001
R² = 6.6%, p = 0.001
R² = 22.1%, p = 0.001
R² = 16.4%, p = 0.001
R² = 8.2%, p = 0.001
Host genus Phaseolus Raphanus Glycine Brassica Triticum Medicago Daucus Solanum Lactuca Diplotaxis Arabidopsis Petrose
Plant family Brassicaceae Fabaceae Apiaceae Asteraceae Solanaceae Poaceae