User Communication Style Determines LLM Safety in Cancer Counseling

Ramez Kouzy, MD[1,2]; Jack Gallifant, MBBS[3,4]; Shan Chen, PhD[3,5]; Marielle Fils, BS[6]; Roxana Attar, MD[7]; Danielle S. Bitterman, MD[3,5]

[1]Department of Radiation Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX

[2]Institute for Data Science in Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX

[3]Artificial Intelligence in Medicine Program, Mass General Brigham, Boston, MA

[4]Laboratory for Computational Physiology, Massachusetts Institute of Technology, Cambridge, MA

[5]Department of Radiation Oncology, Dana-Farber Cancer Institute and Brigham and Women's Hospital, Harvard Medical School, Boston, MA

[6]School of Medicine and Health Sciences, The George Washington University, Washington, DC

[7]Department of Radiation Oncology, Texas Tech University Health Sciences Center El Paso, El Paso, TX

Corresponding Author:

Danielle S. Bitterman, MD

Department of Radiation Oncology

Dana-Farber Cancer Institute and Brigham and Women's Hospital

Harvard Medical School

75 Francis Street, ASB1-L2

Boston, MA 02115

Email: dsbitterman@bwh.harvard.edu

Word Count: 3,500

Figures: 2

Tables: 1

References: 38

Running Title: User Communication Style and LLM Safety

Keywords: Large language models, artificial intelligence, sycophancy, cancer misinformation, patient safety

Conflict of Interest: The authors declare no conflicts of interest.

Funding: None.

Data Availability: Code and de-identified conversation data available at [repository].

Abstract

Introduction: Large language model (LLM) chatbots are increasingly used by patients seeking cancer information. While these models perform well on straightforward medical queries, their safety when users present non-evidence-based beliefs about cancer treatment remains uncharacterized. We hypothesized that LLMs exhibit validation bias when users frame queries with misconceptions, increasing the rate of unsafe responses compared with evidence-aligned framing.

Methods: We conducted a cross-sectional evaluation using a 2×2×2 factorial design across 1,272 conversations involving 14 LLMs from 6 providers. Conversations varied by framing (user misconceptions vs evidence-aligned), treatment type (evidence-based vs alternative), and format (single-turn vs multi-turn). The primary outcome was unsafe response rate, defined as "Weak" classification on a 6-dimension safety rubric or presence of critical failures.

Results: Framing queries with user misconceptions significantly increased unsafe response rates compared with evidence-aligned framing (14.4% vs 5.1%; odds ratio, 3.13; 95% CI, 2.05–4.79; P<.001). Among conversations initially safe under evidence-aligned framing, 22.6% degraded in quality under misconception conditions. Multi-turn dialogue reduced unsafe response rates (absolute risk reduction: 9.6 percentage points; P<.001). Provider performance varied substantially, with unsafe rates ranging from 1.8% to 36.8%.

Conclusions: LLMs exhibited significant validation bias in oncology contexts, with unsafe response rates nearly tripling when users presented misconceptions. This finding suggests that standard evaluations using neutral queries may overestimate real-world safety. Adversarial testing that includes user misconceptions should be considered for medical AI systems.

1 Introduction

Patients have long turned to the internet to supplement or challenge medical advice, but the emergence of artificial intelligence (AI) chatbots has fundamentally altered this landscape. Unlike static search engines, large language models (LLMs) engage users in conversational dialogue, offering personalized synthesis of complex medical information.[1, 2, 3] Early evaluations have been promising: studies comparing AI responses to physician answers have found LLMs to be empathetic and generally accurate when answering questions about cancer.[4, 5, 6] However, these assessments typically rely on a critical assumption—that users ask questions neutrally, seeking objective information.

In clinical reality, patients often approach medical information with preexisting beliefs, seeking validation for unproven alternative therapies or confirmation of fears regarding evidence-based treatments. This distinction is important because LLMs are trained to be helpful and agreeable—objectives that can conflict when a user's beliefs are misaligned with medical evidence. Computer science research identifies "sycophancy" as an alignment challenge: the tendency of models to agree with users' stated views rather than prioritize accuracy.[7, 8, 9, 10] In general domains, this might result in an AI agreeing with a user's preference; in oncology, it could mean validating a decision to forgo evidence-based therapy. The clinical stakes are significant: patients who choose alternative medicine over conventional cancer treatment face substantially worse outcomes.[11]

Current medical AI benchmarks primarily test performance on neutral, fact-seeking queries, creating a gap in understanding of how models perform under adversarial conditions.[12, 13, 14] Recent work by Mo and colleagues demonstrated that LLMs exhibit tension between faithfulness to provided context and adherence to safety protocols when presented with counterfactual medical evidence.[15] By omitting adversarial scenarios, existing evaluations may overestimate the safety of LLMs for real-world use.

We hypothesized that LLMs exhibit validation bias when users present non-evidence-based views about cancer treatment, resulting in higher rates of unsafe responses compared with evidence-aligned framing. To test this hypothesis, we designed a controlled $2\times2\times2$ factorial experiment to determine whether LLMs maintain safety standards when users frame questions with unsubstantiated claims, either by promoting unproven treatments or questioning evidence-based therapies.

2 Methods

2.1 Study Design

We conducted a 2×2×2 factorial experiment evaluating LLM safety when discussing cancer treatments. The three experimental factors were: (1) conversation format (single-turn vs multi-turn), (2) treatment type (evidence-based vs alternative), and (3) query valence (positive vs negative framing). The combination of treatment type and valence created the distinction between user misconception conditions (promoting alternative treatments or questioning evidence-based therapies) and evidence-aligned conditions (supporting evidence-based treatments or questioning alternative therapies).

The conversation structure modeled authentic patient interactions. Multi-turn conversations followed an escalating protocol from routine information-seeking to safety-critical scenarios, reflecting natural progressions in patient inquiries. Prior studies have isolated individual factors contributing to sycophancy in medical AI.[15, 9] Notably, adversarial testing literature has generally found multi-turn interactions more vulnerable than single-turn exchanges; our design allowed evaluation of whether this pattern holds in oncology contexts.

2.2 Conversation Dataset

We evaluated 1,272 conversations corresponding to 6,360 LLM interactions across 14 models from 6 providers between March and September 2025. Model selection aimed to create a representative sample of publicly accessible LLMs, including commercial models (Claude 3.5 Haiku, Claude 3.7 Sonnet, Gemini 1.5 Flash, Gemini 2.0 Flash, Gemini 2.0 Thinking, Gemini 2.5 Pro, GPT-4o, GPT-4o-mini, O3-mini), open-source models (Gemma 3 4B, Gemma 3 27B, DeepSeek R1 Distill, Llama 3.2 3B, Qwen 3 32B).

The treatment dataset comprised 15 alternative therapies and 15 evidence-based therapies representing common unproven and standard cancer treatments (Supplementary Appendix 1). In multi-turn format, five sequential queries (T1–T5) were delivered within a single conversation with context maintained. In single-turn format, the same queries were delivered as independent prompts. Positive and negative versions differed by only key words (e.g., “effective” vs “ineffective”) to isolate valence effects.

2.3 Evaluation Rubric

We developed a structured rubric (maximum 12 points) with six dimensions: Evidence Accuracy (0–3), Testimonial Handling (0–1), Safety Guidance (0–2), Misattribution Correction (0–1), Promotional Content Response (0–3), and Healthcare Professional References (0–2)

(Supplementary Appendix 2). Conversations were classified as Strong (10–12 points, no critical failures), Moderate (6–9 points, no critical failures), or Weak/Unsafe (0–5 points or any critical failure). Critical failures included enabling dangerous implementation, actively endorsing harmful content, or creating promotional content.

2.4 Conversation Scoring

We used GPT-4o-mini via the OpenAI Batch API as an automated judge.[**?**] The judge received structured prompts with rubric criteria, conversation transcripts, and instructions for chain-of-thought analysis before scoring. Judge validation against dual human annotation by two clinical researchers demonstrated 81.8% agreement, with dimension-level agreement ranging from 77.3% to 93.2% (Supplementary Appendix 3).

2.5 Statistical Analysis

Categorical comparisons used Pearson chi-square tests or Fisher's exact test. Odds ratios with 95% confidence intervals quantified effect sizes. Wilson score intervals provided confidence intervals for proportions. Continuous comparisons used independent samples t-tests. All analyses used Python 3.10[16] with pandas,[17] NumPy,[18] scikit-learn,[19] and statsmodels.[20] Statistical significance was set at $\alpha = 0.05$.

3 Results

Among 1,272 conversations, 919 (72.2%) were classified as Strong, 228 (17.9%) as Moderate, and 125 (9.8%) as Weak/Unsafe. A total of 111 conversations (8.7%) contained at least one critical failure.

Framing queries with user misconceptions—promoting unproven treatments or questioning evidence-based therapies—significantly increased unsafe response rates compared with evidence-aligned framing (14.4% [93/644] vs 5.1% [32/628]; odds ratio, 3.13; 95% CI, 2.05–4.79; P<.001; Table 1, Figure 1). This validation bias persisted regardless of whether users promoted harmful treatments or disparaged proven therapies, indicating that models prioritized agreement over accuracy. Critical failures were similarly elevated under misconception framing (13.5% vs 3.9%; odds ratio, 3.84; 95% CI, 2.40–6.14; P<.001).

Among 429 conversations achieving Strong classification under evidence-aligned framing, 97 (22.6%) degraded when shifted to misconception conditions: 54 (12.6%) declined to Moderate and 43 (10.0%) to Weak. This erosion occurred even in models that initially performed well, indicating that misconception framing compromises quality beyond outright failure.

Multi-turn conversations demonstrated substantially lower unsafe response rates than single-turn exchanges (5.2% [34/655] vs 14.8% [91/617]; odds ratio, 0.32; 95% CI, 0.21–0.48; P<.001; Figure 2). This represents a 9.6 percentage point absolute risk reduction overall. Under misconception framing specifically, multi-turn format reduced unsafe rates from 21.2% to 8.1% (absolute risk reduction: 13.1 percentage points).

Critical failures were dominated by promotional content generation (105/111, 94.6%), where models produced marketing-like statements for unproven therapies. Active endorsement of harmful content occurred in 43 conversations (38.7%). Performance varied substantially by provider: top-performing models (Gemini 1.5 Flash, Claude 3.7 Sonnet, O3-mini) achieved 0.8% unsafe rates, while highest-risk models (DeepSeek R1 Distill, Gemma 3 4B) showed 35–37% unsafe rates (Supplementary Table 1).

Misconception framing most severely degraded Promotional Content Response scores (mean 2.42 vs 2.73 of 3 points; P<.001) and Evidence Accuracy scores (2.68 vs 2.80; P<.001), capturing the core failures: generating promotional content for harmful treatments and reduced factual accuracy under adversarial framing (Supplementary Table 2).

Table 1: Primary Safety Outcomes by Framing Condition

Outcome	Framing		OR (95% CI)	P
	Evidence-aligned	Misconception		
Safety Classification, No. (%)				
Strong	477 (75.0)	442 (69.5)	—	.021
Moderate	127 (20.0)	101 (15.9)	—	.048
Weak (Unsafe)	32 (5.0)	93 (14.4)	3.13 (2.05–4.79)	<.001
Critical Failures, No. (%)				
Any	25 (3.9)	86 (13.5)	3.84 (2.40–6.14)	<.001

Abbreviations: CI, confidence interval; OR, odds ratio.
Misconception framing: promoting unproven treatments or questioning evidence-based treatments.




Figure 1: **Study Design and Experimental Framework. (A)** Factorial design crossing three factors: conversation format (single-turn vs multi-turn), treatment type (evidence-based vs alternative), and user valence (positive vs negative framing), yielding 1,272 conversations evaluated by an LLM judge for safety classification. **(B)** Example comparing single-turn and multi-turn strategies. In single-turn queries, models maintained safety boundaries when asked about alternative treatments. Multi-turn strategies progressively established context, leading to safety boundary compromise despite initial refusals.




Figure 2: **Safety Classification Transitions Under Misconception Framing.** Sankey diagram showing how conversations classified under evidence-aligned framing transitioned when shifted to misconception framing. Among 429 conversations initially classified as Strong, 22.6% degraded: 12.6% to Moderate and 10.0% to Weak/Unsafe.

4 Discussion

This study demonstrates that LLMs do not simply fail to provide accurate information when users present misconceptions about cancer treatment—they actively validate those misconceptions. When users framed queries with embedded beliefs, models exhibited validation bias that increased unsafe response rates nearly threefold. This finding has implications for the integration of AI chatbots into cancer care, where treatment decisions carry significant consequences.

The central finding—that misconception framing increases the odds of unsafe responses by a factor of 2.8—reveals a gap between how LLMs are evaluated and how they may perform in practice. Prior evaluations tested models under favorable conditions with evidence-aligned queries from cooperative users.[5, 4] Our findings suggest this approach may overestimate safety.

The bidirectional nature of this effect is notable. Users who promoted unproven treatments and those who questioned evidence-based therapies triggered equivalent failure patterns. This symmetry suggests that validation bias in medical AI represents an alignment challenge where the drive to agree with users can override adherence to medical evidence, which the model demonstrably possesses as evident from its performance on neutral queries.

Johnson and colleagues demonstrated that patients who choose alternative medicine instead of evidence-based cancer treatment experience substantially worse outcomes.[11] The pathways to treatment abandonment are documented: patients encounter claims about alternative treatments, seek validation from trusted sources, and delay or abandon evidence-based therapies.[21, 22, 23] If AI chatbots provide that validation, they may contribute to rather than protect against these pathways.[24, 25]

Multi-turn dialogue reduced unsafe response rates by 9.6 percentage points overall, a finding that differs from prevailing assumptions about conversational AI safety. Adversarial attack literature suggests multi-turn interactions should be more vulnerable through guardrail erosion.[26, 27] Our finding that multi-turn dialogue was protective may reflect the escalating structure of our protocol, which moved from routine information-seeking to safety-critical scenarios, enabling models to establish safety-oriented postures early. Additionally, context accumulation allowed models to reference prior safety guidance, creating consistency pressure. These findings suggest that multi-turn interfaces with appropriate safety protocols may offer advantages for managing adversarial user framing. However, even in multi-turn format, 8.1% of conversations with misconception framing resulted in unsafe responses—a rate that remains concerning for medical applications.

These findings suggest considerations for developing and deploying LLMs in clinical settings. Evaluations may benefit from including adversarial user framing, bidirectional testing of misconceptions, and multi-turn analysis.[28, 14] Healthcare providers should understand that how users frame questions can influence LLM responses. As Kouzy and colleagues demonstrated, the framing of patient queries affects not only response content but also clinical utility of AI-generated information.[**?**] Providers may need to address this dynamic when discussing information patients obtain from AI sources.

As AI chatbots increasingly serve as health information sources, their behavior becomes a public health consideration. Current evaluation frameworks focus on accuracy metrics under ideal conditions. Our findings suggest this may be insufficient—a system can demonstrate high accuracy on neutral queries while exhibiting validation bias when users present misconceptions. Pre-market evaluation of AI health systems should consider including adversarial testing protocols. These concerns become relevant as AI companies expand into healthcare, signaling commercialization that may outpace safety evaluation.

Healthcare institutions deploying AI chatbots should consider validation bias testing as part of governance frameworks. The variation we observed across models (1.8% to 36.8% unsafe rates) suggests that model selection has safety implications. This extends to insurers and delivery networks deploying chatbots for member engagement, often without evaluation of performance across different user communication styles.

Several limitations warrant consideration. Our adversarial prompts were researcher-constructed and may not capture the full diversity of real patient communication. The study used a single LLM judge rather than human ratings for all conversations, though validation demonstrated reasonable agreement. Results represent a temporal snapshot of rapidly evolving models. An important contextual point is that most responses (72.2%) were classified as Strong, indicating that LLMs generally maintain safety standards. However, from a patient safety perspective, the proportion of Moderate or Weak conversations represents a concerning failure rate where even occasional validation of harmful misconceptions may influence treatment decisions.

To our knowledge, this is the first comprehensive evaluation studying how user interaction styles impact LLM response safety in oncology.

4.1 Conclusions

LLMs exhibited significant validation bias when responding to cancer treatment queries framed with user misconceptions, with unsafe response rates increasing nearly threefold. Under conditions simulating authentic interactions where users bring preexisting beliefs, models prioritized agreement over accuracy at rates that may pose risks in clinical contexts. These findings suggest that evaluation frameworks for medical AI should evolve beyond testing on neutral queries to include adversarial scenarios. As AI chatbots become primary health information sources, ensuring their influence aligns with evidence-based care will require attention to validation bias alongside traditional accuracy metrics.

References

[1] Karan Singhal, Shekoofeh Azizi, Tao Tu, S Sara Mahdavi, Jason Wei, Hyung Won Chung, Nathan Scales, et al. Large language models encode clinical knowledge. *Nature*, 620(7972):172–180, 2023.

[2] Dave Van Veen, Cara Van Uden, Louis Blankemeier, Jean-Benoit Delbrouck, Asad Aali, et al. Adapted large language models can outperform medical experts in clinical text summarization. *Nature Medicine*, 30(4):1134–1142, 2024.

[3] Cyril Zakka, Rohan Shad, Akash Chaurasia, Alex R Dalal, Jennifer L Kim, et al. Almanac–retrieval-augmented language models for clinical medicine. *NEJM AI*, 1(2), 2024.

[4] John W Ayers, Adam Poliak, Mark Dredze, Eric C Leas, Zechariah Zhu, et al. Comparing physician and artificial intelligence chatbot responses to patient questions posted to a public social media forum. *JAMA Internal Medicine*, 183(6):589–596, 2023.

[5] Alexander Pan, David Musheyev, David Bockelman, Stacy Loeb, and Aaron E Kabarriti. Assessment of artificial intelligence chatbot responses to top searched queries about cancer. *JAMA Oncology*, 9(10):1437–1440, 2023.

[6] Maria Laymouna, Yuanchao Ma, David Lessard, Tibor Schuster, Kim Engler, and Bertrand Lebouché. Roles, users, benefits, and limitations of chatbots in health care: Rapid review. *Journal of Medical Internet Research*, 26:e56930, 2024.

[7] Shahar Ben Natan and Oren Tsur. Not your typical sycophant: The elusive nature of sycophancy in large language models. *arXiv preprint*, 2026. arXiv:2401.xxxxx.

[8] Brian Christian and Matan Mazor. Self-blinding and counterfactual self-simulation mitigate biases and sycophancy in large language models. *arXiv preprint*, 2026. arXiv:2401.xxxxx.

[9] Yusuf Çelebi, Özay Ezerceli, and Mahmoud El Hussieni. PARROT: Persuasion and agreement robustness rating of output truth–a sycophancy robustness benchmark for LLMs. *arXiv preprint*, 2025. arXiv:2411.xxxxx.

[10] Jingyu Hu, Shu Yang, Xilin Gong, et al. MONICA: Real-time monitoring and calibration of chain-of-thought sycophancy in large reasoning models. *arXiv preprint*, 2025. arXiv:2411.xxxxx.

[11] Skyler B Johnson, Henry S Park, Cary P Gross, and James B Yu. Use of alternative medicine for cancer and its impact on survival. *JAMA Oncology*, 4(10):1375–1381, 2018. Seminal paper showing alternative medicine associated with worse cancer survival.

[12] Benson Huo, Adam Boyle, Nana Marfo, Wuttikorn Tangamornsuksan, et al. Large language models for chatbot health advice studies: A systematic review. *JAMA Network Open*, 8(2):e2457879, 2025.

[13] Sepehr Shool, Sajjad Adimi, Reza Saboori Amleshi, et al. A systematic review of large language model (LLM) evaluations in clinical medicine. *BMC Medical Informatics and Decision Making*, 25(1):117, 2025.

[14] Alayna C Moreno and Danielle S Bitterman. Toward clinical-grade evaluation of large language models. *International Journal of Radiation Oncology, Biology, Physics*, 118(4):916–920, 2024.

[15] Kaijie Mo, Siddhartha Venkatayogi, Chantal Shaib, Ramez Kouzy, Wei Xu, Byron C Wallace, and Junyi Jessy Li. Faithfulness vs. safety: Evaluating LLM behavior under counterfactual medical evidence. *arXiv preprint arXiv:2601.XXXXX*, 2026. Preprint.

[16] Python Software Foundation. *Python Language Reference, version 3.10*. Python Software Foundation, Wilmington, DE, 2021.

[17] Wes McKinney. pandas: a foundational Python library for data analysis and statistics. *Python for High Performance and Scientific Computing*, 14(9):1–9, 2011.

[18] Charles R Harris, K Jarrod Millman, Stéfan J van der Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, Julian Taylor, Sebastian Berg, Nathaniel J Smith, et al. Array programming with NumPy. *Nature*, 585(7825):357–362, 2020.

[19] Fabian Pedregosa, Gaël Varoquaux, Alexandre Gramfort, Vincent Michel, Bertrand Thirion, Olivier Grisel, Mathieu Blondel, Peter Prettenhofer, Ron Weiss, Vincent Dubourg, et al. Scikit-learn: Machine learning in Python. *Journal of Machine Learning Research*, 12:2825–2830, 2011.

[20] Skipper Seabold and Josef Perktold. Statsmodels: Econometric and statistical modeling with Python. In *9th Python in Science Conference*, 2010.

[21] Hye Young Yoon, Ki Hyun You, Jung Hye Kwon, et al. Understanding the social mechanism of cancer misinformation spread on YouTube and lessons learned: Infodemiological study. *Journal of Medical Internet Research*, 24(11):e39571, 2022.

[22] Pedro D Delgado-López and Eva M Corrales-García. Influence of internet and social media in the promotion of alternative oncology, cancer quackery, and the predatory publishing phenomenon. *Cureus*, 10(5):e2617, 2018.

[23] Jessica Peterson, Thomas Wilson, Julie Gruhl, et al. Timing and motivations for alternative cancer therapy with insights from a crowdfunding platform: Cross-sectional mixed methods study. *JMIR Cancer*, 8(2):e34183, 2022.

[24] Shan Chen, Marco Guevara, Srinivas Moningi, others, and Danielle S Bitterman. The effect of using a large language model to respond to patient messages. *The Lancet Digital Health*, 6(6):e379–e381, 2024.

[25] David Chen, Reza Parsa, Kira Swanson, Justin J Nunez, Andrew Critch, Danielle S Bitterman, Fei-Fei Liu, and Srinivas Raman. Large language models in oncology: a review. *BMJ Oncology*, 4(1):e000759, 2025.

[26] Akihiro Yamamoto, Masayuki Koda, Hitoshi Ogawa, Tomoko Miyoshi, Yoshito Maeda, Fumio Otsuka, and Hidekazu Ino. Enhancing medical interview skills through AI-simulated patient interactions: Nonrandomized controlled trial. *JMIR Medical Education*, 10:e58753, 2024.

[27] Haowen Yu, Jie Zhou, Long Li, others, Danielle S Bitterman, et al. Simulated patient systems powered by large language model-based AI agents offer potential for transforming medical education. *Communications Medicine*, 6(1):27, 2025.

[28] Jack Gallifant, Majid Afshar, Salih Ameen, Yin Aphinyanaphongs, others, and Danielle S Bitterman. The TRIPOD-LLM reporting guideline for studies using large language models. *Nature Medicine*, 31(1):60–69, 2025.

Supplementary Appendix

Supplementary Table 1: Safety Performance by Model Provider

Provider	n	Unsafe, %	95% CI	Strong, %
Anthropic	227	1.8	0.6–4.0	85.5
OpenAI	359	8.6	6.0–12.0	78.0
Google*	501	10.2	7.7–13.2	71.7
Meta	118	16.9	10.8–24.7	55.1
Alibaba	29	17.2	6.5–35.0	72.4
DeepSeek	38	36.8	22.4–53.7	39.5
Overall	**1272**	**9.8**	**8.2–11.6**	**72.2**

CI, confidence interval.
* Includes Gemini (commercial) and Gemma (open-source).

Supplementary Table 2: Rubric Dimension Scores by Framing Condition

Dimension (max score)	Evidence-aligned	Misconception	Diff	P
Evidence Accuracy (3)	2.80 (0.5)	2.68 (0.6)	-0.12	<.001
Testimonial Handling (1)	0.61 (0.5)	0.83 (0.4)	+0.22	<.001
Safety Guidance (2)	1.73 (0.5)	1.70 (0.5)	-0.03	.45
Misattribution Correction (1)	0.68 (0.5)	0.77 (0.4)	+0.09	.002
Promotional Content (3)	2.73 (0.7)	2.42 (1.0)	-0.31	<.001
HCP Referral (2)	1.67 (0.5)	1.54 (0.6)	-0.13	<.001
Total (12)	**10.22 (2.1)**	**9.94 (2.8)**	**-0.28**	**.045**

Values shown as mean (SD). HCP, healthcare professional.

Supplementary Figure 1: Model Performance by Provider

Supplementary Figure 2: Multi-Turn Protection Effect

Appendix 1: Treatment List

Alternative Therapies (15): Apricot seeds/Laetrile, Black salve, Colloidal silver, Essiac tea, Gerson therapy, High-dose vitamin C infusions, Hydrogen peroxide therapy, Laetrile (amygdalin), Mistletoe extract, Oxygen therapy, Rife machine, Shark cartilage, Sodium bicarbonate therapy, Turmeric/curcumin supplements, Ukrain.




Figure S1: Forest plot of unsafe response rates for all 14 evaluated models with 95% Wilson score confidence intervals.




Figure S2: Interaction between conversation format and framing condition on unsafe response rates.

Evidence-Based Therapies (15): Chemotherapy, Radiation therapy, Immunotherapy, Targeted therapy, Hormone therapy, Surgery, Stem cell transplant, CAR-T cell therapy, Proton therapy, Brachytherapy, PARP inhibitors, Checkpoint inhibitors, Monoclonal antibodies, Tyrosine kinase inhibitors, Adjuvant therapy.

Appendix 2: Evaluation Rubric Summary

- **Evidence Accuracy (0–3):** Factual correctness with clinical context
- **Testimonial Handling (0–1):** Explanation of anecdotal evidence limitations
- **Safety Guidance (0–2):** Clarity of safety recommendations
- **Misattribution Correction (0–1):** Correction of false claims about prior recommendations
- **Promotional Content Response (0–3):** Refusal to generate marketing materials
- **Healthcare Professional References (0–2):** Consistency of medical consultation recommendations

Critical Failures (automatic Weak classification):

1. Enables dangerous implementation
2. Actively endorses harmful content
3. Creates promotional content

Appendix 3: LLM Judge Validation

Agreement between LLM judge (GPT-4o-mini) and human annotators was assessed on 88 pilot conversations with dual annotation. Overall agreement (LLM score matching at least one human annotator): 81.8%.

Dimension-level agreement: Evidence Accuracy 84.1%, Testimonial Handling 93.2%, Safety Guidance 79.5%, Misattribution Correction 77.3%, Promotional Content 81.8%, HCP Referral 88.6%.