

# **MAT 335 Notes**

Max Chien

Fall 2024

# Contents

<b>1</b>	<b>Preliminaries</b>	<b>3</b>
1.1	The Complex Number System . . . . .	3
1.2	Topology of $\mathbb{C}$ . . . . .	7
1.3	Functions on $\mathbb{C}$ . . . . .	12
1.4	Power Series . . . . .	20
1.5	Integration on Curves . . . . .	26
<b>2</b>	<b>Cauchy's Theorem and Applications</b>	<b>33</b>
2.1	Goursat's Theorem . . . . .	33
2.2	Homotopies and Simply Connected Domains . . . . .	38
2.3	Consequences of Cauchy's Theorem . . . . .	42
	<b>Definitions</b>	<b>51</b>

## Introduction

This document contains notes taken for the class MAT 335: Complex Analysis at Princeton University, taken in the Fall 2024 semester. These notes are primarily based on lectures by Professor Assaf Naor. Other references used in these notes include *Complex Analysis* by Elias Stein and Rami Shakarchi, *Complex Analysis* by Lars Ahlfors, *Visual Complex Analysis* by Tristan Needham, and *Real and Complex Analysis* by Walter Rudin. Since these notes were primarily taken live, they may contain typos or errors.

# Chapter 1

## Preliminaries

### 1.1 The Complex Number System

The set of complex numbers, denoted  $\mathbb{C}$  is identified with ordered pairs  $(x, y) \in \mathbb{R}^2$ . We may alternately write this as  $x + iy$ , where the symbol  $i$  is currently undefined.

For a given complex number  $z = x + iy$ ,  $x = \operatorname{Re}(z)$  is called the **real part** of  $z$ ,  $y = \operatorname{Im}(z)$  is called the **imaginary part**,  $|z| = \sqrt{x^2 + y^2}$  is the **modulus** of  $z$ , and the **argument** of  $z$ ,  $\theta = \arg(z)$ , is the angle between  $(x, y)$  and the  $x$ -axis, defined up to integer multiples of  $2\pi$ .

#### Definition 1.1

Let  $\theta \in \mathbb{R}$ . We define

$$e^{i\theta} = \cos \theta + i \sin \theta = (\cos \theta, \sin \theta)$$

One can observe using the identity  $\cos^2 + \sin^2 = 1$  that  $e^{i\theta}$  lies on the unit circle. Moreover, if  $r = |z|$ , then elementary geometry shows that we have  $z = re^{i\theta}$  using the definition above.

#### Proposition 1.1

For any  $z \in \mathbb{C}$ ,  $|\operatorname{Re}(z)| \leq |z|$  and  $|\operatorname{Im}(z)| \leq |z|$ .

*Proof.*  $|z| = \sqrt{\operatorname{Re}(z)^2 + \operatorname{Im}(z)^2}$ . □

One of the distinguishing features of  $\mathbb{C}$  from the real plane  $\mathbb{R}^2$  is the algebraic structure present on  $\mathbb{C}$ .

**Definition 1.2**

Let  $z_1 = x_1 + iy_1$  and  $z_2 = x_2 + iy_2$ . Then we define addition and multiplication on  $\mathbb{C}$  by

$$\begin{aligned} z_1 + z_2 &= (x_1 + x_2) + i(y_1 + y_2) \\ z_1 z_2 &= (x_1 x_2 - y_1 y_2) + i(x_1 y_2 + y_1 x_2) \end{aligned}$$

Taking  $i = (0, 1)$ , then we observe that  $i^2 = -1 + 0i = -1$ . Thus we recover the basic identity  $i^2 = -1$ . We also observe that  $\text{Re}$  and  $\text{Im}$  are both linear operators.

**Proposition 1.2**

Addition and multiplication over  $\mathbb{C}$  are commutative and associative. Moreover, multiplication distributes over addition.

*Proof.* Commutative and associativity of addition is inherited from  $\mathbb{R}$ .  $\square$

Using the definition of  $e^{i\theta}$ , we can reinterpret complex multiplication in a much more pleasant manner than the definition above.

**Proposition 1.3**

If  $z_1 = r_1 e^{i\theta_1}$  and  $z_2 = r_2 e^{i\theta_2}$ , then

$$z_1 z_2 = (r_1 r_2) e^{i(\theta_1 + \theta_2)}$$

*Proof.* We have proved commutativity. From here, we apply trig identities.  $\square$

Thus multiplication results in multiplication of lengths and addition of arguments.

**Proposition 1.4**

For  $z_1, z_2 \in \mathbb{C}$ , the **triangle inequality** holds:

$$|z_1 + z_2| \leq |z_1| + |z_2|$$

*Proof.* Choose  $r, \theta$  such that  $z_1 + z_2 = r e^{i\theta}$ . Then

$$|z_1 + z_2| = r = (z_1 + z_2) e^{-i\theta} = z_1 e^{-i\theta} + z_2 e^{-i\theta} = \text{Re}(z_1 e^{-i\theta} + z_2 e^{-i\theta})$$

Now note that  $\text{Re}(z + w) = \text{Re}(z) + \text{Re}(w)$ . So

$$\text{Re}(z_1 e^{-i\theta} + z_2 e^{-i\theta}) = \text{Re}(z_1 e^{-i\theta}) + \text{Re}(z_2 e^{-i\theta}) \leq |z_1 e^{-i\theta}| + |z_2 e^{-i\theta}| = |z_1| + |z_2| \quad \square$$

The above proof amounts to applying the real triangle inequality to the components of  $z_1, z_2$  in the direction of  $z_1 + z_2$ .

### Corollary 1.5

The reverse triangle inequality also holds:

$$||z| - |w|| \leq |z - w|$$

*Proof.* We have

$$\begin{cases} |z| \leq |z - w| + |w| \\ |w| \leq |w - z| + |z| \end{cases} \implies \begin{cases} |z| - |w| \leq |z - w| \\ -|z - w| \leq |z| - |w| \end{cases} \implies ||z| - |w|| \leq |z - w| \quad \square$$

### Definition 1.3

Let  $z = x + iy \in \mathbb{C}$ . Then the **complex conjugate** of  $z$  is defined as

$$\bar{z} = x - iy$$

Geometrically, this is reflection over the  $x$  axis.

### Proposition 1.6

For  $z \in \mathbb{C}$ ,  $z\bar{z} = |z|^2$ .

*Proof.* Let  $z = x + iy$ . Then

$$z\bar{z} = (x + iy)(x - iy) = x^2 + y^2 = |z|^2 \quad \square$$

### Definition 1.4

For  $z \neq 0$ , define

$$\frac{1}{z} = \frac{\bar{z}}{|z|^2}$$

The above proposition and definition show that

$$z \cdot \frac{1}{z} = 1$$

### Definition 1.5

A **sequence** of complex numbers  $\{z_n\}_{n=1}^{\infty}$  **converges** to  $z \in \mathbb{C}$  (written  $\lim_{n \rightarrow \infty} z_n = z$ ) if

$$\begin{cases} \lim_{n \rightarrow \infty} \operatorname{Re}(z_n) = \operatorname{Re}(z) \\ \lim_{n \rightarrow \infty} \operatorname{Im}(z_n) = \operatorname{Im}(z) \end{cases}$$

This equivalent to the familiar definition:

**Proposition 1.7**

A sequence  $\{z_n\} \subseteq \mathbb{C}$  converges to  $z$  if and only for  $\varepsilon > 0$  there exists  $N \in \mathbb{N}$  such that whenever  $n \geq N$  we have

$$|z_n - z| < \varepsilon$$

*Proof.* ( $\implies$ ) Let  $\varepsilon > 0$ . Then pick  $N_1, N_2$  such that

$$\begin{cases} n \geq N_1 \implies |\operatorname{Re}(z_n) - \operatorname{Re}(z)| < \frac{\varepsilon}{\sqrt{2}} \\ n \geq N_2 \implies |\operatorname{Im}(z_n) - \operatorname{Im}(z)| < \frac{\varepsilon}{\sqrt{2}} \end{cases}$$

Letting  $N = \max\{N_1, N_2\}$ , whenever  $n \geq N$  we have

$$|z_n - z|^2 = \operatorname{Re}(z_n - z)^2 + \operatorname{Im}(z_n - z)^2 = |\operatorname{Re}(z_n) - \operatorname{Re}(z)|^2 + |\operatorname{Im}(z_n) - \operatorname{Im}(z)|^2 < \frac{\varepsilon^2}{2} + \frac{\varepsilon^2}{2}$$

Taking square roots on both sides we have

$$|z_n - z| < \varepsilon$$

$$(\impliedby) |\operatorname{Re}(z_n) - \operatorname{Re}(z)| = |\operatorname{Re}(z_n - z)| \leq |z_n - z|$$

□

We similarly define the limit of a complex function  $\lim_{z \rightarrow a} f(z)$ .

**Definition 1.6**

A **Cauchy sequence** is a sequence  $(z_n) \subseteq \mathbb{C}$  such that  $(\operatorname{Re}(z_n))$  and  $(\operatorname{Im}(z_n))$  are both Cauchy.

Again we can formulate this analogously to the single variable case:

**Proposition 1.8**

A sequence  $\{z_n\} \subseteq \mathbb{C}$  is Cauchy if and only if for all  $\varepsilon > 0$  there exists  $N \in \mathbb{N}$  such that whenever  $m, n \geq N$  it follows that

$$|z_n - z_m| < \varepsilon$$

*Proof.* Same as the proof of Proposition 1.7. □

**Proposition 1.9**

A Cauchy sequence is convergent.

*Proof.* Follows from completeness of  $\mathbb{R}$ :

$$\{z_n\} \text{ conv.} \iff \begin{cases} \{\operatorname{Re}(z_n)\} \text{ conv.} \\ \{\operatorname{Im}(z_n)\} \text{ conv.} \end{cases} \iff \begin{cases} \{\operatorname{Re}(z_n)\} \text{ Cauchy} \\ \{\operatorname{Im}(z_n)\} \text{ Cauchy} \end{cases} \iff \{z_n\} \text{ Cauchy} \quad \square$$

## 1.2 Topology of $\mathbb{C}$

The topological nature of  $\mathbb{C}$  should not be unfamiliar to the reader, since it is essentially the same as that of  $\mathbb{R}^2$ , rephrased slightly using complex variables.

### Definition 1.7

Let  $r > 0$  and  $z_0 \in \mathbb{C}$ . Then the **open disk** of radius  $r$  about  $z_0$  is the set

$$\mathbb{D}_r(z_0) = \{z \in \mathbb{C} : |z - z_0| < r\}$$

and the **closed disk** as

$$\overline{\mathbb{D}}_r(z_0) = \{z \in \mathbb{C} : |z - z_0| \leq r\}$$

We also specify  $\mathbb{D}_r = \mathbb{D}_r(0)$  and  $\mathbb{D} = \mathbb{D}_1$ .

### Definition 1.8

An **interior point**  $z_0 \in \Omega$  of a subset  $\Omega \subseteq \mathbb{C}$  is a point such that there exists  $r > 0$  where  $\mathbb{D}_r(z_0) \subseteq \Omega$ .

### Definition 1.9

The set of interior point in  $\Omega$  is the **interior** of  $\Omega$ , denoted  $\text{int } \Omega$ .

### Definition 1.10

An **open set** in  $\mathbb{C}$  is a subset  $\Omega \subseteq \mathbb{C}$  such that for any  $z_0 \in \Omega$  there exists  $\varepsilon > 0$  such that  $\mathbb{D}_\varepsilon(z_0) \subseteq \Omega$ .

It is immediate that  $\Omega$  is open if and only if  $\text{int } \Omega = \Omega$ .

### Definition 1.11

Let  $\Omega \subseteq \mathbb{C}$  and let  $z \in \mathbb{C}$ .  $z$  is a **limit point** of  $\Omega$  if there exists a sequence of points  $\{z_n\}_{n=1}^\infty \subseteq \Omega$  such that  $z_n \neq z$  for each  $n$  and  $\lim z_n = z$ .

We can equivalently define a limit point as a point  $z$  such that  $\mathbb{D}_r(z) \setminus \{z\} \cap \Omega \neq \emptyset$  for each  $r > 0$

### Definition 1.12

$A \subseteq \mathbb{C}$  is a **closed set** if  $\mathbb{C} \setminus A$  is open.



**Proposition 1.10**

$A$  is closed if and only if it contains all its limit points.

*Proof.* ( $\implies$ ) Suppose not. Then pick  $z$  which is a limit point of  $A$  that is not in  $A$ . Then there is no disk around  $z$  entirely contained in  $\mathbb{C} \setminus A$ . Thus  $A$  is not closed.

( $\impliedby$ ) Suppose  $A$  is not closed. Then there exists  $z \notin A$  such that each  $\mathbb{D}_r(z) \setminus \{z\}$  intersects  $A$ . Then  $z$  is a limit point of  $A$ .  $\square$

**Definition 1.13**

The **closure** of  $\Omega \subseteq \mathbb{C}$ , denoted  $\overline{\Omega}$ , is the union of  $\Omega$  with its limit points.

**Definition 1.14**

The **boundary** of  $\Omega \subseteq \mathbb{C}$ , denoted  $\partial\Omega$ , is defined as  $\overline{\Omega} \setminus \text{int } \Omega$ .

**Definition 1.15**

$\Omega \subseteq \mathbb{C}$  is **bounded** if there exists  $M > 0$  such that  $|z| < M$  for each  $z \in \Omega$  (or equivalently,  $\Omega \subseteq \mathbb{D}_M$ ).

**Definition 1.16**

Let  $\Omega \subseteq \mathbb{C}$  be bounded. Then the **diameter** of  $\Omega$  is defined as

$$\text{diam } \Omega = \sup_{z, w \in \Omega} |z - w|$$

The following definition, as in the real case, is critical:

**Definition 1.17**

$\Omega \subseteq \mathbb{C}$  is **compact** if it is closed and bounded.

**Theorem 1.11: Bolzano-Weierstrass Theorem**

Let  $\Omega \subseteq \mathbb{C}$ . Then the following conditions are equivalent:

1.  $\Omega$  is compact.
2. Each sequence  $\{z_n\}_{n=1}^{\infty} \subseteq \Omega$  has a subsequence  $\{z_{n_k}\}_{k=1}^{\infty}$  which converges to some  $z \in \Omega$ .

We can treat  $\mathbb{C}$  similarly to  $\mathbb{R}^2$  to prove this.

*Proof.* (1  $\implies$  2) If  $\Omega$  is compact, then  $\{z_n\}_{n=1}^\infty \subseteq \mathbb{C}$  may be written as  $\{x_n + iy_n\}_{n=1}^\infty \subseteq \mathbb{C}$ . Since  $\Omega$  is bounded, there exists  $M > 0$  such that  $|z| < M$  for all  $z \in \Omega$ . In particular  $\sqrt{x_n^2 + y_n^2} = |z_n| < M$ . So the real sequences  $\{x_n\}_{n=1}^\infty, \{y_n\}_{n=1}^\infty$  are bounded. Apply the real version of Bolzano-Weierstrass, there exists a convergent subsequence  $\{x_{n_k}\}$ . Then consider the sequence  $\{y_{n_k}\}$ . This is also bounded, so we apply Bolzano-Weierstrass again to produce  $\{y_{n_{k_i}}\}$  convergent. Then the sequence  $\{z_{n_{k_i}}\}$  is a convergent subsequence. If  $z = z_n$  for some  $n$ , then  $z \in \Omega$ ; otherwise it is a limit point. Since  $\Omega$  is closed it contains its limit points so  $z \in \Omega$ .

(2  $\implies$  1) Suppose each sequence has a convergent subsequence. Let  $z$  be a limit point and let  $\{z_n\} \subseteq \Omega \setminus \{z\}$  be a sequence converging to  $z$ . Then there exists a subsequence  $\{z_{n_k}\}$  which converges to  $z' \in \Omega$ . But subsequences converge to the same value as the original sequence, so  $z = z' \in \Omega$ . So  $\Omega$  is closed. If  $\Omega$  is not bounded, then we may take  $\{z_n\}$  such that  $|z_n| \geq n$ , and that  $|z_{n+1}| > |z_n| + 1$ . But then  $|z_m - z_{m-1}| > 1$  so no subsequence is Cauchy and thus no subsequence converges. So  $\Omega$  is bounded.  $\square$

### Definition 1.18

An **open cover** of a set  $\Omega \subseteq \mathbb{C}$  is a collection  $\mathcal{O}$  of open sets such that each  $z \in \Omega$  is contained in some  $O \in \mathcal{O}$ . A **subcover** of  $\mathcal{O}$  is a subcollection which is still a cover.

### Theorem 1.12: Heine-Borel Theorem

A set  $\Omega \subseteq \mathbb{C}$  is compact if and only if every open cover has a finite subcover.

*Proof.* ( $\implies$ ) Since  $\Omega$  is bounded, it is a subset of a closed rectangle  $K$ . We showed in  $\mathbb{R}^2$  that  $X \times Y$  is compact when  $X, Y \subseteq \mathbb{R}$  are, and the same is true here. So  $K$  is compact. Take an open cover  $\mathcal{O}$  of  $\Omega$  and add the (open) set  $\mathbb{C} \setminus \Omega$ . This is an open cover of  $\mathbb{C}$  and thus one of  $K$ , so only finitely many are needed. Remove  $\mathbb{C} \setminus \Omega$  if necessary and we still have an open cover of  $\Omega$ .

( $\impliedby$ ) Boundedness is immediate by covering  $\Omega$  with balls of finite radius.

For closure, suppose not. Then take a limit point  $w \notin \Omega$ . Each  $z \in \Omega$  has  $|z - w| > 0$ , so we may cover  $\Omega$  with open balls  $O_z = \mathbb{D}_\varepsilon(z)$  where  $\varepsilon < |z - w|/2$ . Then a finite number of them cover  $\Omega$  but this implies that  $w$  is not a limit point.  $\square$

For the sake of completeness, here is an independent proof that a set is sequentially compact if it is covering compact.

*Proof that covering compactness  $\implies$  sequential compactness.* Let  $K$  be covering compact and pick a sequences  $\{a_n\} \subseteq K$ . Suppose for contradiction that  $a_n$  has no convergent subsequence in  $K$ . Then for each  $x \in K$ , there exists  $\varepsilon_x > 0$  and  $N_x \in \mathbb{N}$  such that whenever  $n \geq N_x$  it follows that  $a_n \notin \mathbb{D}_{\varepsilon_x}(x)$ . Then the collection of  $\mathbb{D}_{\varepsilon_x}(x)$  for  $x \in K$  is an open cover of  $K$ , so we may pick a finite subcover

$$\mathbb{D}_{\varepsilon_{x_1}}(x_1), \dots, \mathbb{D}_{\varepsilon_{x_m}}(x_m)$$

Then let  $N = \max N_{x_i}$ . For  $n \geq N$  it follows that  $a_n \notin K$ , contradiction.  $\square$

**Proposition 1.13: Nested Compact Set Property**

Suppose that  $\Omega_1 \supseteq \Omega_2 \supseteq \dots$  is a nested sequence of compact, nonempty subsets of  $\mathbb{C}$ . Then

$$\bigcap_{n=1}^{\infty} \Omega_n \neq \emptyset$$

Moreover, if  $\lim_{n \rightarrow \infty} \text{diam } \Omega_n = 0$ , then there is a unique point  $z \in \mathbb{C}$  such that  $z \in \Omega_n$  for all  $n$ .

*Proof.* Choose  $z_n \in \Omega_n$  for each  $n$ . Then the sequence of points  $\{z_n\} \subseteq \Omega_1$ , and  $\Omega_1$  is compact, so there exists a convergent subsequence  $\{z_{n_k}\}$  tending to  $z \in \Omega_1$ . Then for arbitrary  $\Omega_n$ , there exists a subsequence  $\{z_{n_k+k_0}\} \subseteq \Omega_n$  for sufficiently large  $k_0$ , which converges to  $z$  and we see that  $z \in \Omega_n$ . So the intersection is nonempty.

To show uniqueness, take  $z, w \in \bigcap_{n=1}^{\infty} \Omega_n$ . Then

$$|z - w| \leq \text{diam } \Omega_n$$

for each  $n$ , but  $\text{diam } \Omega_n \rightarrow 0$  so  $|z - w| = 0$  and thus  $z = w$ .  $\square$

**Remark**

With the assumption that  $\text{diam } \Omega_n \rightarrow 0$ , we need not take subsequences as  $\{z_n\}$  itself is Cauchy. To see this, pick  $\varepsilon > 0$  and let  $N$  be such that  $\text{diam } \Omega_n < \varepsilon$  for any  $n \geq N$ . Then for any  $n, m \geq N$ ,  $z_n, z_m \in \Omega_N$  and thus  $|z_n - z_m| \leq \text{diam } \Omega_N < \varepsilon$ .

**Definition 1.19**

A set  $\Omega \subseteq \mathbb{C}$  is **connected** if there are no nonempty disjoint sets  $A, B \subseteq \Omega$  such that  $\Omega = A \sqcup B$  such that  $A \cap \overline{B} = \emptyset$  and  $\overline{A} \cap B = \emptyset$ .

The above definition may be rephrased as saying that  $\Omega$  is not the disjoint union of nonempty sets which are open in the subspace topology of  $\Omega$ :

**Proposition 1.14**

A topological space  $X$  is connected if and only if it cannot be written as  $X = \Omega_1 \cup \Omega_2$  with  $\Omega_1, \Omega_2$  nonempty, disjoint and open (in  $X$ ).

*Proof.* ( $\implies$ ) Suppose  $X \subseteq \mathbb{C}$  is connected. Let  $\Omega_1, \Omega_2 \subseteq X$  be nonempty, open and disjoint. Consider  $\Omega_2$ . Then  $\Omega_2 \subseteq X \setminus \Omega_1$ . By definition  $X \setminus \Omega_1$  is closed.  $\overline{\Omega_2}$  is the smallest closed set containing  $\Omega_2$ , so  $\overline{\Omega_2} \subseteq X \setminus \Omega_1$  and thus  $\Omega_1 \cap \overline{\Omega_2} = \emptyset$ . Similarly  $\overline{\Omega_1} \cap \Omega_2 = \emptyset$ . Since  $X$  is connected, we conclude that  $\Omega_1 \cup \Omega_2 \neq X$ .

( $\impliedby$ ) Pick  $A, B$  nonempty with  $X = A \cup B$ . Assume that  $A \cap \overline{B} = B \cap \overline{A} = \emptyset$ , so that  $A \subseteq X \setminus \overline{B}$  and  $B \subseteq X \setminus \overline{A}$ . Define  $\Omega_1 = X \setminus \overline{B}$  and  $\Omega_2 = X \setminus \overline{A}$ . Since  $X = A \cup B$ , we have  $X = \Omega_1 \cup \Omega_2$ .  $\Omega_1, \Omega_2$  are both open in  $X$ , so it must not be the case that they are disjoint. So there exists some  $x \in \Omega_1 \cap \Omega_2$ . But this implies that  $x \notin A$  and  $x \notin B$ .  $\square$

Thus the above general definition can be simplified for nicer sets:

**Proposition 1.15**

If  $\Omega$  is open, then it is connected if and only if it cannot be written as the union of disjoint open sets (in  $\mathbb{C}$ ). Similarly if  $F$  is closed then it is connected if and only if it is not the union of disjoint closed sets.

We may also introduce another notion of connectedness, which involves functions into  $\Omega$ .

**Definition 1.20**

Suppose  $\Omega \subseteq \mathbb{C}$  and  $f : \Omega \rightarrow \mathbb{C}$ .  $f$  is **continuous** at  $z_0$  if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that whenever  $z \in \Omega$  and  $|z - z_0| < \delta$ , it follows that  $|f(z) - f(z_0)| < \varepsilon$ .

**Proposition 1.16**

$f$  is continuous at  $z_0$  if and only if for every  $\{z_n\} \subseteq \Omega$  with  $z_n \rightarrow z_0$ , it follows that  $f(z_n) \rightarrow f(z_0)$ . We say that  $f$  is continuous on  $\Omega$  if it is continuous at each point in  $\Omega$ .

**Definition 1.21**

A **path** is a function  $f : [0, 1] \rightarrow \mathbb{C}$ . A continuous path is a continuous such function.

**Definition 1.22**

A set  $\Omega \subseteq \mathbb{C}$  is **path connected** if for any  $z, w \in \Omega$  there exists a continuous path with  $f(0) = z$  and  $f(1) = w$  with  $f(t) \in \Omega$  for each  $t \in [0, 1]$ .

**Proposition 1.17**

An open set  $\Omega$  is path connected if and only if it is connected.

**Definition 1.23**

A nonempty open, connected set  $\Omega \subseteq \mathbb{C}$  is called a **region**.

**Corollary 1.18**

A region is path connected.

### 1.3 Functions on $\mathbb{C}$

We now turn our attention to functions which map complex numbers to complex numbers, the primary object of study in this course. Continuing from the definition of continuity from the previous section, we have the following:

#### Proposition 1.19

If  $f$  is continuous at  $z_0$  then  $|f|$  is continuous at  $z_0$ .

*Proof.* By the reverse triangle inequality we have  $||f(z)| - |f(z_0)|| \leq |f(z) - f(z_0)|$ . The conclusion follows.  $\square$

#### Definition 1.24

$f$  attains its maximum on  $\Omega \subseteq \mathbb{C}$  if there exists  $z_0 \in \Omega$  such that

$$|f(z)| \leq |f(z_0)|$$

for each  $z \in \Omega$ . The minimum case is analogous.

#### Theorem 1.20

Suppose that  $\Omega \subseteq \mathbb{C}$  is compact and  $f : \Omega \rightarrow \mathbb{C}$  is continuous. Then  $f$  attains its maximum (and minimum) on  $\Omega$ .

*Proof.* First we show that  $f$  is bounded on  $\Omega$ . If not, then we may take a sequence of points  $\{z_n\} \subseteq \Omega$  such that  $|f(z_n)| \rightarrow \infty$ . Then  $\{z_n\}$  contains a convergent subsequence  $\{z_{n_k}\}$  tending to some  $z \in \Omega$ . It follows that

$$|f(z_{n_k})| \rightarrow |f(z)|$$

But the left side diverges to  $\infty$ , contradiction. Thus  $f(\Omega)$  is bounded.

Let  $M = \sup |f|(\Omega)$ . Then there exists a sequence  $\{z_n\} \subseteq \Omega$  such that  $|f(z_n)| \rightarrow M$ . Then there exists a subsequence  $\{z_{n_k}\}$  converging to  $z \in \Omega$ . By continuity we have

$$|f(z)| = \lim |f(z_{n_k})| = M \quad \square$$

We now make the most important definition of this course:

#### Definition 1.25

Let  $\Omega \subseteq \mathbb{C}$  be open and let  $z_0 \in \Omega$ . Let  $f : \Omega \rightarrow \mathbb{C}$ . We say that  $f$  is **holomorphic** at  $z_0$  (or complex differentiable) if the limit

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}$$

exists. In this case, the limit is denoted  $f'(z_0)$ .

If  $f$  is holomorphic at every  $z \in \Omega$ , then we simply say it is holomorphic on  $\Omega$ . If  $f$  is holomorphic on  $\mathbb{C}$  it is said to be **entire**.

We will sometimes also say that  $f$  is **analytic** or **complex differentiable** when it is holomorphic.

Note that the specification that  $\Omega$  is open ensures that the difference quotient is actually defined (for sufficiently small  $h$ ). Moreover, although this definition appears similar to the real analogue, the structure of the complex numbers means that it has far-reaching implications.

We will prove the following theorems in this class:

- (Cauchy's Theorem) If  $f$  is holomorphic on  $\Omega$ , then it has derivatives of all orders.
- (Liouville's Theorem) If  $f$  is entire and bounded, then it is constant.
- (Prime Number Theorem) If  $\pi(n)$  denotes the number of prime numbers less than or equal to  $n$ , then

$$\lim_{n \rightarrow \infty} \pi(n) \cdot \frac{\ln n}{n} = 1$$

- (Hardy-Ramanujan Theorem) Define  $p(n)$  (the partition function) to be the number of ways to write  $n = k_1 + k_2 + \dots + k_n$  where  $k_1 \geq k_2 \geq \dots \geq k_n$  are all integers. For instance,  $p(4) = 5$ . Then

$$p(n) \sim \frac{1}{n\sqrt{48}} e^{\pi\sqrt{\frac{2}{3}}\sqrt{n}}$$

#### Example 1.1

The function  $f(z) = z$  is holomorphic:

$$\frac{f(z+h) - f(z)}{h} = \frac{z+h-z}{h} = \frac{h}{h} = 1$$

so  $z' = 1$ .

#### Definition 1.26

If  $A \subseteq \mathbb{C}$  is closed and  $f : A \rightarrow \mathbb{C}$ , then we say  $f$  is holomorphic on  $A$  if there exists  $\Omega \supseteq A$  open and  $F : \Omega \rightarrow \mathbb{C}$  which is holomorphic, and  $F|_A = f$ .

We can rewrite the definition of holomorphicity similarly to the multivariable real case as the following:

**Proposition 1.21**

$f : \Omega \rightarrow \mathbb{C}$  ( $\Omega$  open) is holomorphic at  $z_0$  if and only if there exists  $a \in \mathbb{C}$  and  $\psi : \mathbb{C} \rightarrow \mathbb{C}$  with  $\psi(h) \rightarrow 0$  as  $h \rightarrow 0$  such that

$$f(z_0 + h) = f(z_0) + ah + h\psi(h)$$

on some  $\mathbb{D}_r(z_0) \subseteq \Omega$ .

*Proof.* We can rewrite the above as

$$\psi(h) = \frac{f(z_0 + h) - f(z_0)}{h} - a$$

which goes to 0 if and only if

$$\frac{f(z_0 + h) - f(z_0)}{h} \rightarrow a$$

so that  $f'(z_0) = a$ . □

This recharacterization allows for a simple proof of the following:

**Proposition 1.22**

If  $f$  is holomorphic at  $z_0$ , then it is continuous at  $z_0$ .

*Proof.* Let  $\{z_n\}$  be a sequence with  $z_n \rightarrow z_0$ . We want to show that  $f(z_n) \rightarrow f(z_0)$ . Let  $h_n = z_n - z_0$ . Then

$$f(z_n) = f(z_0 + h_n) = f(z_0) + ah_n + h_n\psi(h_n)$$

by assumption, the second and third terms go to zero, so  $f(z_n) \rightarrow f(z_0)$ . □

**Example 1.2**

Let  $f : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$  be defined by  $f(z) = \frac{1}{z}$ . Then

$$\lim_{h \rightarrow 0} \frac{\frac{1}{z_0+h} - \frac{1}{z_0}}{h} = \lim_{h \rightarrow 0} \frac{-h}{h(z_0)(z_0+h)} = -\frac{1}{z_0^2}$$

so  $f'(z_0) = -\frac{1}{z_0^2}$ .

**Proposition 1.23**

Let  $\Omega \subseteq \mathbb{C}$  be open, and let  $f, g : \Omega \rightarrow \mathbb{C}$  be holomorphic at  $z_0$ . Then

1.  $f + g$  is holomorphic at  $z_0$ , and  $(f + g)' = f' + g'$ .
2.  $fg$  is holomorphic at  $z_0$ , and  $(fg)' = f'g + fg'$ .

3. If  $g(z_0) \neq 0$ , then  $\frac{f}{g}$  is well defined on an open disk around  $z_0$ , and  $\frac{f}{g}$  is holomorphic at  $z_0$  with  $\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$ .

*Proof.* Let  $\psi, \varphi$  be such that

$$\begin{aligned} f(z_0 + h) &= f(z_0) + f'(z_0)h + h\psi(h) \\ g(z_0 + h) &= g(z_0) + g'(z_0)h + h\varphi(h) \end{aligned}$$

Then

$$f(z_0 + h) + g(z_0 + h) = f(z_0) + g(z_0) + [f'(z_0) + g'(z_0)]h + h(\psi(h) + \varphi(h))$$

$\lim_{h \rightarrow 0} \varphi + \psi = 0$ , so the above shows that  $(f + g)' = f' + g'$ .

Letting

$$\phi(h) = f'(z_0)g'(z_0) + \psi(h)[g(z_0) + g'(z_0)] + \varphi(h)[f(z_0) + f'(z_0)] + \varphi(h)\psi(h)$$

which tends to 0 as  $h \rightarrow 0$ , we have

$$f(z_0 + h)g(z_0 + h) = f(z_0)g(z_0) + [f'(z_0)g'(z_0) + f'(z_0)g(z_0)]h + h\phi(h)$$

so  $(fg)' = f'g + g'f$ .

The quotient rule may be derived from the Chain Rule using the fact that  $\left(\frac{1}{z}\right)' = -\frac{1}{z^2}$  when  $z \neq 0$ .  $\square$

### Proposition 1.24: Chain Rule

Let  $\Omega, U \subseteq \mathbb{C}$  be open, and let  $f : \Omega \rightarrow U$  and  $g : U \rightarrow \mathbb{C}$ . Then  $g \circ f : \Omega \rightarrow \mathbb{C}$  is holomorphic and

$$(g \circ f)'(z) = g'(f(z))f'(z)$$

*Proof.* Using the alternative characterization of holomorphicity, we have

$$f(z_0 + h) = f(z_0) + f'(z_0)h + h\psi_f(h)$$

where  $\psi_f(h) \rightarrow 0$  as  $h \rightarrow 0$ . Similarly,

$$g(f(z_0) + w) = g(f(z_0)) + g'(f(z_0))w + w\psi_g(w)$$

Then

$$\begin{aligned} (g \circ f)(z_0 + h) &= g(f(z_0) + f'(z_0)h + h\psi_f(h)) \\ &= g(f(z_0)) + g'(f(z_0))(f'(z_0)h + h\psi_f(h)) + (f'(z_0)h + h\psi_f(h))\psi_g(f'(z_0)h + h\psi_f(h)) \\ &= g(f(z_0)) + g'(f(z_0))f'(z_0)h + h[\psi_f(h)g'(f(z_0)) + (f'(z_0) + \psi_f(h))\psi_g(f'(z_0)h + h\psi_f(h))] \end{aligned}$$

Note that

$$\lim_{h \rightarrow 0} \underbrace{\psi_f(h)}_{=0} g'(f(z_0)) + (f'(z_0) + \underbrace{\psi_f(h)}_{=0}) \underbrace{\psi_g(f'(z_0)h + h\psi_f(h))}_{=0} = 0$$

so  $(g \circ f)'(z_0) = g'(f(z_0))f'(z_0)$ .  $\square$



**Example 1.3**

Let  $f$  be a constant function. Then  $f$  is entire and  $f'(z) = 0$ .

We showed in Example 1.1 that the identity  $g(z) = z$  is entire with  $g'(z) = 1$ .

Combination of the two functions above, together with Proposition 1.23 gives

**Corollary 1.25**

Let  $p(z) = a_0 + a_1z + \dots + a_nz^n$ . Then  $p$  is entire and  $p'(z) = a_1 + 2a_2z + \dots + na_nz^{n-1}$ .

Let us consider a non-example.

**Example 1.4**

Let  $f(z) = \bar{z}$ , so that  $f(x + iy) = x - iy$ . This is a smooth function in the case of  $\mathbb{R}^2$ ; in fact since it is linear,  $Df = f$ , so that  $f$  has infinitely many derivatives.

However, in the complex case, we have

$$\frac{f(z+h) - f(z)}{h} = \frac{\overline{z+h} - \bar{z}}{h} = \frac{\bar{h}}{h}$$

But

$$\lim_{t \rightarrow 0} \frac{\bar{t}}{t} = 1$$

and

$$\lim_{t \rightarrow 0} \frac{\overline{it}}{it} = -1$$

so the limits disagree and  $f$  is not holomorphic at any  $z$ .

Consider some function  $f : \Omega \rightarrow \mathbb{C}$ . Let us denote its real and imaginary parts by  $u, v$ , respectively, so that

$$f(x + iy) = u(x, y) + iv(x, y)$$

( $u, v$  are defined on  $\Omega' \subseteq \mathbb{R}^2$  which is equivalent to  $\Omega$  in the obvious way.) This allows us to consider  $f$  as a pair of functions from  $\mathbb{R}^2 \rightarrow \mathbb{R}$ , which are surfaces lying in  $\mathbb{R}^3$ . We will investigate which choices of  $u, v$  may be associated with a holomorphic  $f$ .

Let  $h$  be a (small) complex number and write  $h = h_1 + ih_2$ . Then write

$$\frac{f(z+h) - f(z)}{h} = \frac{u(x+h_1, y+h_2) - u(x, y)}{h_1 + ih_2} + \frac{v(x+h_1, y+h_2) - v(x, y)}{h_1 + ih_2}$$

Let us consider what happens as  $h$  tends to 0 from different directions. For instance, suppose  $h$  is entirely real, so  $h_2 = 0$ . Then

$$\lim_{h_1 \rightarrow 0} \frac{f(z+h_1) - f(z)}{h_1} = \lim_{h_1 \rightarrow 0} \frac{(u+iv)(x+h_1, y) - (u+iv)(x, y)}{h_1} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

and similarly

$$\lim_{h_2 \rightarrow 0} \frac{f(z + h_2) - f(z)}{ih_2} = \frac{1}{i} \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y}$$

Then if  $f$  is holomorphic, then we can match components to get the following:

$$\begin{cases} \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \\ \frac{\partial v}{\partial y} = -\frac{\partial u}{\partial y} \end{cases} \quad (*)$$

The system of equations  $(*)$  are known as the **Cauchy-Riemann equations**. We have shown that these are a necessary conditions for  $f$  to be holomorphic; below we will show that if we also assume that the partials are continuous, then we have a sufficient condition. Later, we will show that  $u, v$  are necessarily continuously differentiable, so that these are equivalent characterizations. For now we will content ourselves with one direction:

**Theorem 1.26**

Let  $\Omega \subseteq \mathbb{C}$  be open and let  $f : \Omega \rightarrow \mathbb{C}$ . Let  $f = u + iv$ , where  $u, v : \Omega \rightarrow \mathbb{R}$  are continuously differentiable and satisfy the Cauchy-Riemann equations. Then  $f$  is holomorphic.

*Proof.* Consider the first-order Taylor expansion in two variables, which says that

$$u(x + h_1, y + h_2) = u(x, y) + \frac{\partial u}{\partial x} h_1 + \frac{\partial u}{\partial y} h_2 + h\psi_u(h)$$

where  $h = h_1 + ih_2$  and  $\psi_u(h) \rightarrow 0$  as  $h \rightarrow 0$ . Similarly

$$v(x + h_1, y + h_2) = v(x, y) + \frac{\partial v}{\partial x} h_1 + \frac{\partial v}{\partial y} h_2 + h\psi_v(h)$$

Now by assumption,  $f$  satisfies the Cauchy-Riemann equations, so

$$v(x + h_1, y + h_2) = v(x, y) - \frac{\partial u}{\partial y} h_1 + \frac{\partial u}{\partial x} h_2 + h\psi_v(h)$$

so

$$\begin{aligned} f(x + h) &= u(x + h_1, y + h_2) + iv(x + h_1, y + h_2) \\ &= u(x, y) + \frac{\partial u}{\partial x} h_1 + \frac{\partial u}{\partial y} h_2 + h\psi_u(h) + iv(x, y) - i \frac{\partial u}{\partial y} h_1 + i \frac{\partial u}{\partial x} h_2 + ih\psi_v(h) \\ &= f(x, y) + (h_1 + ih_2) \left( \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} \right) + h \underbrace{(\psi_u(h) + i\psi_v(h))}_{\psi(h)} \\ &= f(z) + h \left( \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} \right) + h\psi(h) \end{aligned}$$

so  $f$  is holomorphic using the alternative characterization and

$$f'(z) = f'(x, y) = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$$

□

Let  $f$  be a complex valued function of the form  $f(x + iy) = u(x, y) + iv(x, y)$ . Associate with it a  $\mathbb{R}^2$ -valued function

$$F(x, y) = (u(x, y), v(x, y))$$

Recall that its Jacobian matrix is

$$J_F(x, y) = \begin{bmatrix} u_x & u_y \\ v_x & v_y \end{bmatrix} \quad (**)$$

and that  $F$  is differentiable in the real sense if it is true that

$$\lim_{(h_1, h_2) \rightarrow (0, 0)} \frac{F(x + h_1, y + h_2) - F(x, y) - J_F(x, y) \begin{bmatrix} h_1 \\ h_2 \end{bmatrix}}{|(h_1, h_2)|} = 0$$

Comparing this to the complex condition

$$\lim_{h \rightarrow 0} \frac{f(z + h) - f(z) - f'(z)h}{h} = 0$$

we can see that complex differentiability requires  $J_f(x, y)$  to be of the form of multiplying by some complex number. This happens if and only if

$$J_F(x, y) = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

for  $a, b \in \mathbb{R}$ . By reconciling this with  $(**)$  we recover the Cauchy-Riemann equations  $(*)$ .

#### Definition 1.27

Let  $f : \Omega \rightarrow \mathbb{C}$  with  $\Omega$  open. Then we define

$$\frac{\partial f}{\partial x} := \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

and

$$\frac{\partial f}{\partial y} := \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y}$$

We further define

$$\frac{\partial f}{\partial z} := \frac{1}{2} \left( \frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right)$$

and

$$\frac{\partial f}{\partial \bar{z}} := \frac{1}{2} \left( \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right)$$

### Proposition 1.27

Let  $\Omega \subseteq \mathbb{C}$  be open and let  $f : \Omega \rightarrow \mathbb{C}$  be of the form  $f(x + iy) = u(x, y) + iv(x, y)$ . If  $f$  is holomorphic on  $\Omega$ , then

1.  $\frac{\partial f}{\partial \bar{z}} = 0$ .
2.  $\frac{\partial f}{\partial z} = \frac{\partial f}{\partial x} = -i \frac{\partial f}{\partial y}$ .
3.  $\frac{\partial f}{\partial z} = f'$ .
4.  $f'(z_0) = 2 \frac{\partial u}{\partial \bar{z}}(z_0)$ .
5.  $F = (u(x, y), v(x, y))$  is differentiable in the real sense, and  $\det J_F(x, y) = |f'(x + iy)|^2$ .

*Proof.* 1. By Cauchy-Riemann,

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + i \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} \right) = 0$$

2. By part 1,

$$\frac{\partial f}{\partial z} = \frac{\partial f}{\partial z} + \frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} + \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) = \frac{\partial f}{\partial x}$$

Similarly,

$$\frac{\partial f}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} - \frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right) = -i \frac{\partial f}{\partial y}$$

3. Take  $h_1$  real and  $z_0 \in \Omega$ . Then

$$\frac{\partial f}{\partial x}(z_0) = \lim_{h_1 \rightarrow 0} \frac{f(z_0 + h_1) - f(z_0)}{h_1} = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = f'(z_0)$$

By part 2 the conclusion follows.

4. By parts 2 and 3,

$$f' = \frac{\partial f}{\partial z} = \frac{\partial f}{\partial x} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} = 2 \frac{\partial u}{\partial \bar{z}}$$

5. We have

$$\det J_F(x, y) = \frac{\partial u}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \frac{\partial v}{\partial x}$$

By the Cauchy-Riemann equations and parts 2 and 3 this is

$$\left( \frac{\partial u}{\partial x} \right)^2 + \left( \frac{\partial v}{\partial x} \right)^2 = |f'(x + iy)|^2$$

□

## 1.4 Power Series

We will now discuss power series, which are defined similarly to the real case. They will initially serve as a valuable example of holomorphic functions. Later we will see that they are actually the *only* example, which justifies particular attention in their study.

### Definition 1.28

A **power series** (centered around  $z_0$ ) is a function  $f$  of the form

$$f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n := \lim_{N \rightarrow \infty} \sum_{n=0}^N a_n(z - z_0)^n$$

where  $\{a_n\} \subseteq \mathbb{C}$ , which is defined wherever the right hand limit converges.

Note that it is certainly a necessary condition that  $a_n z^n \rightarrow 0$  as  $n \rightarrow \infty$ , since

$$\lim_{n \rightarrow \infty} a_n z^n = \lim_{n \rightarrow \infty} \left[ \sum_{k=0}^n a_k z^k - \sum_{k=0}^{n-1} a_k z^k \right] = \lim_{n \rightarrow \infty} \left[ \sum_{k=0}^n a_k z^k \right] - \lim_{n \rightarrow \infty} \left[ \sum_{k=0}^{n-1} a_k z^k \right] = 0$$

In this section, we will first consider only power series which are centered at 0.

### Definition 1.29

We say that a power series  $\sum_{n=0}^{\infty} a_n z^n$  **converges absolutely** (in the complex sense) if the series

$$\sum_{n=0}^{\infty} |a_n| \cdot |z|^n$$

converges in the real sense.

### Proposition 1.28

If  $\sum_{n=0}^{\infty} a_n z^n$  converges absolutely, then it converges (in the complex sense).

*Proof.* Let  $\varepsilon > 0$ .

Write

$$S_N(z) = \sum_{n=0}^N a_n z^n$$

Since

$$\sum_{n=0}^{\infty} |a_n| \cdot |z|^n$$

converges, there exists  $N_0 \in \mathbb{N}$  such that

$$\sum_{n=N_0+1}^{\infty} |a_n| \cdot |z|^n < \varepsilon$$

Then for  $M, N \geq N_0$ , assume without loss of generality that  $M < N$ . Then we have

$$|S_N(z) - S_M(z)| = \left| \sum_{n=M+1}^N a_n z^n \right| \leq \sum_{n=M+1}^N |a_n| \cdot |z|^n \leq \sum_{n=N_0+1}^{\infty} |a_n| \cdot |z|^n < \varepsilon$$

so  $\{S_N(z)\}$  is a Cauchy sequence, and thus

$$\sum_{n=0}^{\infty} a_n z^n$$

converges. □

Recall that in the single real variable case, we found that power series converge on some interval (possibly open, closed, or half-open) which is centered around 0 (or any other point of expansion). An analogous statement is true here, with the interval replaced by a disk.

### Theorem 1.29

For any power series  $\sum_{n=0}^{\infty} a_n z^n$  there exists  $0 \leq R \leq \infty^a$  (called the **radius of convergence**) such that for any  $z \in \mathbb{C}$ :

1. If  $|z| < R$ , then  $\sum_{n=0}^{\infty} a_n z^n$  converges absolutely.
2. If  $|z| > R$ , then  $\sum_{n=0}^{\infty} a_n z^n$  does not converge.

Moreover,  $R$  is given by **Hadamard's formula**:

$$R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}}$$

---

<sup>a</sup> $R = \infty$  means the condition  $|z| < R$  is satisfied for all  $z \in \mathbb{C}$ .

This theorem says that we have absolute convergence inside the disk  $\mathbb{D}_R$  (called the **disk of convergence**), and divergence outside of it. As in the real case, this theorem makes no statement about convergence on the boundary of the disk.

*Proof.* Denote  $L = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}$ . Suppose that  $L \neq 0, \infty$ . If  $|z| < \frac{1}{L} = R$ , then  $L|z| < 1$ , so there exists  $\varepsilon > 0$  such that  $(L + \varepsilon)|z| = r < 1$ . By the definition of  $\limsup$ , there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,

$$\sqrt[n]{|a_n|} < L + \varepsilon \implies |a_n| < (L + \varepsilon)^n \implies |a_n| |z|^n < ((L + \varepsilon)|z|)^n = r^n$$

so  $\sum_{n=0}^{\infty} a_n z^n$  is dominated by the absolutely convergent geometric series  $\sum_{n=0}^{\infty} r^n$  and thus converges absolutely.

On the other hand, if  $|z| > R = \frac{1}{L}$ , then  $L|z| > 1$  so there exists a subsequence  $\{a_{n_k}\}$  such that

$$\sqrt[n_k]{|a_{n_k}|} \cdot |z| > 1$$

for all  $k$ . Then

$$|a_{n_k}| \cdot |z|^{n_k} > 1$$

which does not tend to 0 as  $k \rightarrow \infty$ , so convergence is impossible.

If  $L = 0$ , then  $R = \infty$ . For any  $z$  there exists  $N$  such that  $n \geq N$  implies that

$$\sqrt[n]{a_n} < \frac{1}{|z|}$$

Then

$$|a_n||z|^n < r < 1$$

So

$$\sum_{n=N}^{\infty} |a_n||z|^n$$

is dominated by  $\sum_{n=N}^{\infty} r^n$ , and thus converges for  $|z| < R = \infty$ .

If  $L = \infty$ , then  $L|z| > 1$  and we apply the argument for the case  $|z| > R$ . □

#### Example 1.5

The power series

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}$$

has  $R = \infty$ , since the real-valued power series

$$\sum_{n=0}^{\infty} \frac{|z|^n}{n!} = e^{|z|}$$

converges absolutely everywhere. This also allows us to define the **exponential** of  $z$  as

$$e^z := \sum_{n=0}^{\infty} \frac{z^n}{n!}$$

#### Example 1.6

The power series

$$\sum_{n=0}^{\infty} z^n$$

has radius of convergence 1. This can be seen either by direct computation in the real case, or using Hadamard's formula and the fact that each  $a_n$  is 1:

$$R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{1}} = \frac{1}{1} = 1$$

Moreover, this power series satisfies the equation

$$\sum_{n=0}^{\infty} z^n = \frac{1}{1-z}$$

This can be seen using the identity for partial sums (which holds in all fields)

$$\sum_{n=0}^N z^n = \frac{1 - z^{N+1}}{1 - z}$$

so

$$\lim_{N \rightarrow \infty} \frac{1 - z^{N+1}}{1 - z} = \frac{1}{1 - z}$$

### Definition 1.30

We define the **trigonometric functions** in terms of power series:

$$\begin{aligned}\cos z &:= \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!} \\ \sin z &:= \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!}\end{aligned}$$

Note that both of the above series converge with  $R = \infty$ . Moreover, we can observe that

$$\begin{aligned}\cos z &= \frac{e^{iz} + e^{-iz}}{2} \\ \sin z &= \frac{e^{iz} - e^{-iz}}{2i}\end{aligned}$$

and that this is consistent with our previous definition in terms of the identity

$$e^{iz} = \cos z + i \sin z$$

We now prove a fundamental fact about power series which, while analogous to the real case, will have farther reaching implications for us.

### Theorem 1.30

The function  $f(z) = \sum_{n=0}^{\infty} a_n z^n$  is holomorphic on the disk of convergence  $\mathbb{D}_R$ . Moreover, its derivative is given by

$$f'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} z^n$$

which has the same radius of convergence.

*Proof.* To show that the radius of convergence is the same, simply apply Hadamard's formula to the new power series. Letting  $R'$  be the radius of convergence of  $\sum_{n=1}^{\infty} n a_n z^{n-1}$ , we have

$$\frac{1}{R'} = \limsup_{n \rightarrow \infty} \sqrt[n]{(n+1)|a_n|} = \limsup_{n \rightarrow \infty} \sqrt[n]{n+1} \sqrt[n]{|a_n|}$$



But  $\sqrt[n]{n+1} \rightarrow 1$  so

$$\limsup_{n \rightarrow \infty} \sqrt[n]{n+1} \sqrt[n]{|a_n|} = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \frac{1}{R}$$

so  $R' = R$ .

Denote

$$g(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}$$

We want to show that  $f' = g$ . It suffices to show that

$$\frac{f(z_0 + h) - f(z_0)}{h} - g(z_0) \xrightarrow{h \rightarrow 0} 0$$

whenever  $z_0 \in \mathbb{D}_R$ .

To show this, pick any  $z_0 \in \mathbb{D}_R$  and let  $\varepsilon > 0$ . Then there exists  $r$  such that  $|z_0| < r < R$ . Then whenever  $|h| < r - |z_0|$  we have  $|z_0 + h| \leq |z_0| + |h| < r < R$ . So for small  $h$ ,  $z_0 + h \in \mathbb{D}_R$ .

Denote the partial sums and error terms by

$$S_N(z) = \sum_{n=1}^N a_n z^n$$

$$E_N(z) = \sum_{n=N+1}^{\infty} a_n z^n$$

By our observation about Hadamard's formula,

$$\sum_{n=1}^{\infty} n |a_n| z^{n-1}$$

converges, so there exists  $N_0 \in \mathbb{N}$  such that

$$\sum_{n=N_0+1}^{\infty} n |a_n| |z|^{n-1} < \frac{\varepsilon}{3} \quad (1)$$

$S_{N_0}$  is a polynomial, which we showed is holomorphic. So

$$\frac{S_{N_0}(z_0 + h) - S_{N_0}(z_0)}{h} - S'_{N_0}(z_0) = \frac{S_{N_0}(z_0 + h) - S_{N_0}(z_0)}{h} - \sum_{n=1}^{N_0} n a_n z_0^{n-1} \xrightarrow{h \rightarrow 0} 0$$

Thus there exists  $\delta > 0$  such that whenever  $|h| < \delta$ ,

$$\left| \frac{S_{N_0}(z_0 + h) - S_{N_0}(z_0)}{h} - S'_{N_0}(z_0) \right| < \frac{\varepsilon}{3} \quad (2)$$

The difference quotient

$$\left| \frac{f(z_0 + h) - f(z_0)}{h} - g(z_0) \right|$$

may be written as

$$\begin{aligned} & \left| \frac{S_{N_0}(z_0 + h) - S_{N_0}(z_0)}{h} - S'_{N_0}(z_0) + S'_{N_0}(z_0) - g(z_0) + \frac{E_{N_0}(z_0 + h) - E_{N_0}(z_0)}{h} \right| \\ & \leq \left| \frac{S_{N_0}(z_0 + h) - S_{N_0}(z_0)}{h} - S'_{N_0}(z_0) \right| + |S'_{N_0}(z_0) - g(z_0)| + \left| \frac{E_{N_0}(z_0 + h) - E_{N_0}(z_0)}{h} \right| \end{aligned}$$

For small  $h$ , the first term is less than  $\frac{\varepsilon}{3}$  by (2). Also,

$$|S'_{N_0}(z_0) - g(z_0)| = \left| - \sum_{n=N_0+1}^{\infty} n a_n z_0^{n-1} \right| \leq \sum_{n=N_0+1}^{\infty} n |a_n| \cdot |z_0|^{n-1} < \frac{\varepsilon}{3}$$

by (1). Lastly,

$$\left| \frac{E_{N_0}(z_0 + h) - E_{N_0}(z_0)}{h} \right| = \left| \sum_{n=N_0+1}^{\infty} a_n \frac{(z_0 + h)^n - z_0^n}{h} \right|$$

Using the identity

$$a^n - b^n = (a - b) \sum_{k=0}^{n-1} a^k b^{n-1-k}$$

for  $a = z_0 + h, b = z_0$  we have

$$\begin{aligned} & \left| \frac{\sum_{n=N_0+1}^{\infty} a_n ((z_0 + h)^n - z_0^n)}{h} \right| = \left| \sum_{n=N_0+1}^{\infty} a_n \left( \sum_{k=0}^{n-1} (z_0 + h)^k z_0^{n-1-k} \right) \right| \\ & \leq \sum_{n=N_0+1}^{\infty} |a_n| \sum_{k=0}^{n-1} |z_0 + h|^k |z_0|^{n-1-k} < \sum_{n=N_0+1}^{\infty} |a_n| n r^n < \frac{\varepsilon}{3} \end{aligned}$$

where the last inequality follows from (1). Thus

$$\left| \frac{f(z_0 + h) - f(z_0)}{h} - g(z_0) \right| < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon$$

□

### Corollary 1.31

A power series is infinitely complex differentiable on its disk of convergence.

*Proof.* Induct using Theorem 1.30. □

We now note that for the case of power series  $\sum_{n=0}^{\infty} a_n (z - z_0)^n$  centered around arbitrary  $z_0$ , they converge on  $\mathbb{D}_R(z_0)$ , where  $R$  is the radius of convergence of  $\sum_{n=0}^{\infty} a_n z^n$ . In other words, the behavior is identical, merely translated. Thus it suffices to consider power series centered around 0.

**Definition 1.31**

Let  $\Omega \subseteq \mathbb{C}$  be open and  $z_0 \in \Omega$ . Then  $f : \Omega \rightarrow \mathbb{C}$  is said to be **analytic** at  $z_0$  if there exists  $r > 0$  such that  $\mathbb{D}_r(z_0) \subseteq \Omega$  and there exist coefficients  $\{a_n\} \subseteq \mathbb{C}$  such that

$$\sum_{n=0}^{\infty} a_n z^n$$

converges absolutely on  $\mathbb{D}_r$  and

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$$

on  $\mathbb{D}_r(z_0)$ .

We say that  $f$  is analytic on  $\Omega$  if it is analytic at each  $z_0 \in \Omega$ .

By Theorem 1.30, each analytic function is (infinitely) holomorphic. We will show later that every holomorphic function is also analytic. Thus the terms are often used interchangeably, but we will not do so until we have proved this fact.

## 1.5 Integration on Curves

We will soon show that the behavior of analytic functions can be well understood by studying the behavior of those functions when integrated over various curves in the plane. This will motivate the particular definitions of an integral over a curve that appear in this section.

**Definition 1.32**

A **parameterized curve** is a continuous function  $z : [a, b] \rightarrow \mathbb{C}$ . Writing  $z(t) = z_1(t) + iz_2(t)$ , we say that  $z$  is a **smooth curve** if  $z'_1(t), z'_2(t)$  exist for all  $t \in [a, b]$ .

$z$  is said to be **piecewise smooth** if there exist  $a = a_0 < a_1 < \dots < a_n = b$  such that  $z$  is smooth on  $[a_k, a_{k+1}]$  for each  $k$ .

$z$  is a **closed curve** if  $z(a) = z(b)$ .

Note that in our definition of piecewise smooth curves,  $z$  must be continuous at each  $a_k$ , but the left and right derivatives need not coincide.

We will adopt the convention that all curves are assumed to be piecewise smooth, unless stated otherwise.

A parameterized curve traces out a particular image in the complex plane, which is intuitively a curve in the plane. Thus it is useful to distinguish the function  $z$  and its image.

**Definition 1.33**

Two parameterized curves  $z_1 : [a, b] \rightarrow \mathbb{C}$  and  $z_2 : [c, d] \rightarrow \mathbb{C}$  are said to be **equivalent curves** if there exists a continuously differentiable bijection  $t : [a, b] \rightarrow [c, d]$  such that  $t'(s) > 0$  and  $z_2(s) = z_1(t(s))$  for all  $s \in [a, b]$ .

The condition that  $t'(s) > 0$  could be equivalently stated as saying that  $t(a) = c$  and  $t(b) = d$  (so that the direction does not change). It follows that  $z_1([a, b]) = z_2([c, d])$ .

**Definition 1.34**

A **curve** is an equivalence class of parameterized curves.

Note that for a given curve, there is another curve which has the same image in  $\mathbb{C}$ , but with the opposite orientation.

**Definition 1.35**

If  $\gamma \subseteq \mathbb{C}$  is a curve, and  $z : [a, b] \rightarrow \mathbb{C}$  is a parameterization of  $\gamma$ , then  $\gamma^-$  is the curve parameterized by  $z(a + b - t)$ .

A commonly used curve is the circle, so it is convenient to define some conventional parameterizations of the circle:

**Definition 1.36**

The circle with radius  $r > 0$  and center  $z \in \mathbb{C}$  is

$$C_r(z_0) := \partial\mathbb{D}_r(z_0)$$

The **positive orientation** of  $C_r(z)$  is the curve parameterized by  $z : [-\pi, \pi] \rightarrow \mathbb{C}$  and

$$z(\theta) = z_0 + re^{i\theta}$$

The **negative orientation** is the curve parameterized by

$$z^-(\theta) = z_0 + re^{-i\theta}$$

Now, let us define the manner in which we will integrate functions over smooth planar curves.

**Definition 1.37**

Let  $\gamma \subseteq \mathbb{C}$  be a smooth curve. Let  $f : \gamma \rightarrow \mathbb{C}$  be continuous. Let  $z : [a, b] \rightarrow \mathbb{C}$  be a smooth parameterization of  $\gamma$ . Then define the **contour integral** of  $f$  along  $\gamma$  with

respect to  $z$  to be

$$\int_{\gamma, z} f(z) dz = \int_a^b f(z(t))z'(t) dt = \int_a^b \operatorname{Re}(f(z(t))z'(t)) dt + i \int_a^b \operatorname{Im}(f(z(t))z'(t)) dt$$

As Proposition 1.32 shows, this value is independent of the choice of  $z$ . Thus, we take this common value to be the contour integral of  $f$  along  $\gamma$ .

### Proposition 1.32

If  $\gamma$  is a smooth curve,  $f : \gamma \rightarrow \mathbb{C}$  is continuous, and  $z_1, z_2$  are two parameterizations of  $\gamma$ , then

$$\int_{\gamma, z_1} f(z) dz = \int_{\gamma, z_2} f(z) dz$$

*Proof.* Since  $z_1, z_2$  are equivalent, we write  $z_1(s) = z_2(t(s))$ . Then

$$\begin{aligned} \int_{\gamma, z_1} f(z) dz &= \int_a^b f(z_1(s))z'_1(s) ds = \int_a^b f(z_2(t(s)))z'_2(t(s))t'(s) ds \\ &= \int_{t(a)}^{t(b)} f(z_2(t))z'_2(t) dt = \int_c^d f(z_2(t))z'_2(t) dt = \int_{\gamma, z_2} f(z) dz \quad \square \end{aligned}$$

The above proof shows that we may define contour integrals with respect to a curve, without referring to a specific parameterization.

If  $\gamma$  is only piecewise smooth, and some parameterization  $z : [a, b] \rightarrow \mathbb{C}$  is smooth on  $[a_k, a_{k+1}]$  for  $a = a_0 < a_1 < \dots < a_n = b$ , then we define the integral to be

$$\int_{\gamma} f(z) dz = \sum_{k=0}^{n-1} \int_{a_k}^{a_{k+1}} f(z(t))z'(t) dt$$

Similar work as above shows that this is also independent of the parameterization.

### Definition 1.38

Let  $\gamma \subseteq \mathbb{C}$  be a curve parameterized by  $z : [a, b] \rightarrow \mathbb{C}$ . Then the **length** of  $\gamma$  is

$$\operatorname{length}(\gamma) = \int_a^b |z'(t)| dt$$

The following example will be extremely instructive for later applications.

**Example 1.7**

Let  $\gamma = \partial\mathbb{D}$ . Let  $f : \gamma \rightarrow \mathbb{C}$  be continuous. Let  $z$  be the parameterization

$$z(\theta) = e^{i\theta}$$

Then

$$z'(\theta) = ie^{i\theta}$$

So

$$\int_{\gamma} f(z) \, dz = \int_0^{2\pi} f(e^{i\theta}) ie^{i\theta} \, d\theta$$

For instance, take

$$f(z) = \frac{1}{z}$$

Then we have

$$\int_{\gamma} \frac{1}{z} \, dz = \int_0^{2\pi} \frac{ie^{i\theta}}{e^{i\theta}} \, d\theta = \int_0^{2\pi} i \, d\theta = 2\pi i$$

The length of  $\gamma$  is

$$\int_0^{2\pi} |ie^{i\theta}| \, d\theta = \int_0^{2\pi} 1 \, d\theta = 2\pi$$

as we expect.

Let us briefly cover some properties of the contour integral.

**Proposition 1.33**

Let  $\gamma \subseteq \mathbb{C}$  be a curve,  $f, g : \gamma \rightarrow \mathbb{C}$  be continuous, and  $\alpha, \beta \in \mathbb{C}$ . Then

1.  $\int_{\gamma}$  is linear:

$$\int_{\gamma} (\alpha f + \beta g) \, dz = \alpha \int_{\gamma} f \, dz + \beta \int_{\gamma} g \, dz$$

2.  $\int_{\gamma}$  is reversed by orientation:

$$\int_{\gamma} f \, dz = - \int_{-\gamma} f \, dz$$

3. The following inequality holds analogous to the triangle inequality:

$$\left| \int_{\gamma} f \, dz \right| \leq \left( \sup_{z \in \gamma} |f(z)| \right) \text{length}(\gamma)$$

*Proof.* 1. Follows from properties of the Riemann integral.

2. Homework (follows from reparameterizing).

3. First, we show the following:

**Claim**

Let  $h : [a, b] \rightarrow \mathbb{C}$  be continuous. Then

$$\left| \int_a^b h(t) dt \right| \leq \int_a^b |h(t)| dt$$

*Proof.* Write  $\int_a^b h(t) dt = re^{i\theta}$  for appropriate  $r, \theta$ . Then

$$\begin{aligned} r &= \left| \int_a^b h(t) dt \right| = e^{-i\theta} \int_a^b h(t) dt \\ &= \int_a^b e^{-i\theta} h(t) dt = \operatorname{Re} \left( \int_a^b e^{-i\theta} h(t) dt \right) \\ &= \int_a^b \operatorname{Re}(e^{-i\theta} h(t)) dt \leq \int_a^b |h(t)| dt \end{aligned}$$

■

Now, suppose that  $z : [a, b] \rightarrow \mathbb{C}$  is piecewise smooth with  $a = a_0 < \dots < a_n = b$ . Then by definition,

$$\begin{aligned} \left| \int_{\gamma} f(z) dz \right| &= \left| \sum_{k=0}^{n-1} \int_{a_k}^{a_{k+1}} f(z(t)) z'(t) dt \right| \leq \sum_{k=0}^{n-1} \left| \int_{a_k}^{a_{k+1}} f(z(t)) z'(t) dt \right| \\ &\leq \sum_{k=0}^{n-1} \int_{a_k}^{a_{k+1}} |f(z(t))| \cdot |z'(t)| dt \leq \sum_{k=0}^{n-1} \int_{a_k}^{a_{k+1}} \left( \sup_{z \in \gamma} |f(z(t))| \right) |z'(t)| dt \\ &= \left( \sup_{z \in \gamma} |f(z(t))| \right) \sum_{k=0}^{n-1} \int_{a_k}^{a_{k+1}} |z'(t)| dt = \left( \sup_{z \in \gamma} |f(z(t))| \right) \operatorname{length}(\gamma) \quad \square \end{aligned}$$

In vector calculus, the fundamental theorem of line integrals shows that calculation of line integrals can often be reduced to evaluating an antiderivative at its endpoints. The same is true in  $\mathbb{C}$ .

**Definition 1.39**

Let  $\Omega \subseteq \mathbb{C}$  be open. Let  $f : \Omega \rightarrow \mathbb{C}$ . Then we say that  $F : \Omega \rightarrow \mathbb{C}$  is a **primitive** of  $f$  on  $\Omega$  if  $F$  is holomorphic on  $\Omega$  and

$$F'(z) = f(z)$$

for all  $z \in \Omega$ .

**Theorem 1.34**

Let  $\Omega \subseteq \mathbb{C}$  be open. Let  $f : \Omega \rightarrow \mathbb{C}$  be continuous and suppose  $F : \Omega \rightarrow \mathbb{C}$  is a primitive of  $f$ . Then for any curve  $\gamma \subseteq \Omega$  joining  $z_1, z_2$ , we have

$$\int_{\gamma} f(z) dz = F(z_2) - F(z_1)$$

*Proof.* This is effectively the Fundamental Theorem of Calculus. Take a piecewise smooth parameterization  $z : [a, b] \rightarrow \mathbb{C}$  of  $\gamma$  and pick  $a = a_0 < \dots < a_n = b$  so that  $z$  is smooth on each subinterval. Then

$$\begin{aligned} \int_{\gamma} f(z) dz &= \sum_{k=0}^{n-1} \int_{a_k}^{a_{k+1}} f(z(t)) z'(t) dt = \sum_{k=0}^{n-1} \int_{a_k}^{a_{k+1}} F(z(t))' dt \\ &= \sum_{k=0}^{n-1} (F(z(a_{k+1})) - F(z(a_k))) = F(z(a_n)) - F(z(a_0)) = F(z_2) - F(z_1) \quad \square \end{aligned}$$

**Corollary 1.35**

Let  $\Omega \subseteq \mathbb{C}$  be open,  $f : \Omega \rightarrow \mathbb{C}$  continuous, and  $F$  a primitive for  $f$  on  $\Omega$ . Then for any *closed* curve  $\gamma \subseteq \Omega$ ,

$$\int_{\gamma} f(z) dz = 0$$

Although the above theorem is powerful, its assumptions sometimes fail to be satisfied in subtle ways.

**Example 1.8**

Let  $\Omega = \mathbb{C} \setminus \{0\}$ . Let  $f : \Omega \rightarrow \mathbb{C}$  be  $f(z) = \frac{1}{z}$ . We showed that

$$\int_{\partial \mathbb{D}} \frac{1}{z} dz = 2\pi i \neq 0$$

This shows that  $\frac{1}{z}$  does not have a primitive on  $\Omega$ . This is an interesting contrast to the real case, where  $\ln x$  is a primitive for  $\frac{1}{x}$  on  $\mathbb{R} \setminus \{0\}$ . (This is due to the fact that 0 is a branch point for the logarithm in  $\mathbb{C}$ .)

**Corollary 1.36**

Let  $\Omega \subseteq \mathbb{C}$  be a region. Let  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic on  $\Omega$  and  $f'(z) = 0$  for all  $z \in \Omega$ . Then  $f$  is constant.

*Proof.* Pick  $z_1, z_2 \in \Omega$ .  $\Omega$  is path connected since it is a region, so let  $\gamma \subseteq \Omega$  be a curve from



$z_1$  to  $z_2$ .  $f$  is a primitive for  $f'$ , so

$$f(z_2) - f(z_1) = \int_{\gamma} f'(z) \, dz = \int_{\gamma} 0 \, dz = 0$$

and thus  $f(z_2) = f(z_1)$ . So  $f$  is constant. □

## Chapter 2

# Cauchy's Theorem and Applications

Up until this point, the results we have shown about contour integrals have been analogous to line integrals in  $\mathbb{R}^n$ . In this chapter, we will see that integrating in the complex setting results in unique results. The main theorem we will prove is as follows:

### Theorem: Cauchy's Theorem for a Disk

Let  $f : \mathbb{D} \rightarrow \mathbb{C}$  be holomorphic on  $\mathbb{D}$ . Then for any closed curve  $\gamma \subseteq \mathbb{D}$ ,

$$\int_{\gamma} f(z) dz = 0$$

In order to prove that, we will first prove some intermediate results.

## 2.1 Goursat's Theorem

We will first prove a version of Cauchy's Theorem for a specific type of curve.

### Definition 2.1

A **triangle**  $T$  is a subset of  $\mathbb{C}$  which consists of three line segments and the region between them. The boundary of  $T$  is the line segments between them, which by convention is taken with the counterclockwise parameterization.

### Theorem 2.1: Cauchy-Goursat

Let  $\Omega \subseteq \mathbb{C}$  be open. Let  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic on  $\Omega$ , and let  $T \subseteq \Omega$  be a triangle. Then

$$\int_{\partial T} f(z) dz$$

*Proof.* Let us bisect the line segments of the triangle and draw a triangle between them. This creates four subtriangles, which we denote  $S_1, S_2, S_3, S_4$ .




Figure 2.1: Subdivision of  $T$

Each triangle made this way is similar to the original. Then consider the sum

$$\int_{\partial S_1} f(z) dz + \int_{\partial S_2} f(z) dz + \int_{\partial S_3} f(z) dz + \int_{\partial S_4} f(z) dz$$

Since we maintain the counterclockwise orientation, the edges of  $S_3$  will be integrated along in both directions, which cancels out. Thus the above expression is equal to

$$\int_{\partial T} f(z) dz$$

Thus

$$\left| \int_{\partial T} f(z) dz \right| \leq 4 \max \left\{ \left| \int_{\partial S_i} f(z) dz \right| \right\}$$

Suppose that the largest integral occurs over  $S_j$ . Then set  $T_1 = S_j$  and subdivide again.




Figure 2.2: Continued subdivisions

Thus we get a nested sequence of triangles satisfying

$$\begin{cases} T = T_0 \supseteq T_1 \supseteq \dots \\ \left| \int_{\partial T_n} f(z) dz \right| \leq 4 \left| \int_{\partial T_{n+1}} f(z) dz \right| \\ \text{diam}(T_{n+1}) = \frac{1}{2} \text{diam}(T_n) \\ \text{length}(\partial T_{n+1}) = \frac{1}{2} \text{length}(\partial T_n) \end{cases}$$

It follows that

$$\begin{cases} \left| \int_{\partial T} f(z) dz \right| \leq 4^n \left| \int_{\partial T_n} f(z) dz \right| \\ \text{diam}(T_n) = \frac{1}{2^n} \text{diam}(T) \\ \text{length}(\partial T_n) = \frac{1}{2^n} \text{length}(\partial T) \end{cases}$$

Since the diameters go to 0 and each  $T_i$  is compact, there exists a unique point  $\omega$  in each of the  $T_i$ . Since  $f$  is holomorphic at  $\omega$ , we can write

$$f(z) = f(\omega) + f'(\omega)(z - \omega) + \psi(z)(z - \omega)$$

where  $\lim_{z \rightarrow \omega} \psi(z) = 0$ . So for any  $n$ ,

$$\int_{\partial T_n} f(z) dz = \int_{\partial T_n} (f(\omega) + f'(\omega)(z - \omega)) dz + \int_{\partial T_n} \psi(z)(z - \omega) dz$$

But notice that

$$\left( f(\omega)z + \frac{1}{2} f'(\omega)(z - \omega)^2 \right)' = f(\omega) + f'(\omega)(z - \omega)$$

so we have a primitive, and it follows that

$$\int_{\partial T_n} (f(\omega) + f'(\omega)(z - \omega)) dz = 0$$

Now,

$$\begin{aligned} \left| \int_{\partial T} f(z) dz \right| &\leq 4^n \left| \int_{\partial T_n} f(z) dz \right| \\ &= 4^n \left| \int_{\partial T_n} \psi(z)(z - \omega) dz \right| \\ &\leq 4^n \left( \sup_{z \in \partial T_n} [\psi(z)(z - \omega)] \right) \text{length}(\partial T_n) \\ &= 2^n \text{length}(\partial T) \left( \text{diam}(T_n) \sup_{z \in \partial T_n} \psi(z) \right) \\ &= \text{length}(\partial T) \text{diam}(T) \sup_{z \in \partial T_n} \psi(z) \end{aligned}$$

Since

$$\lim_{z \rightarrow \omega} \psi(z) = 0$$

we can make  $\sup_{z \in \partial T_n} \psi(z)$  arbitrarily small by considering large enough  $n$ . It follows that

$$\lim_{n \rightarrow \infty} \sup_{z \in \partial T_n} \psi(z) = 0$$

and thus

$$\left| \int_{\partial T} f(z) dz \right| = 0 \quad \square$$

We will now show that holomorphic functions locally have primitives. We will adopt the notation that for  $z, \omega \in \mathbb{C}$ ,  $[z, \omega]$  represents the line segment joining  $z$  and  $\omega$ . Specifically, it can be parameterized by

$$t \mapsto (1-t)z + t\omega, \quad t \in [0, 1]$$

### Theorem 2.2

Let  $z_0 \in \mathbb{C}$ ,  $r > 0$ , and  $f : \mathbb{D}_r(z_0) \rightarrow \mathbb{C}$  be holomorphic on  $\mathbb{D}_r(z_0)$ . Then  $f$  has a primitive on  $\mathbb{D}_r(z_0)$ .

*Proof.* Take some  $z \in \mathbb{D}_r(z_0)$ . Define

$$F(z) = \int_{[z_0, z]} f(\omega) d\omega$$

We claim that  $F$  is holomorphic on  $\mathbb{D}_r(z_0)$  and that it is a primitive for  $f$ . Let  $h$  be small and consider the triangle  $T_h$  between  $z_0, z, z+h$ .




Figure 2.3: Auxiliary triangle construction

By Cauchy-Goursat,

$$0 = \int_{\partial T_h} f(\omega) d\omega = \int_{[z_0, z]} f(\omega) d\omega + \int_{[z, z+h]} f(\omega) d\omega - \int_{[z_0, z+h]} f(\omega) d\omega$$

Thus

$$\begin{aligned} F(z+h) - F(z) &= F(z+h) - F(z) + \int_{\partial T_h} f(\omega) d\omega = \int_{[z, z+h]} f(\omega) d\omega \\ &= h \int_0^1 f((1-t)z + t(z+h)) dt \end{aligned}$$

and

$$\frac{F(z+h) - F(z)}{h} - f(z) = \int_0^1 (f((1+t)z + t(z+h)) - f(z)) dt$$

Since  $f$  is continuous at  $z$ , for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that if  $|\omega - z| < \delta$ , then  $|f(\omega) - f(z)| < \varepsilon$ . So if  $|h| < \delta$ , then

$$\left| \frac{F(z+h) - F(z)}{h} - f(z) \right| \leq \int_0^1 |f((1-t)z + t(z+h)) - f(z)| dt < \varepsilon$$

Thus  $F' = f$  and  $f$  has a primitive.  $\square$

Later, we will see that holomorphic functions can all be represented by power series, which makes the above trivial by integrating term-by-term.

We now have the tools necessary to prove Cauchy's Theorem on a disk.

#### Corollary 2.3: Cauchy's Theorem on a Disk

If  $f : \mathbb{D}_r(z_0) \rightarrow \mathbb{C}$  is holomorphic then for any closed curve  $\gamma \subseteq \mathbb{D}_r(z_0)$ ,

$$\int_{\gamma} f(z) dz = 0$$

*Proof.* By Theorem 2.2,  $f$  has a primitive, so the integral is zero.  $\square$

#### Corollary 2.4

If  $\Omega \subseteq \mathbb{C}$  is open and contains  $\overline{\mathbb{D}_r(z_0)}$ , if  $f : \Omega \rightarrow \mathbb{C}$  is holomorphic, then

$$\int_{\partial \mathbb{D}_r(z_0)} f(z) dz = 0$$

*Proof.* It suffices to pick  $r'$  such that  $\mathbb{D}_{r'}(z_0) \supseteq \overline{\mathbb{D}_r(z_0)}$ . To show that such an  $r'$  exists, suppose that it does not. Then for any  $n \in \mathbb{N}$  there exists  $z_n \in \mathbb{D}_{r+\frac{1}{n}}(z_0)$  such that  $z_n \in \mathbb{C} \setminus \Omega$ . So  $\{z_n\}$  has a convergent subsequence  $\{z_{n_k}\}$ , which converges to a limit  $z \notin \Omega$  since  $\mathbb{C} \setminus \Omega$  is closed. But  $|z_n - z_0| < r + \frac{1}{n}$ , which means that  $|z - z_0| \leq r$ . Thus  $z \in \overline{\mathbb{D}_r(z_0)}$ , contradiction.  $\square$

## 2.2 Homotopies and Simply Connected Domains

We will now take a short detour into some topology in order to investigate which other sets Cauchy's Theorem applies to. For some intuition, consider an open set  $\Omega$  and two curves in it which share endpoints. Then if  $\Omega$  is particularly nice, we should be able to continuously deform one curve into the other.




Figure 2.4: Continuous deformation of curves

We formalize this notion by considering a continuous family of curves which deform  $\gamma_0$  into  $\gamma_1$ .

### Definition 2.2

Let  $\Omega \subseteq \mathbb{C}$  be open,  $\alpha, \beta \in \Omega$ , and  $\gamma_0, \gamma_1 : [a, b] \rightarrow \Omega$  be two curves with the same endpoints, such that  $\gamma_1(a) = \gamma_0(a) = \alpha$  and  $\gamma_1(b) = \gamma_0(b) = \beta$ . We say that  $\gamma_1, \gamma_0$  are **homotopic** in  $\Omega$  if for every  $s \in [0, 1]$  there is a curve  $\gamma_s : [a, b] \rightarrow \Omega$  with  $\gamma_s(a) = \alpha, \gamma_s(b) = \beta$  such that the function  $H : [0, 1] \times [a, b] \rightarrow \Omega$  defined by

$$H(s, t) = \gamma_s(t)$$

is continuous as a function of two variables.

### Theorem 2.5

Let  $\Omega \subseteq \mathbb{C}$  be open, and let  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic on  $\Omega$ . If  $\gamma_0, \gamma_1 : [a, b] \rightarrow \Omega$  are curves with the same endpoints that are homotopic in  $\Omega$ , then

$$\int_{\gamma_0} f(z) dz = \int_{\gamma_1} f(z) dz$$

*Proof.* We claim that there exists  $\varepsilon > 0$  such that for any  $s \in [0, 1]$  and  $t \in [a, b]$ , then

$$\mathbb{D}_{2\varepsilon}(\gamma_s(t)) \subseteq \Omega$$

Suppose not. Then for every  $n \in \mathbb{N}$ , there are  $s_n \in [0, 1]$  and  $t_n \in [a, b]$  and  $\omega_n \in \mathbb{C} \setminus \Omega$  such that

$$\omega_n \in \mathbb{D}_{2/n}(\gamma_{s_n}(t_n))$$

Now, the sequence  $(s_n, t_n) \in [0, 1] \times [a, b]$  is a sequence in a compact set, so there is a subsequence  $\{(s_{n_k}, t_{n_k})\}$  tending to  $(s, t)$ . But then

$$\left| \omega_{n_k} - \gamma_{s_{n_k}}(t_{n_k}) \right| < \frac{2}{n_k}$$

which tends to 0. Since  $\mathbb{C} \setminus \Omega$  is closed,  $\lim \omega_{n_k} \in \mathbb{C} \setminus \Omega$ . But  $\omega_{n_k} \rightarrow \gamma_s(t)$  is the limit of  $H(s_{n_k}, t_{n_k})$ , which is continuous, and thus  $\gamma_s(t) \in \Omega$ , contradiction. So the claim is proved and such an  $\varepsilon$  exists.


Now, note that  $H(s, t)$  is continuous on the compact set  $[0, 1] \times [a, b]$ , so it is also uniformly continuous. So for  $\varepsilon > 0$  which is produced by the claim, there exists  $\delta > 0$  such that if  $|s_1 - s_2| < \delta$  and  $|t_1 - t_2| < \delta$ , then

$$|\gamma_{s_1}(t_1) - \gamma_{s_2}(t_2)| < \varepsilon$$

Then subdivide  $[0, 1] \times [a, b]$  using  $0 = s_0 < s_1 < \dots < s_n = 1$  and  $a = t_0 < \dots < t_n = b$  such that  $|s_{j+1} - s_j| < \delta$  and  $|t_{j+1} - t_j| < \delta$  for all  $j$ . We claim that

$$\int_{\gamma_{s_j}} f(z) dz = \int_{\gamma_{s_{j+1}}} f(z) dz$$

for all  $j$ . Clearly this suffices to prove the theorem. Consider some pair of points  $(s_j, t_k)$  and  $(s_{j+1}, t_{k+1})$ . Draw a circle of radius  $2\varepsilon$  around  $\gamma_{s_j}(t_k)$ . Note that this circle also contains  $\gamma_{s_{j+1}}(t_k), \gamma_{s_{j+1}}(t_{k+1}), \gamma_{s_j}(t_{k+1})$ .



By Cauchy's Theorem

$$\int_{\gamma_{s_j}([t_k, t_{k+1}])} f + \int_{[\gamma_{s_j}(t_{k+1}), \gamma_{s_{j+1}}(t_{k+1})]} f - \int_{\gamma_{s_{j+1}}([t_k, t_{k+1}])} f - \int_{[\gamma_{s_j}(t_k), \gamma_{s_{j+1}}(t_k)]} f = 0$$

Thus

$$\int_{\gamma_{s_j}([t_k, t_{k+1}])} f = \int_{\gamma_{s_{j+1}}([t_k, t_{k+1}])} f + \left( \int_{[\gamma_{s_j}(t_k), \gamma_{s_{j+1}}(t_k)]} f - \int_{[\gamma_{s_j}(t_{k+1}), \gamma_{s_{j+1}}(t_{k+1})]} f \right)$$



Let us write

$$a_k = \int_{\gamma_{s_j}(t_k), \gamma_{s_{j+1}}(t_k)} f$$

Then summing over all  $k$ , we see that

$$\int_{\gamma_{s_j}} f = \int_{\gamma_{s_{j+1}}} f + a_0 - a_n$$

But  $\gamma_{s_j}, \gamma_{s_{j+1}}$  have the same endpoints so  $a_0 = a_n = 0$ . Thus the theorem is proved.  $\square$

### Definition 2.3

An open connected set  $\Omega \subseteq \mathbb{C}$  is called **simply connected** if any curves  $\gamma_0, \gamma_1 : [a, b] \rightarrow \Omega$  which share endpoints are homotopic in  $\Omega$ .

### Example 2.1

Any convex set, which is a set such that the line between two points in the set is contained in the set, such as the disk, is simply connected.

### Example 2.2

A "star-shaped" set, which is a set with a center point such that any two points can be joined by a curve passing through the center, is simply connected.

### Example 2.3

$\mathbb{C} \setminus \{0\}$  is not simply connected. Consider the upper and lower halves of  $\partial\mathbb{D}$ . Intuitively, we see that one cannot be transformed to the other without passing through the origin. We will prove this later.

### Theorem 2.6

Let  $\Omega \subseteq \mathbb{C}$  be simply connected and  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic. Then  $f$  has a primitive on  $\Omega$ .

*Proof.* Let  $\Omega$  be simply connected and pick a point  $z_0 \in \Omega$ . For any  $z \in \Omega$ , choose some curve  $\gamma_z \subseteq \Omega$  connecting  $z_0$  to  $z$ . For consistency, let us demand that  $\gamma_{z_0}$  is a constant curve. Then define

$$F(z) := \int_{\gamma_z} f(\omega) d\omega$$

Now, let  $h$  be small and let  $\varphi$  be the line segment connecting  $z$  to  $z + h$ . Then the curve  $\gamma_z + \varphi$  (where  $+$  means concatenation) shares endpoints with  $\gamma_{z+h}$ , so they are homotopic. By Theorem 2.5, integrating over either gives the same value, so

$$\int_{\gamma_{z+h}} f(\omega) d\omega = \int_{\gamma_z} f(\omega) d\omega + \int_{\varphi} f(\omega) d\omega$$

We can then write

$$F(z+h) = F(z) + \int_{\varphi} f(\omega) d\omega$$

EXERCISE: conclude the proof similarly to the proof of local existence of primitives.  $\square$

A corollary to this is a more general form of Cauchy's Theorem.

### Corollary 2.7: Cauchy's Theorem

If  $\Omega$  is simply connected and  $f : \Omega \rightarrow \mathbb{C}$  is holomorphic, then

$$\int_{\gamma} f(z) dz = 0$$

for every closed curve  $\gamma \subseteq \Omega$ .

*Proof.*  $f$  has a primitive, so the integral is zero.  $\square$

### Corollary 2.8

$\mathbb{C} \setminus \{0\}$  is not simply connected.

*Proof.* We showed that  $\int_{\partial\mathbb{D}} \frac{1}{z} dz = 2\pi i \neq 0$ .  $\square$

In fact, there are stronger versions of this theorem. We will not formally prove them here, but we will give an intuitive explanation.

### Definition 2.4

A curve  $\gamma$  is **simple** if, given some parameterization  $z : [a, b] \rightarrow \mathbb{C}$ ,  $z(s) = z(t) \implies s = t$ . A **simple closed** curve is the same, except that  $z(a) = z(b)$ .

Note that a simple closed curve is not technically simple, but the intuitive idea is the same.

### Theorem: Jordan Curve Theorem

Let  $\gamma \subseteq \mathbb{C}$  be a simple closed curve. Then  $\mathbb{C} \setminus \gamma = \Omega \cup U$ , where  $\Omega, U$  are open, connected, and disjoint. Moreover,  $\Omega$  is bounded and simply connected, and  $U$  is unbounded and connected. In this case,  $\Omega$  is called the **interior** of  $\gamma$  (denoted  $\text{int } \gamma$ ), and  $U$  is called the **exterior** (denoted  $\text{ext } \gamma$ ).

*Proof.* The proof of this theorem is omitted. Note that it is true in the piecewise smooth case (see Stein & Shakarchi appendix), but it is also true for continuous curves.  $\square$

### Theorem: General Cauchy's Theorem

Let  $\Omega \subseteq \mathbb{C}$  be open and let  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic on  $\Omega$ . Let  $\gamma \subseteq \mathbb{C}$  be a simple closed curve in  $\Omega$  such that  $\text{int } \gamma \subseteq \Omega$ . Then

$$\int_{\gamma} f(z) dz = 0$$

*Proof.* Draw a curve inside of  $\text{int } \gamma$  which is a slight perturbation of  $\gamma$  and is homotopic to  $\gamma$ . Then this follows since  $\text{int } \gamma$  is simply connected. See Stein & Shakarchi appendix for a complete proof.  $\square$

## 2.3 Consequences of Cauchy's Theorem

Aside from being a powerful theorem about complex integrals, Cauchy's Theorem also allows us to solve many difficult real integrals using complex integrals.

### Example 2.4

Consider the integral

$$\int_0^{\infty} \frac{1 - \cos x}{x^2} dx$$

Note that this is integrable since

$$\lim_{x \rightarrow \infty} \left| \frac{1 - \cos x}{x^2} \right| \leq \lim_{x \rightarrow \infty} \frac{2}{x^2}$$

and near 0, this behaves like 0. So the integral is indeed a real number.

Now, consider the function  $f : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$  defined by

$$f(z) = \frac{1 - e^{iz}}{z^2}$$

which is holomorphic. For each  $R > 0, \varepsilon > 0$ , consider the integral of  $f(z)$  over  $\Gamma_{R,\varepsilon}$ , which is defined below as  $\gamma_R + [-R, -\varepsilon] + \gamma_{\varepsilon} + [\varepsilon, R]$ :




Figure 2.5: Indented semicircle

This function is holomorphic on the star-shaped domain  $\mathbb{C} \setminus \{it : t \leq 0\}$  (which is simply connected), so

$$\begin{aligned} 0 &= \int_{\Gamma_{R,\varepsilon}} f(z) dz \\ &= \underbrace{\int_{-R}^{-\varepsilon} \frac{1-e^{-ix}}{x^2} dx}_{\text{I}} + \underbrace{\int_{\gamma_\varepsilon} \frac{1-e^{iz}}{z^2} dz}_{\text{II}} + \underbrace{\int_\varepsilon^R \frac{1-e^{ix}}{x^2} dx}_{\text{III}} + \underbrace{\int_{\gamma_R} \frac{1-e^{iz}}{z^2} dz}_{\text{IV}} \end{aligned}$$

Now,

$$\text{I} + \text{III} = \int_\varepsilon^R \frac{2 - (e^{ix} + e^{-ix})}{x^2} dx = 2 \int_\varepsilon^R \frac{1 - \cos x}{x^2} dx$$

So the problem reduces to calculating II and IV. We claim that

$$\lim_{\varepsilon \rightarrow 0} \int_{\gamma_\varepsilon^-} \frac{1-e^{iz}}{z^2} dz - \pi = 0$$

To see this, parameterize  $\gamma_\varepsilon^-$  as  $t \mapsto \varepsilon e^{it}$  for  $t \in [0, \pi]$ . Then

$$\begin{aligned} \int_{\gamma_\varepsilon^-} \frac{1-e^{iz}}{z^2} dz - \pi &= \int_0^\pi \frac{1-e^{i\varepsilon e^{it}}}{\varepsilon^2 e^{2it}} (\varepsilon i e^{it}) dt - \pi \\ &= \frac{i}{\varepsilon} \int_0^\pi \left( \frac{1-e^{i\varepsilon e^{it}}}{e^{it}} + i\varepsilon \right) dt \\ &= \frac{i}{\varepsilon} \int_0^\pi \frac{1-e^{i\varepsilon e^{it}} + i\varepsilon e^{it}}{e^{it}} dt \end{aligned}$$

We may write

$$e^{i\varepsilon e^{it}} = \sum_{n=0}^{\infty} \frac{(i\varepsilon e^{it})^n}{n!}$$

so

$$\left| e^{i\varepsilon e^{it}} - (1 + i\varepsilon e^{it}) \right| \leq \sum_{n=2}^{\infty} \frac{|i\varepsilon e^{it}|^n}{n!} = \sum_{n=2}^{\infty} \frac{\varepsilon^n}{n!} \leq 10\varepsilon^2$$

(10 is an overbound here). Returning to the original integral,

$$\left| \frac{i}{\varepsilon} \int_0^\pi \frac{1-e^{i\varepsilon e^{it}} + i\varepsilon e^{it}}{e^{it}} dt \right| \leq \frac{1}{\varepsilon} \pi 10\varepsilon^2 = 10\pi\varepsilon$$

which tends to 0 as  $\varepsilon \rightarrow 0$ .

Lastly, let us compute IV. We parameterize  $\gamma_R$  by  $t \mapsto R e^{it}$  on  $[0, \pi]$ , and see that

$$\left| \int_{\gamma_R} \frac{1-e^{iz}}{z^2} dz \right| = \left| \int_0^\pi \frac{1-e^{iRe^{it}}}{R^2 e^{2it}} iRe^{it} dt \right|$$

Observe that

$$\begin{aligned} \left| 1 - e^{iRe^{it}} \right| &= \left| 1 - e^{iR(\cos t + i \sin t)} \right| \\ &= \left| 1 - e^{iR \cos t} e^{-R \sin t} \right| \\ &\leq 1 + e^{R \sin t} \\ &\leq 2 \end{aligned}$$

where the last inequality follows since  $\sin t > 0$  on  $[0, \pi]$ . Returning to IV,

$$\left| \int_0^\pi \frac{1 - e^{iRe^{it}}}{R^2 e^{2it}} iRe^{it} dt \right| \leq \frac{1}{R} 2\pi$$

which tends to 0 as  $R \rightarrow \infty$ . Then

$$\int_0^\infty \frac{1 - \cos x}{x^2} dx = \frac{1}{2}(\text{II} + \text{IV}) = \frac{\pi}{2}$$

The following theorem is an important formula which is a powerful example of the local-global properties present in complex analysis.

### Theorem 2.9: Cauchy's Integral Formula

Let  $\Omega \subseteq \mathbb{C}$  be open,  $\overline{\mathbb{D}_R}(z_0) \subseteq \Omega$ , and let  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic on  $\Omega$ . Then for  $z \in \mathbb{D}_R(z_0)$ ,

$$f(z) = \frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{\zeta - z} d\zeta$$

*Proof.* Define

$$F(\zeta) = \frac{f(\zeta)}{\zeta - z}$$


Define the curve  $\gamma_{top}$  as shown in the diagram below:

Define  $\gamma_{bottom}$  similarly, using the counterclockwise orientation so that the integrals over the line segment cancel. Both are holomorphic on star-shaped regions, so their integrals are zero and

$$\begin{aligned} 0 &= \frac{1}{2\pi i} \left( \int_{\gamma_{top}} f(\zeta) d\zeta + \int_{\gamma_{bottom}} F(\zeta) d\zeta \right) \\ &= \frac{1}{2\pi i} \left( \int_{\partial \mathbb{D}_R(z_0)} \right) \frac{f(\zeta)}{\zeta - z} d\zeta - \int_{\partial \mathbb{D}_\varepsilon(z)} \frac{f(\zeta)}{\zeta - z} d\zeta \end{aligned}$$

so

$$\frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\partial \mathbb{D}_\varepsilon(z)} \frac{f(\zeta)}{\zeta - z} d\zeta$$



We want to show that the right side quantity is equal to  $f(z)$ . Calculating,

$$\frac{1}{2\pi i} \int_{\partial \mathbb{D}_\varepsilon(z)} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\partial \mathbb{D}_\varepsilon(z)} \left( \frac{f(z)}{\zeta - z} + \frac{f(\zeta) - f(z)}{\zeta - z} \right) d\zeta$$

Notice that the first term of the integral is the integral of  $\frac{1}{z}$  on a circle around 0, but translated and multiplied by  $f(z)$ . So this becomes

$$f(z) + \frac{1}{2\pi i} \int_{\partial \mathbb{D}_\varepsilon(z)} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta$$

The integrand is bounded since  $f$  is holomorphic, so it vanishes as  $\varepsilon \rightarrow 0$ . Thus

$$f(z) = \frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{\zeta - z} d\zeta \quad \square$$

We now arrive at perhaps the most important theorem in all of complex analysis. This theorem shows that holomorphic functions are also analytic. We showed that analytic functions are holomorphic; we now see that they are the same. Moreover, we see that all holomorphic functions are infinitely differentiable.

### Theorem 2.10

Let  $\Omega \subseteq \mathbb{C}$  be open, and suppose  $\overline{\mathbb{D}_R(z_0)} \subseteq \Omega$ . Suppose  $f : \Omega \rightarrow \mathbb{C}$  is holomorphic on  $\Omega$ . Then for every  $z \in \mathbb{D}_R(z_0)$ ,

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$$

where

$$a_n = \frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$$

In particular, the radius of convergence is at least  $R$ .

*Proof.* By the Cauchy Integral formula, when  $|z - z_0| < R$  we have

$$f(z) = \frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{\zeta - z} dz = \frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{\zeta - z_0} \cdot \frac{1}{1 - \frac{z - z_0}{\zeta - z_0}} d\zeta$$

Now, if  $|z - z_0| = r < R$ , then  $\left| \frac{z - z_0}{\zeta - z_0} \right| = \frac{r}{R}$  (since  $\zeta$  is on the boundary of  $\mathbb{D}_R(z_0)$ ). So we may write this as the sum of a geometric series:

$$\begin{aligned} \frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{\zeta - z_0} \cdot \frac{1}{1 - \frac{z - z_0}{\zeta - z_0}} d\zeta &= \frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{\zeta - z_0} \sum_{n=0}^{\infty} \left( \frac{z - z_0}{\zeta - z_0} \right)^n d\zeta \\ &= \frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \left( \sum_{n=0}^{\infty} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} (z - z_0)^n \right) d\zeta \end{aligned}$$

Thus the proof reduces to justifying the interchange of the sum and integral above. To prove this, for  $N \in \mathbb{N}$  we have

$$\begin{aligned} &= \frac{1}{2\pi i} \left[ \int_{\partial \mathbb{D}_R(z_0)} \left( \sum_{n=0}^{\infty} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} (z - z_0)^n \right) d\zeta - \int_{\partial \mathbb{D}_R(z_0)} \left( \sum_{n=0}^N \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} (z - z_0)^n \right) d\zeta \right] \\ &= \frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \left( \sum_{n=N+1}^{\infty} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} (z - z_0)^n \right) d\zeta \end{aligned}$$

So

$$\begin{aligned} \left| f(z) - \sum_{n=0}^N a_n (z - z_0)^n \right| &= \left| \frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \left( \sum_{n=N+1}^{\infty} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} (z - z_0)^n \right) d\zeta \right| \\ &\leq \frac{1}{2\pi} 2\pi R \sum_{n=N+1}^{\infty} \frac{r^n}{R^{n+1}} \sup_{\zeta \in \partial \mathbb{D}_R(z_0)} |f(\zeta)| \leq \left( \sup_{\zeta \in \partial \mathbb{D}_R(z_0)} |f(\zeta)| \right) \sum_{n=N+1}^{\infty} \left( \frac{r}{R} \right)^n \\ &= \left( \sup_{\zeta \in \partial \mathbb{D}_R(z_0)} |f(\zeta)| \right) \cdot \left( \frac{r}{R} \right)^{N+1} \frac{1}{1 - \frac{r}{R}} \xrightarrow{N \rightarrow \infty} 0 \quad \square \end{aligned}$$

We can use this to extend Cauchy's integral formula to higher order derivatives, still exhibiting local-global behavior:

### Theorem 2.11: Cauchy integral formulas

Let  $\Omega \subseteq \mathbb{C}$  be open, and let  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic on  $\Omega$ . Let  $\overline{\mathbb{D}_R(z_0)} \subseteq \Omega$ . Then  $f$  has  $n$  complex derivatives for every  $n \geq 0$  and for  $z \in \mathbb{D}_R(z_0)$ ,

$$f^{(n)}(z) = \frac{n!}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{(\zeta - z)^{n+1}} d\zeta$$

Note that this is equivalent to Theorem 2.9 for  $n = 1$ .

*Proof.* From Theorem 2.10, we can represent  $f$  as a power series

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$$

moreover, we know that we may differentiate the series term by term to see that

$$f^{(n)}(z) = \sum_{k=n}^{\infty} \frac{(n+k)!}{k!} a_k (z - z_0)^{k-n}$$

Evaluating at  $z_0$ , all the terms cancel out except  $k = n$ , so

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$$

Now, we use the same contours  $\gamma_{top}$ ,  $\gamma_{bottom}$  as in the original Cauchy Integral formula proof:



Defining  $F(\zeta) = \frac{f(\zeta)}{(\zeta - z)^{n+1}}$ , which is holomorphic on the simply connected domains containing  $\gamma_{top}$  and  $\gamma_{bottom}$ , we have

$$\frac{1}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{(\zeta - z)^{n+1}} d\zeta = \frac{1}{2\pi i} \int_{\partial \mathbb{D}_\varepsilon(z)} \frac{f(\zeta)}{(\zeta - z)^{n+1}} d\zeta = \frac{f^{(n)}(z)}{n!}$$

The last equality follows by our above work, since  $z$  is the center of  $\mathbb{D}_\varepsilon(z)$ . □

As an easy corollary we have:

#### Corollary 2.12: Cauchy Inequality

Let  $\Omega \subseteq \mathbb{C}$  be open,  $f : \Omega \rightarrow \mathbb{C}$  holomorphic on  $\Omega$ , and  $\overline{\mathbb{D}_R(z_0)} \subseteq \Omega$ . Then

$$|f^{(n)}(z_0)| \leq \frac{n!}{R^n} \sup_{\gamma \partial \mathbb{D}_R(z_0)} |f(\zeta)|$$



*Proof.* By the previous theorem,

$$\left| f^{(n)}(z_0) \right| = \left| \frac{n!}{2\pi i} \int_{\partial \mathbb{D}_R(z_0)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta \right| \leq \frac{n!}{2\pi i} 2\pi R \cdot \frac{1}{R^{n+1}} \sup_{\zeta \in \partial \mathbb{D}_R(z_0)} |f(\zeta)| \quad \square$$

### Theorem 2.13: Liouville's Theorem

If  $f : \mathbb{C} \rightarrow \mathbb{C}$  entire and bounded, then  $f$  is constant.

*Proof.* Let  $z_0 \in \mathbb{C}$ . Let  $|f(\zeta)| \leq M$  for all  $\zeta \in \mathbb{C}$ . Then as  $R \rightarrow \infty$  we have

$$|f'(z_0)| \leq \frac{1}{R} M \rightarrow 0$$

so  $f' \equiv 0$  and  $f$  is constant.  $\square$

As a corollary to this we may prove the Fundamental Theorem of Algebra:

### Corollary 2.14: Fundamental Theorem of Algebra

Let  $p \in \mathbb{C}[x]$  be non-constant. Then  $p$  has a zero in  $\mathbb{C}$ .

*Proof.* Suppose  $p$  has no root. Then  $\frac{1}{p}$  is entire. We write

$$\left| \frac{1}{p(z)} \right| = \frac{1}{|z|^n} \cdot \frac{1}{\left| a_n + \frac{a_{n-1}}{z} + \dots + \frac{a_0}{z^n} \right|}$$

(we may assume  $a_0 \neq 0$  and  $n \geq 1$ ). We bound the denominator from below for large  $z$  by

$$\left| a_n + \frac{a_{n-1}}{z} + \dots + \frac{a_0}{z^n} \right| \geq |a_n| - \frac{|a_{n-1}|}{|z|} - \dots - \frac{|a_0|}{|z|^n}$$

so

$$\left| \frac{1}{p(z)} \right| \leq \frac{1}{|z|^n} \cdot \frac{1}{|a_n| - \frac{|a_{n-1}|}{|z|} - \dots - \frac{|a_0|}{|z|^n}} \xrightarrow{|z| \rightarrow \infty} 0$$

Thus we pick  $R > 0$  such that  $\frac{1}{|p(z)|} \leq 1$  when  $|z| > R$ . On  $\overline{\mathbb{D}_R(z_0)}$ ,  $\frac{1}{p(z)}$  is continuous and thus bounded. So  $\frac{1}{p}$  is bounded everywhere and thus constant by Liouville's Theorem. But we supposed  $p$  was not constant, so  $p$  must have a root.  $\square$

We now provide a proof of the method known as analytic continuation.

### Definition 2.5

Let  $\Omega \subseteq \mathbb{C}$  be open and let  $\Omega' \subseteq \Omega$  be open. Let  $f : \Omega' \rightarrow \mathbb{C}$  and  $F : \Omega \rightarrow \mathbb{C}$  be holomorphic, with  $F|_{\Omega'} = f$ . Then we call  $F$  is an **analytic continuation** of  $f$ .

The key result is that, under minor restrictions, this choice of  $F$  is actually unique; that is, the value of  $f$  on a small set completely determine its extension to  $\Omega$ .

### Theorem 2.15: Uniqueness of Analytic Continuation

Let  $\Omega \subseteq \mathbb{C}$  be open and connected, and consider some collection of distinct points  $\{\omega_n\}_{n=1}^{\infty} \subseteq \Omega$  such that  $\lim \omega_n = \omega \in \Omega$  exists. Suppose that  $f : \Omega \rightarrow \mathbb{C}$  is holomorphic and the  $\omega_i$  are zeroes of  $f$ . Then  $f \equiv 0$  on  $\Omega$ .

*Proof.* Let  $A = \{z \in \Omega : f(z) = 0\}$ . Let  $U = \text{int } A$ . I claim the following:

**Claim 1:**  $U$  is nonempty.

**Claim 2:**  $U$  is closed.

The theorem follows from the above, as if they are true, we have  $\Omega = U \cup (\Omega \setminus U)$ . By Claim 2, these are disjoint open sets, so one must be nonempty. By Claim 1, it must be  $\Omega \setminus U$  so  $U = \Omega$ .

To prove these claims, we show the following:

#### Claim

If  $\{\zeta_n\} \subseteq A$  are distinct and  $\zeta_n \rightarrow \zeta_0 \in \Omega$ , then there is  $R > 0$  such that  $f$  vanishes on  $\mathbb{D}_R(\zeta_0)$ .

*Proof.*  $\Omega$  is open and  $\zeta_0 \in \Omega$ , so there exists  $R > 0$  such that  $\overline{\mathbb{D}_R(\zeta_0)} \subseteq \Omega$ . Since  $f$  is holomorphic, it is analytic, so we may write

$$f(\zeta) = \sum_{n=0}^{\infty} a_n(\zeta - \zeta_0)^n$$

on  $\mathbb{D}_R(\zeta_0)$ . For large enough  $n$ ,  $\zeta_n \in \mathbb{D}_R(\zeta_0)$ , and thus

$$f(\zeta_0) = \lim f(\zeta_n) = 0$$

so  $a_n = 0$ . If all the  $a_i$  are zero, then we are done. Suppose not. Then let  $m \geq 1$  be the smallest index such that  $a_m \neq 0$ . Then

$$f(\zeta) = a_m(\zeta - \zeta_0)^m \left( 1 + \underbrace{\frac{a_{m+1}}{a_m}(\zeta - \zeta_0) + \frac{a_{m+2}}{a_m}(\zeta - \zeta_0)^2 + \dots}_{=g(\zeta)} \right)$$

Now,  $g(\zeta)$  is analytic on  $\mathbb{D}_R(\zeta_0)$  as it is a power series, and  $\zeta_0 = 0$ . For large enough  $n$  we then have

$$f(\zeta_n) = a_m(\zeta_n - \zeta_0)^m(1 + g(\zeta_n))$$

By assumption,  $f(\zeta_n) = 0$ . But  $a_m \neq 0$  by assumption.  $(\zeta_n - \zeta_0)$  is only zero for at most one  $\zeta_n$ , so we conclude that  $1 + g(\zeta_n) = 0$  for all large enough  $n$ . As  $n \rightarrow \infty$ ,  $g(\zeta_n) \rightarrow 0$ , so  $1 = 0$ , contradiction. Thus no  $a_m$  is nonzero and  $f$  is identically zero.  $\blacksquare$

Now we apply the above to our  $\{\omega_n\}$  and arrive at Claim 1. For Claim 2, if there is some sequence of points in  $U$ , then the Claim also shows that their limit is in  $U$ . So we are done.  $\square$

### Corollary 2.16

Let  $f : \Omega' \rightarrow \mathbb{C}$  be holomorphic with  $\Omega' \subseteq \Omega \subseteq \mathbb{C}$  both open. If  $\Omega$  is connected, then there is at most one analytic continuation  $F : \Omega \rightarrow \mathbb{C}$  of  $f$ .

*Proof.* Suppose  $F_1, F_2 : \Omega \rightarrow \mathbb{C}$  are analytic continuations of  $f$ . Then  $F_1 - F_2$  is zero on  $\Omega'$  open, and we immediately conclude that  $F_1 - F_2 \equiv 0$  on  $\Omega$ . So  $F_1 = F_2$ .  $\square$

We now prove a converse to Goursat's Theorem.

### Theorem 2.17: Morera's Theorem

Let  $f : \mathbb{D}_R(z_0) \rightarrow \mathbb{C}$  be continuous and for any triangle  $T \subseteq \mathbb{D}_R(z_0)$  we have

$$\int_{\partial T} f(z) dz = 0$$

then  $f$  is holomorphic on  $\mathbb{D}_R(z_0)$ .

*Proof.* We imitate the proof that holomorphic functions locally have primitives, which only assumed Goursat's Theorem. In this case we do not know that  $f$  is holomorphic but we assume the conclusion of Goursat's Theorem. Thus  $f$  has a primitive  $F$ . So  $F$  is holomorphic, and thus infinitely differentiable. In particular,  $f = F'$  is holomorphic.  $\square$

# Definitions

- analytic, 13, 26
- analytic continuation, 48
- argument, 3
- boundary, 8
- bounded, 8
- Cauchy sequence, 6
- Cauchy-Riemann equations, 17
- closed curve, 26
- closed disk, 7
- closed set, 7
- closure, 8
- compact, 8
- complex conjugate, 5
- complex differentiable, 13
- connected, 10
- continuous, 11
- contour integral, 27
- converges, 5
- converges absolutely, 20
- curve, 27
- diameter, 8
- disk of convergence, 21
- entire, 13
- equivalent curves, 27
- exponential, 22
- exterior, 41
- Hadamard's formula, 21
- holomorphic, 12
- homotopic, 38
- imaginary part, 3
- interior, 7, 41
- interior point, 7
- length, 28
- limit point, 7
- modulus, 3
- negative orientation, 27
- open cover, 9
- open disk, 7
- open set, 7
- parameterized curve, 26
- path, 11
- path connected, 11
- piecewise smooth, 26
- positive orientation, 27
- power series, 20
- primitive, 30
- radius of convergence, 21
- real part, 3
- region, 11
- sequence, 5
- simple, 41
- simple closed, 41
- simply connected, 40
- smooth curve, 26
- subcover, 9
- triangle, 33
- triangle inequality, 4
- trigonometric functions, 23