The art and design of genetic screens: Yeast

Article in Nature Reviews Genetics · October 2001 DOI: 10.1038/35088500 · Source: PubMed					
CITATIONS 164		READS 2,050			
1 author	rthor:				
	Susan Forsburg University of Southern California 136 PUBLICATIONS 6,793 CITATIONS SEE PROFILE				

REVIEWS

THE ART AND DESIGN OF GENETIC SCREENS: YEAST

Susan L. Forsburg

Understanding the biology of complex systems is facilitated by comparing them with simpler organisms. Budding and fission yeasts provide ideal model systems for eukaryotic cell biology. Although they differ from one another in terms of a range of features, these yeasts share powerful genetic and genomic tools. Classical yeast genetics remains an essential element in discovering and characterizing the genes that make up a eukaryotic cell.

ASCOMYCETE
Free-living fungus that
reproduces sexually through
the formation of spores
packaged in a sac called an
ascus. Some taxonomists
include non-sexually
reproducing fungi with DNA
sequences, which indicates a
close degree of relatedness.

COMPLEMENTATION TEST Determines whether two recessive mutations are in the same functional unit or gene. Two recessive mutant strains, a1 and a2, crossed together complement each other if the resulting diploid has a wildtype phenotype; as each provides the function missing in the other, they are assumed to affect independent genes. If, instead, the diploid has the mutant phenotype, then a1 and a2 do not complement and are assumed to affect the same gene.

Molecular and Cell Biology Laboratory, The Salk Institute, 10010 N. Torrey Pines Road, La Jolla, California 92037, USA. email: forsburg@salk.edu Yeast geneticists half-mockingly talk about the cult of APYG: the 'awesome power of yeast genetics'. But mocking aside, these simple, single-celled fungi have proven themselves to be the workhorses of cell biology because of the ease of their genetic manipulation. The budding yeast *Saccharomyces cerevisiae* and the fission yeast *Schizosaccharomyces pombe* are quite different in their biology (FIG. 1), but they share a similar tool set that makes the process of gene discovery, and subsequent characterization of gene function, remarkably easy.

Historically, *S. cerevisiae* has been the more popular experimental system. The first eukaryote to be transformed by plasmids, it was also the first eukaryote for which precise gene knockouts were constructed, and the first to have its genome sequenced¹⁻³. The cell biological issues that have been explored in this ASCOMYCETE range from signal transduction to cell-cycle control, chromosome structure to secretion. The identified genes have been used as probes to uncover further pathways and to identify metazoan homologues. Despite the completion of its genome sequence several years ago¹, the roles of many of its 6,000+ genes remain unclear. The process of mutant analysis and discovery of gene function continues with the added tools of genomics^{4,5}.

By contrast, the experimental history of *S. pombe* involves a smaller, but growing, community. Its genome sequence is essentially complete but shares no conserved synteny (gene order) with the budding yeast in its 4,900+ genes⁶. Although yeast phylogeny is still unclear, *S. pombe* is thus quite distinct from *S. cerevisiae* and filamentous fungi^{7–9}. The fission yeast

has a symmetrical pattern of cell division and has been particularly popular for studies of cell growth and division, and chromosome dynamics. Other cell biological questions have been addressed more recently, inspired by the power of a comparative approach between these two superficially similar organisms. Each of them offers unique insights as a model organism for elucidating the biology of more complex cell types¹⁰.

Both yeasts are adaptable to several forms of genetic analysis that allow the identification of new genes, or the functional analysis of previously identified genes^{11,12}. Because both can grow and divide as haploids, recessive mutations are easily recovered. However, a diploid sexual cycle exists for both, allowing facile genetic analysis, including tests of COMPLEMENTATION, RECOMBINATION and EPISTASIS. The identification of replication origin sequences allows plasmids to be




Figure 1 | **Morphology of the two yeasts.** The fission yeast *Schizosaccharomyces pombe* and the budding yeast *Saccharomyces cerevisiae* are free-living haploid cells that are easily grown in the laboratory. Left, fission yeast; right, budding yeast; visualized with the DNA stain DAPI (4',6-diamidino-2-phenylindole) to highlight the nucleus.

Table 1 | Nomenclature in the two yeast species

	S. cerevisiae	S. pombe
Wild-type gene	YFG1	yfg1+
Deletion (null) mutant	Δ yfg1 yfg1 Δ	Δ yfg1 yfg1 Δ
Recessive mutant	yfg1-1	yfg1-1
Dominant mutant	YFG1-2	yfg1-2
Protein	Yfg1 YFG1p	Yfg1 yfg1p

Yfg typically means 'your favourite gene'. The 'p' designation for proteins (for example, Yfg1p) is occasionally used. S. cerevisiae, Saccharomyces cerevisiae or budding yeast; S. pombe, Schizosaccharomyces pombe or fission yeast.

RECOMBINATION Any process in a diploid or partially diploid cell that generates new gene or chromosomal combinations not found in that cell or in its progenitors. At meiosis, recombination (or crossing over) is the process of reciprocal exchange between homologous chromosomal segments that generates a haploid product genotypically distinct from the two haploid genotypes of the original meiotic diploid.

EPISTASIS

The phenotype caused by a mutation in one gene is masked by a mutation in another gene. Epistatic analysis requires that two mutants have distinguishable phenotypes. It can be used to determine the order of gene function by testing whether the phenotype of the double mutant ab is similar to that of mutant a, or mutant h

HETEROLOGOUS RECOMBINATION Recombination between DNA molecules with significantly different sequences, for example when a transgenic construct integrates randomly in the

maintained as free episomes, and these are easily introduced into the cell by transformation. In addition, both species have high rates of homologous recombination, allowing precise manipulation of the genome for the construction of gene disruptions and allele-specific replacements. Their nomenclature is distinct (TABLE 1), but they benefit from similar genetic and molecular tools (see examples in TABLE 2), which provide the basic requirements for the genetic screens described below.

With the genome sequences of both species now completed, the challenge is to identify the functions associated with their genes. The classical genetic analysis described here complements the newest genomic approaches13: genetics moves from a function, defined by mutation, to identify the gene responsible, whereas genomics moves from the catalogue of genes to identify their function. The true power of genetics is its predictive value; genetic interactions predict physical interactions, and these can be tested using standard molecular and biochemical techniques. Genetics is, therefore, one of a trio of methods, the others being molecular biology/biochemistry and cell biology, which are required to understand the function of individual genes in vivo. In this review, I discuss principles and examples of genetic screening methods in both yeasts that facilitate functional gene discovery, and that continue to provide crucial insights into the mechanisms that make up a single cell.

Table 2 | Corresponding tools in the two yeast species

	S. cerevisiae	S. pombe
Regulated promoter	GAL (galactose regulated)	nmt (thiamine regulated)
Plasmid replication origins	ARS1 or 2µ	ars1
Auxotrophic markers		
Uracil, orotidine 5'-phosphate decarboxylase Select against with 5-FOA	URA3	ura4+
Leucine, β-isopropylmalate dehydrogenase	LEU2	leu1+
Adenine, phosphoribosyl- aminoimidazole carboxylase Accumulates red colour	ADE2	ade6+

 2μ (2 micron), an endogenous plasmid DNA molecule found in some yeast cells, with a circumference of 2µ; 5-FOA, 5'-fluoro-orotic acid. S. cerevisiae, Saccharomyces cerevisiae or budding yeast; S. pombe, Schizosaccharomyces pombe or fission yeast

Classical genetics and mutant isolation

The essence of any genetic screen or selection is to identify a process of interest, to predict the likely phenotype of a mutant unable to carry out that process, and to devise a method of identifying mutants with that phenotype. The precise methods used vary according to the needs of the investigator and the limitations associated with each approach, but the broad goals remain the same across different screens.

A particular advantage of the yeasts as genetic models is their haploid life cycle. This facilitates the recovery of recessive mutations; these are usually loss-of-function alleles and are particularly useful in determining the normal function of the gene. Null mutations that result in a complete loss of gene function can be generated in several ways. Specific genes can be disrupted using a construct that contains a nutritional or drug marker and homologous sequences. If no gene-specific sequences are used, the construct can integrate randomly throughout the genome. For example, fission yeast has recently been subject to random insertional mutagenesis 14,15 — a technique that is assisted by its relative promiscuity in HET-EROLOGOUS RECOMBINATION¹⁶. The advantage is the ease of identifying the gene of interest, which is tagged by the inserted cassette, and the knowledge that the gene function is abolished. The disadvantage is that essential genes cannot be examined, because the mutated cells will be inviable; so, the approach is useful for any non-essential cell activity (for example, meiosis) or redundant gene, or for gene activity that is sensitive to dosage.

However, even lethal mutations can be maintained if the wild-type copy is also present and this is accomplished using diploid heterozygotes. A recent development in the budding yeast community is the use of a pool of strains in which all the open reading frames have already been knocked out (see link to the Saccharomyces genome deletion project) or fused to a transposon (see link to TRIPLES — a database of transposon-insertion phenotypes, localization and expression in Saccharomyces). These collections therefore contain mutations in essential as well as nonessential genes. But there is an additional twist — not all viable heterozygotes in this collection are normal. Several such heterozygous diploids have abnormal chromosome content, presumably as a result of internal selection¹⁷. This is presumably because budding yeast often has several versions of the same gene¹⁸, reflecting ancient genome duplication events; loss of one gene might impose a growth selection that leads to increased copy number of the chromosome that contains another version of the same gene, because it confers increased growth rate. (This leads to the important caveat that any time an investigator grows a yeast strain in the laboratory, the cells are under some form of selection!) Nevertheless, these heterozygous strains can be turned to an advantage and the pool of diploid disruptions can be used for novel screens. For example, the reduced dosage associated with a heterozygous null can generate useful phenotypes owing to HAPLOINSUFFICIENCY, and this has been successfully used in screens for drug sensitivity¹⁹.




Figure 2 | Examples of mutant phenotypes in studies of cell-cycle and chromosome dynamics. a-c, e-g | Saccharomyces pombe; d | Saccharomyces cerevisiae. a | Abnormal mitosis in fission yeast. The top two wild-type cells show a normal mitotic spindle (anti-tubulin, red) and proper segregating anaphase nuclei (DAPI, which stains DNA, blue). The bottom two mutant cells show a fragmentation phenotype of irregularly segregating DNA along the spindle. **b** | Cells entering meiosis from the haploid state in a *pat1* mutant. DAPI is false-coloured green. Note the three or four irregularly segregating nuclei per cell body, after a lethal attempt to go through MI and MII divisions. c | cell division cycle 2 (cdc2, which encodes a cyclindependent kinase (CDK)) mutants elongate without dividing. Composite image of differential interference contrast (DIC) and fluorescence (DAPI). d | cdc28 (CDK) mutants arrest without budding. e-f | Normal yeast cells (smaller cells) are shown next to re-replication mutants in yeast that overexpress cdc18. Mutants undergo several rounds of DNA replication in the absence of mitosis (e, phase contrast; f, propidium iodide-stained DNA in red). g | Cells deleted for tip1, with mispositioned growth sites. h | Overexpression of an unknown gene that affects cytokinesis, as well as the size, number and positioning of growth sites. (Panels e-f courtesy of Hideo Nishitani, Japan; panels **g-h** courtesy of Damian Brunner, EMBL, Germany.)

However, the difficulty of working with lethal, lossof-function mutations in essential genes leads investigators to seek more modest point mutations that change, but do not abolish, gene function. A broad spectrum of these mutations can be generated using different chemical mutagens or irradiation. Geneticists are particularly fond of conditional mutants, in which the encoded gene product is functional under one condition, but not under another. A classic condition is temperature. Temperature- or thermosensitive (ts) mutant proteins function at low temperature, but are inactivated or unfolded at high temperature. By contrast, cold-sensitive (cs) mutant proteins are presumed to affect protein-protein interaction surfaces at the low (restrictive) temperature. In these screens, cells are propagated at the permissive temperature and are then screened at the restrictive temperature for the phenotype of interest. These are identified by REPLICA PLATING, and looking for colonies that can grow under permissive conditions but not under restrictive ones. In fission yeast, the difference can be enhanced using phloxin B, a vital dye that stains sick colonies dark pink or red. The cells are then examined further for the desired phenotype, which could be a particular cell or colony morphology, or nuclear or spindle defect. Some examples are shown in FIG. 2.

For example, cells that have ts mutations in the celldivision-cycle machinery can grow but cannot divide at the restrictive temperature, leading to mutant cells with a distinct morphology, which indicates an arrest at a particular point in the cell cycle (cell division cycle, or cdc, mutants). In fission yeast, cdc mutants are typically elongated; in budding yeast, they have a distinct bud morphology that corresponds to their position in the cell cycle. These characteristics were exploited in a series of classic papers by Lee Hartwell in budding yeast²⁰, followed by Paul Nurse in fission yeast21, and many of the cdc genes isolated were subsequently shown to be conserved in all eukaryotes (reviewed in REF. 22). Another condition might be imposed using specific drugs. For example, drugs that disrupt normal cell-cycle events, such as hydroxyurea (which blocks DNA replication) or benomyl (which disrupts microtubule function), provide other ways of challenging normal cell-cycle dynamics in a mutant cell. Some mutants that are defective in the DNA replication checkpoint in fission yeast were identified because of their inability to form colonies on plates that contain low doses of hydroxyurea when wildtype cells were able to grow²³. The first isolation of spindle-checkpoint mutants in budding yeast relied on a change in cell morphology in response to drug treatment: cells that arrest the cell cycle normally in response to benomyl remain as single cells, whereas mutants with defective checkpoints attempt to divide, forming a minicolony of dying cells^{24,25}. Finally, different nutrient media can also provide conditional selection. For example, strains that are deficient in galactose metabolism will still be able to grow on a medium that contains glucose.

It is important to note that a conditional mutant is mutant at all times, not simply under restrictive conditions. Under permissive conditions, the activity of HAPLOINSUFFICIENCY A gene dosage effect that occurs when a diploid requires both functional copies of a gene for a wild-type phenotype. An organism that is heterozygous for a haploinsufficient locus does not have a wild-type phenotype.

the mutant gene might be near to wild type, or it might be attenuated sufficiently to observe a phenotype such as chromosome loss. Mutants with attenuated activities under permissive conditions can be particularly useful in synthetic enhancement screens (see below). The phenotype of viable cells with partial activity can also be exploited in direct genetic screens. For example, cells with defects in essential genes

required for chromosome maintenance or transmission are often identified by assays that monitor relatively minor defects in plasmid stability under permissive conditions. A useful tool now in wide use is a colour marker that allows colonies containing the plasmid to be scored by the naked eye^{26,27} (BOX 1).

The use of selection, as opposed to screening, is particularly powerful. Whereas a screen requires a strategy to examine large numbers of mutants and then find among them the few with the desired phenotype, a selection establishes conditions in which only the mutants of interest survive. If one considers the difference between screening for mutants that are sensitive to a drug such as hydroxyurea, versus selecting for mutants that are resistant to the drug, it is easy to see that in the latter case only the mutants of interest will be recovered, saving the investigator significant effort.

Screens for dominant, or gain-of-function, mutations are also quite powerful. Sometimes simply increasing the dosage of a gene can result in a mutant phenotype²⁸. Overexpression screens are typically carried out using a HIGH-COPY LIBRARY, in which a cDNA is expressed under a controllable promoter, but can also be carried out using a genomic library, by relying on the increased copy number of the EPISOME relative to the chromosome. The particular advantage of these dosage-sensitivity screens is that the gene of interest is essentially cloned already, and can be identified simply by isolating the plasmid from the yeast cell, partially sequencing it and comparing it to genome databases. The phenotypes screened or selected for by overexpression studies are as varied as those used for loss-of-function screening. In a recent example, a screen of budding yeast genes that confer a cell-cyclerelated phenotype when overproduced, such as arrest at a particular point of the division cycle, identified several new genes with no previous link to cell-cycle progression²⁹. This method can also be used to analyse the function of a previously cloned gene, by mutagenizing it in vitro and screening for variants that cause a dominant-negative phenotype³⁰.


Screening for mutants is a statistical effort. Numerous labs can repeat the same screen and come up with a different spectrum of mutants for each iteration. Generally, a screen is said to approach saturation when mutations in the same genes are isolated repeatedly. However, variations of methods, mutagens, plasmid libraries and even investigators continue to provide fruitful outcomes in these analyses.

Defining genetic networks

In well-studied systems such as the yeasts, there are few processes that have not yet been targeted for genetic analysis, so for many processes, some genes have already been identified. Mutants in these provide a way to bootstrap through a genetic network and identify other genes in the same pathway, or in different pathways that affect the process of interest. These methods rely on synthetic interactions between different mutations, and are perhaps the most potent weapons in the current genetics arsenal31,32.

Box 1 | Plasmid dynamics

Plasmid screening Cells that contain a plasmid can be screened using colour markers^{12,26,27}. A yeast cell that lacks the Saccharomyces cerevisiae ADE2 (or Schizosaccharomyces pombe ade6+) gene accumulates a red intermediate in the adenine biosynthesis pathway (see the illustration). If the wildtype copy of the corresponding ade gene is



placed on a non-essential plasmid or a minichromosome, it complements the chromosomal ade mutation and prevents accumulation of the red compound. The colonies that contain the wild-type ade gene will therefore be white or pale pink, whereas cells that lose the plasmid will form dark red colonies. Plasmid loss can be monitored by determining whether white colonies produce red papillae or red sectors, and whether individual cells produce red, white, or sectored colonies (see illustration). An increased rate of sectoring can indicate defects in plasmid maintenance. Alternatively, cells will form white or pale pink colonies without sectors if they acquire some other mutation that requires the plasmid for viability. A variation on the theme uses a tRNA suppressor encoded by the plasmid and a nonsense allele in the ade gene. The tRNA suppressor incorporates an amino acid at the ade nonsense codon. In this case, the degree of colour is an indicator of plasmid dosage: higher copy number or more stable forms of the plasmid result in white cells, low copy number or unstable plasmids result in pink cells and loss of the plasmid results in red colonies.

Plasmid selection


Cells that contain a plasmid are typically selected using nutritional markers; for example, by putting a LEU2 gene required for leucine biosynthesis on a plasmid and transforming it into a leu2 mutant yeast strain (see TABLE 2 for some commonly used markers). However, some markers can be used for negative (counter-) selection, to select cells that have lost the plasmid. For example, the S. cerevisiae CAN1 (S. pombe can1+) gene, which encodes an arginine/histidine permease, renders cells sensitive to the toxic arginine analogue, canavanine. Mutant cells are resistant to the drug, because they cannot import it. Therefore, can1- cells that have lost a plasmid containing the CAN1 marker will be able to grow on canavanine-containing plates (canavanine resistant), whereas cells that contain the plasmid will be canavanine sensitive. The most useful marker of all is the S. cerevisiae URA3 gene (S. pombe ura4+), which has both positive and negative selection properties. S. cerevisiae ura3 or S. pombe ura4 mutants require the gene to grow in the absence of uracil, but when the gene is present, they become sensitive to the toxic analogue 5-FOA (5'-fluoro-orotic acid). So, cells transformed with plasmids that contain the wild-type genes and a ura marker can be isolated by selecting for growth without uracil, and plasmid-free cells can be recovered by growth in 5-FOA. The S. cerevisiae TRP1 gene can also be used for counterselection with FAA (5'-fluoroanthranilic acid).

a Dosage suppressor: rescues in high copy


Mutant Increased dosage of wild-type partner stabilizes protein Wild type Protein is destabilized

Suppressor

b Interaction suppressor: allele specific, gene specific



c Bypass suppressor: pathway specific, rescues null allele



d Nonsense suppressor: allele specific, gene non-specific




Figure 3 | Suppressor mechanisms. Depending on the allele and gene specificity associated with suppressors, mechanisms can be inferred, as shown. a | Dosage suppressors encode proteins that stabilize the mutant product when they are expressed at high levels. **b** | An interaction suppressor restores the interaction between the mutant product and its partner(s). c | A bypass suppressor activates an alternative pathway to the wild-type pathway. d | A nonsense suppressor encodes a tRNA molecule that recognizes a premature termination codon and inserts an amino acid at that position.

REPLICA PLATING A classic method to duplicate the colonies on an agar plate by stamping them on sterile velvets or filters, and then applying these copies to new (replica) plates. The replica plates can then be used to test the colonies for growth on different nutrient media or at different temperatures.

HIGH-COPY LIBRARY Most plasmid episomes in yeast are present at greater than one copy per cell, leading to an increased dosage of any gene(s) carried by the plasmid. The high copy dosage effect can be enhanced if the cells are transformed with a library that contains cDNAs expressed by strong promoters.

Two broad approaches exist for working a way through the network. The first is to use suppression analysis: trying to rescue the phenotype of the original mutation by an additional mutation, or by increased gene dosage³¹. The second is to search for synthetic enhancement: trying to exacerbate the phenotype of the original mutation by a mutation in a second gene, or by increased gene dosage, perhaps to the point of death (synthetic lethality)32.

Suppressor analysis. Suppressor analysis is particularly attractive because it generally uses selection, rather than screening. A hypothetical mutant yfg1 (your favourite gene 1) that is unable to grow under particular conditions is subject either to mutagenesis, or to transformation with a high-copy plasmid library. Survivors are selected directly in restrictive conditions. Subsequent genetic and molecular analysis reveals whether the suppressor simply reverted the original lesion in YFG1, or whether it defines an independent locus SUP1. However, although conceptually straightforward, the practical

application of suppressor analysis is not always easy, and not all *yfg1* mutants will be tractable for suppression.

How does suppression happen? There are several mechanisms for second-site suppressors31 (FIG. 3). The most informative is the interaction suppressor. Imagine two proteins that interact physically. Mutation of one might disrupt the interaction, but a compensatory mutation of its partner will restore the association and rescue the mutant phenotype. This change is gene specific and allele specific, as not all mutations in the first gene can be rescued by the same compensatory mutation in the second, and a null mutation will not be rescued. This 'lock-and-key' model offers a strong prediction for direct physical interaction that can be readily tested by molecular or cytological assays, such as coimmunoprecipitation or localization. Using this sort of analysis, direct interactions between the fission yeast Cdc2 kinase and the Cdc13 cyclin were predicted33, subsequently allowing definition of Cdc2 as the cyclindependent kinase (CDK) of this organism. A variation of the lock-and-key interpretation was provided in studies of budding yeast actin and its associated proteins, in which suppressor mutations seemed to increase the affinity of the interacting proteins for one another by introducing additional contact surfaces³⁴.

Another helpful example is a bypass suppressor, in which an alternative pathway can be activated. For example, the Cdc7 kinase in budding yeast is required to initiate DNA replication, probably by modifying the minichromosome maintenance (MCM) protein complex. The MCM complex consists of six highly conserved proteins that are essential for the initiation and progression of DNA synthesis (reviewed in REF. 35). A specific mutation in MCM5 called bob1 bypasses the requirement for Cdc7 activity. This is a gene-specific, but allele-nonspecific suppressor, as mcm5-bob1 suppresses the lethal phenotype of the cdc7 null as well as the conditional alleles. The bob1 mutation is thought to induce a conformational change in the MCM complex that would normally require modification by the Cdc7 kinase (REF 36).

Not all second-site suppressors are relevant to the study of a particular process. For example, if a gene has a mutation that results in a termination codon, it is possible to isolate a rescuing mutation in a tRNA that now recognizes the termination codon and inserts an amino acid at that position. This informational suppressor is not specific to the original gene that was mutated, but will rescue any gene with the same termination codon. It is an allele-specific, but gene-nonspecific suppressor. These are relatively rare in most yeast screens, simply because the typical mutations used for suppressor analysis are temperature sensitive and these are less likely to result from premature termination.

The identity of the genes involved in second-site suppression must be established molecularly by cloning the corresponding genes (FIG. 4). Cloning by complementation (the standard yeast approach) requires a selectable phenotype. For this reason, suppressors are often screened for additional phenotypes; for example, one might suppress a ts mutant and screen the rescuers for a




Figure 4 | Cloning suppressors. Cloning of suppressors is significantly easier if they confer an independent phenotype (for example, cold sensitivity). However, strategies exist for cloning suppressors without a separate phenotype, as described in the text and shown here for the fission yeast Schizosaccharomyces pombe. lof, loss-of-function mutation; gof, gain-of-function mutation; yfg1, your favourite gene1. See text for details.

EPISOME

An independent DNA element, such as a plasmid, that can replicate extrachromosomally or that can be maintained by integrating into the genome of the host.

DOMINANT NEGATIVE A mutant allele that interferes with the function of its wildtype version.


SYNTHETIC INTERACTIONS These occur when a double mutant has a phenotype different from either single mutant parent. For suppressors (synthetic viable), the double mutant is viable when at least one of the single mutants is not. For synthetic lethal mutants, the double mutant is inviable under conditions in which both parents are viable.

new, cs phenotype. The suppressor gene would then be identified by transforming a plasmid library into the cs strain and complementing it for growth at the restrictive temperature (a direct selection, FIG. 4a). However, not all proteins are susceptible to cs mutations.

It is possible, although time-consuming, to clone a suppressor even if it lacks an independent phenotype. If the suppressor mutation is recessive, then it can be cloned in a screen for loss of suppression. In this strategy, the plasmid library is transformed into the suppressed strain that carries the original mutation (*yfg1 sup1*), and the transformants are screened for loss of the suppression phenotype. For example, if the original mutation yfg1 was temperature sensitive (FIG. 4b), the suppressed strain yfg1 sup1 would be temperature resistant. But restoring the wild-type copy of the suppressor gene *sup1*+ would again confer temperature sensitivity. This could be determined using replica plating — an entirely feasible, if somewhat laborious approach. Matters become more complicated if the suppressor is dominant, representing a

gain-of-function mutation. Under these circumstances, the only options are either to construct a plasmid library from the suppressor strain, transform it into the original mutant yfg1 and screen for suppression (FIG. 4c), or to genetically map the mutation, and attempt to identify the gene using the complete genome sequence and physical map. This could take months.

The potential for difficulty in cloning second-site suppressors makes high-copy, or dosage suppression, an attractive approach, because the rescuing gene is already cloned. In this strategy, the mutant strain is transformed with a high-copy library, preferably a cDNA library under a controllable promoter. The transformants are screened or selected for survivors under conditions in which the promoter is induced. One likely category of survivors will contain the wildtype version of the mutant gene under study. However, other genes might also be isolated. These dosage suppressors might result from direct interactions that stabilize the mutant protein. Alternatively, increasing the



Loss-of-function mutation in yfg1ts: requires plasmid for viability

Figure 5 | Synthetic lethal screens. Synthetic lethal screens are powerful tools for identifying additional genes in a pathway of interest. The challenge of the screen is to design starting strains that allow rapid determination of whether the new mutation occurs in a different gene, and whether it has an independent phenotype. The figure shows the search for mutations that are synthetically lethal with yfg1 (your favourite gene 1) in the budding yeast Saccharomyces cerevisiae. A strain with a chromosomal mutation in yfg1, and with ade2 and ura3 markers, forms pink colonies. It is rescued by the wild-type YFG1 that is present on a selectable plasmid marked with ADE2 and URA3, forming white colonies. The plasmid-transformed strain is mutagenized and the surviving cells are scored for those that cannot survive in the absence of the plasmid (and therefore require YFG1 for viability). These will include strains with null mutations either in the yfg1 gene itself or in an unrelated gene that is synthetically lethal with yfg1. Cells that do not require the plasmid will form pink colonies and will be discarded. 5-FOA, 5'-fluoro-orotic acid; ts, temperature sensitive.

dosage of a downstream factor might bypass the requirement for the original mutated gene. These possibilities could be distinguished genetically, by determining whether the mutation is allele specific or can rescue the null allele, or biochemically, by determining whether the proteins encoded by the genes can physically interact with one another. Suppressing plasmids are readily recovered from the transformed yeast, and the identity of the open reading frames can be rapidly determined by generating a short sequence from the clone and comparing this to the completed genome sequence databases.

One powerful version of plasmid-based suppression is the identification of functional homologues from other organisms. cDNA libraries that contain mammalian genes in yeast vectors can be used to determine whether, for example, a human gene can complement a yeast mutant. The human version of the fission yeast cdc2+ gene that encodes a CDK kinase was identified in just this way³⁷. Banks of ts yeast strains have been constructed to exploit this approach (for example, REF. 38). However, despite obvious structural conservation between many genes involved in basic cell function, successful crosscomplementation is relatively rare. It might be that precise protein interactions have changed through evolution,

or additional components of the system are more variable than the one under study. So, failure of a metazoan gene to complement a yeast mutant does not mean that they are not functional homologues.

Synthetic lethality. Synthetic lethality is the opposite of suppression³². Instead of restoring function to the original mutant, the synthetic lethal screen aims to further attenuate function, to the point of death. This strategy can be particularly useful for uncovering redundant pathways or direct protein interactions, depending on the nature of the starting mutation. The hallmark of this approach is that only the double mutant has the phenotype: either single mutant is functional (this explains the 'synthetic' in synthetic lethality).

Consider a disruption allele that has no strong phenotype, but for other reasons the gene is assumed to affect the process of interest. Is there a redundant gene or pathway that keeps the cells functioning in the absence of the original gene? If there is a single additional gene that is redundant, then mutating it in a background that already lacks the first gene will confer the desired phenotype. Only the doubly mutant strain would have the phenotype; either mutant alone would be normal. This can be considered as the parallel pathway model: the cells are




Figure 6 | Plasmid shuffle. The plasmid shuffle technique can be used to analyse individual protein function or isolate conditioned alleles. Using a balance between selective and counter-selective pressure, cells can be isolated that either contain a plasmid or are cured of it. Selection can be conferred by a nutritional marker, or by an additional gene on the plasmid that might be required for viability. Using colour-based assays (see illustration in BOX 1), an additional visual screen might be incorporated to allow rapid assessment of plasmid presence. In this example, a null mutation in chromosomal yfg1 (your favourite gene 1) is rescued by the wild-type version ($yfg1^+$) that is present on a selectable plasmid in the fission yeast Schizosaccharomycespombe. This strain is transformed with a library of plasmids that contain mutant variants of yfg1 (yfg1*). If the yfg1* plasmid can be shuffled out of the transformed cells (here, by selecting against the presence of plasmid-borne ura4 and ade6), then this is a sign that the $yfg1^*$ allele can rescue the lethality of the chromosomal yfg1 mutation; the $yfg1^*$ allele can then be isolated and studied further. 5-FOA, 5'-fluoro-orotic acid.

viable as long as one pathway is intact. An example of this is the overlapping activity of the Cln1, Cln2 and Cln3 cyclins that control the G1 phase of the cell cycle in budding yeast: any one is sufficient for viability and severe phenotypes are only observed if all are deleted³⁹.

Conversely, using synthetic lethality as a tool to probe direct interactions requires beginning the screen with alleles that attenuate, but do not abolish, the function of a protein. Consider ts mutants that are grown at the permissive temperature. They are viable but, in many cases, the protein function is attenuated even at the permissive temperature. Mutation of another gene in the pathway can interact genetically to abolish function: the cells can tolerate any single insult to the pathway of interest, but cannot tolerate more than one. This can be considered as the protein complex, or single pathway, model. For example, a network of genetic interactions between a collection of budding yeast replication mutants, including suppression and synthetic lethality, provided the first evidence that the proteins they encode form functional complexes in vivo⁴⁰. The complication of the synthetic lethal approach is that if the process under study is essential, one must use a screen rather than a selection. The investigator is looking for conditions in which a viable cell will die.

To isolate synthetic mutants, one begins with a strain that contains the original mutation and a plasmid with the wild-type version of the gene. The cells are mutagenized and then screened for loss of the plasmid on the appropriate counter-selection agent (BOX 1 and FIG. 5) under permissive conditions, to isolate strains that have lost the plasmid, and with it, the wild-type gene YFG1. The desired mutants will never

survive on the counter-selection agent, because they have a mutation that requires the presence of the wild-type copy of the gene for viability. So, the investigator isolates the mutants that are sensitive to the counter-selection agent. This strategy can be improved by including a colour marker to facilitate rapid identification of colonies that have lost the plasmid (BOXES 1 and 2).

If the starting strain contained a null mutant in the original gene that was caused by deletion of that gene, then any new mutation that results in synthetic lethality must occur in another locus. If the starting strain contained a conditional mutant, then the new mutation could also occur in the original gene and render it completely inactive, or it could represent a mutation in a second gene that produces a synthetic lethal phenotype. These can be distinguished genetically in crosses to appropriately marked strains, to determine whether the new mutation is linked to the original mutant, whether it has a phenotype by itself and whether the effect results from a single genetic lesion. The desired phenotype must result from the interaction between the two mutations yfg1 and syn1.

Again, cloning of the genes that correspond to synthetic lethal mutants can be a challenge. In some cases, the *syn1* mutant causes an independent phenotype by itself, such as heat or cold sensitivity, or drug sensitivity. If this phenotype is recessive, the corresponding gene can be cloned by complementation of the syn1 phenotype. Often, however, there is no obvious independent phenotype and other means must be used. These depend on whether the *syn1* mutation is dominant or recessive. For example, if the haploid strain yfg1^{ts} syn1 is inviable, but the diploid strain yfg1^{ts}/yfg1^{ts}

Box 2 | Genetic screens: a case study

An excellent example of a synthetic lethal screen and plasmid shuffle analysis was published recently⁴⁷. Saccharomyces cerevisiae SGS1 encodes a helicase that is required for genome stability and recovery from replication insults, but is not essential for viability. S. J. Brill and colleagues⁴⁷ isolated mutants in six genes that are required for viability in a $\triangle sgs1$ background. The authors used a variation on the colour assay described in BOX 1, by using a double ade mutant. The starter strain was ade2 ade3, which is unable to grow without adenine. This strain forms white colonies because the ade3 mutation is epistatic to ade2 — ade3 lies upstream in the metabolic pathway and prevents formation of the red pigment that normally accumulates in ade2 mutants. With wild-type ADE3 on a starting plasmid, which also contains SGS1 and URA3, transformed colonies will accumulate the red pigment. Cells were mutagenized and plated to look for colonies that were solidly red without sectors, indicating that they required the plasmid for viability. These clones were unable to form colonies on 5'flouro-orotic acid (5-FOA) plates. Six complementation groups were isolated that depended on the plasmid for viability; because several alleles were isolated for each group, the authors concluded that their screen was approaching saturation. The mutants were called slx1-6. Characterization of these new mutants indicated that they formed three phenotypic classes, with two slx mutants in each.

Cloning of the wild-type SLX genes used a plasmid shuffle method (FIG. 6). Each $\triangle sgs1$ slx strain, containing a plasmid with SGS1, ADE3, and URA3 markers, was transformed with a plasmid library that contains the LEU2 marker. Upon plating on 5-FOA, clones that no longer required the original SGS1 URA3 ADE3 plasmid for viability were selected. These now relied instead on the LEU2 plasmid, which contained either the wild-type SGS1 or the wild-type SLX gene. SGS1-containing plasmids were identified on the basis of their ability to rescue the lethality of all $\sin \Delta sgs1$ slx mutant strains. Isolation of the SLX plasmids allowed identification of the open reading frames (which were for the most part previously uncharacterized), construction of targeted gene disruptions and molecular experiments. Intriguingly, the two SLX genes in each phenotypic class seem to interact with each other in a complex.

This example shows how classical yeast genetics methods can be used to identify genes with no previously known function, to place them in existing networks and to predict molecular interactions.

> syn1/SYN1 is viable, then the synthetic lethal mutation syn1 is recessive. If this is the case, the investigator begins to clone SYN1 using yfg1ts syn1 cells kept alive by YFG1 on the original plasmid, and transforms them with a plasmid library that contains wild-type genes. Upon plating on the counter-selection agent, the original plasmid is driven out and only those cells that contain a library plasmid with YFG1 or SYN1 are able to grow. This is a form of the plasmid shuffle screen (FIG. 6), described in more detail below. An example of a synthetic lethal screen that used this cloning strategy is described in BOX 2.

> As is the case for suppressors, synthetic phenotypes can also be generated by overproduction of wild-type genes from a library in a mutant background⁴¹. This synthetic dosage lethality phenotype might reflect direct interactions, or downstream effectors, and the plasmids are easily recovered and their inserts identified by sequencing.

> More commonly, synthetic dosage lethality is used to probe the response of the cell to a known mutant protein expressed in high dosage. Overexpression of a mutant form of *yfg1* might be tolerated by wild-type cells, but not by mutant cells. This can be tested on a case-by-case basis using previously isolated mutant strains, or can be the basis of a screen by identifying mutants that are sensitive to overproduction of the

defective protein, when wild-type cells are resistant. For example, expression of a mutant version of fission yeast Mcm2 that lacks its nuclear localization sequence (NLS) is tolerated in wild-type cells, but is lethal in mcm2ts mutants even at the permissive temperature. The NLS mutant out-competes the attenuated ts protein and sequesters the other members of the MCM complex in the cytoplasm. Only the wild-type Mcm2 is sufficiently robust to withstand this lethal trap⁴².

Plasmid-based screens and functional analysis

Both of the yeast species are easily transformed by extrachromosomal plasmids that contain a selectable marker and an origin of replication. By combining forward selection for transformants, and counter-selection to isolate cells that are cured of the plasmid, the investigator gains considerable control in manipulating gene expression and plasmid dosage.

Plasmids can be used as markers for chromosomal dynamics, including DNA replication (mediated by the plasmid origin of replication) and segregation. For budding yeast, the centromere is small enough to be encompassed on an average plasmid; for fission yeast, the much larger centromere can be contained on a large minichromosome, which must be transferred genetically from strain to strain⁴³. The presence or absence of a plasmid in a cell can be determined by direct selection, or by a colony colour-based visual screen (BOX 1). The latter screen is readily done and allows the investigator to determine whether all the cells in a colony have lost the plasmid, or whether they are being lost at a rapid rate, by determining whether colony colour is uniform or sectored

Perhaps the most useful plasmid-screening method is the analysis of individual protein function using the plasmid shuffle44 (FIG. 6). This method starts with a null allele of $\Delta yfg1$ in the chromosome, kept alive by wildtype $yfg1^+$ on a counter-selectable plasmid. A library of yfg1* mutants (consisting of mutagenized genes) on a compatible plasmid with a different marker is constructed by in vitro mutagenesis, for example with hydroxylamine. The method can be enhanced by including a colour marker on the counter-selectable plasmid (BOX 1). The starting strain is transformed with the yfg1* library and plated on the counter-selection agent to see whether the wild-type plasmid can be shuffled out, its function provided by one of the yfg1* plasmids. If it can, then the colonies kept alive by yfg1* can be screened to see if they have any associated phenotypes (for example, drug resistance or temperature sensitivity). Ultimately, the yfg1* allele can be recovered and used to replace the endogenous chromosomal copy for additional study. This method is commonly used to isolate ts alleles and replace them in the chromosome (for example, REF. 45).

Conclusion and perspectives

With the completion of the genome sequences of both yeast species, and the classical genetics tools described here, it is clear that the yeasts are entering a new golden age46. Their limited complement of genes

Links

 $\textbf{DATABASE LINKS} \ Cdc2 \ | \ Cdc13 \ | \ actin \ | \ Cdc7 \ | \ MCM5 \ | \ Cln1 \ | \ Cln2 \ | \ Cln3 \ | \ Mcm2 \ | \ ADE2$ | ade6⁺ | LEU2 | CAN1 | can1⁺ | URA3 | ura4⁺ | TRP1 | SGS1 | ade3 | slx1-6 | pat1 | cdc28 |

YEAST PHYLOGENY Ascomycota | Fungal phylogeny

GENOME DATABASES Saccharomyces Genome Database (SGD) | Sanger Centre S. pombe database | Proteome (YPD and pombePD)

PROTOCOLS S. cerevisiae | S. pombe

 $\textbf{OTHER YEAST RESOURCES} \ Yeast \ virtual \ library \ | \ Gene \ conversion \ table \ | \ \textit{Saccharomyces}$ genome deletion project | TRIPLES

LABS AND INVESTIGATORS S. cerevisiae labs | S. pombe labs | Susan Forsburg's lab **COMMUNITY INFORMATION** S. cerevisiae | S. pombe

provides biologists with streamlined systems for understanding fundamental principles in eukaryotic cell biology. Remarkably, the classical methods discussed here are still timely and in wide use; modest refinements are made, but the principles remain the same. These methods help to meet the continuing challenge of the genomic era to identify the function of individual genes. The insights from the yeasts therefore continue to inform experiments in more complex cells, and cell biologists can only benefit from adding these versatile and tractable cells to their toolkits.

- Goffeau, A. et al. Life with 6000 genes, Science 274. 546–567 (1996).
- Beggs, J. D. Transformation of yeast by a replicating hybrid plasmid. *Nature* **275**, 104–109 (1978).
- Rothstein, R. One step gene disruption in yeast. *Methods Enzymol.* **101**, 202–211 (1983).
- Oliver, S. G. From gene to screen with yeast. Curr. Opin Genet, Dev. 7, 405-409 (1997).
- Oliver, S. G., Winson, M. K., Kell, D. B. & Banganz, F. Systematic functional analysis of the yeast genome. Trends Biotechnol. **16**, 373–378 (1998).
- Sipiczki, M. Phylogenesis of fission yeasts contradictions surrounding the origin of a century old genus. *Antonie Van Leeuwenhoek* **68**, 119–149 (1995).
- Paquin, B. et al. The fungal mitochondrial genome project: evolution of fungal mitochondrial genomes and their gene xpression. Curr. Genet. 31, 380-395 (1997).
- Berbee, M. L. & Taylor, J. W. Dating the evolutionary radiations of the true fungi. Can. J. Bot. 71, 1114-1127
- Keogh, R. S., Seoighe, C. & Wolfe, K. H. Evolution of gene order and chromosome number in Saccharomyce Kluyveromyces and related fungi. Yeast 14, 443-457
- Forsburg, S. L. The best yeast. Trends Genet. 15, 340-344 (1999).
 - Summarizes some differences in the biology of the
- two yeast species.
 11. Guthrie, C. & Fink, G. R. (eds) Guide to yeast genetics and molecular biology. Methods Enzymol. 194, 1-863 (1991). Describes more specific methods and protocols for both yeast species.
- Moreno, S., Klar, A. & Nurse, P. Molecular genetic analysis of the fission yeast *Schizosaccharomyces pombe*. Methods Enzymol. 194, 795–823 (1991).
- Kumar, A. & Snyder, M. Emerging technologies in yeast genomics. *Nature Rev. Genet.* 2, 302–312 (2001). The genomics revolution complements the classical genetics approach.
- Hoffman, C. S. & Welton, R. Mutagenesis and gene cloning in *Schizosaccharomyces pombe* using nonhomologous plasmid integration and rescue. Biotechniques 28, 532-539 (2000).
- 15. Chua, G., Taricani, L., Strangle, W. & Young, P. G. Insertional mutagenesis based on illegitimate recombination in Schizosaccharomyces pombe. Nucleic Acids Res. 28, E53 (2000).
- Grallert, B., Nurse, P. & Patterson, T. E. A study of integrative transformation in Schizosaccharomyces pombe. Mol. Gen. Genet. 238, 26-32 (1993).
- Hughes, T. R. et al. Widespread aneuploidy revealed by DNA microarray expression profiling. Nature Genet. 25,

- 333-337 (2000).
- Wolfe, K. H. & Shields, D. C. Molecular evidence for an ancient duplication of the entire yeast genome. Nature 387, 708-713 (1997).
- Giaever, G. et al. Genomic profiling of drug sensitivities via induced haploinsufficiency. Nature Genet. 21, 278-283
- Hartwell, L., Culotti, J. & Reid, B. Genetic control of the cell division cycle in yeast. I. Detection of mutants. Proc. Natl Acad. Sci. USA 66, 352-359 (1970).
- Nurse, P. Genetic control of cell size at cell division in yeast. Nature 256, 547-551 (1975).
- Nurse, P. Universal control mechanism regulating onset of M phase. Nature 344, 503-508 (1990).
- Enoch, T., Carr, A. M. & Nurse, P. Fission yeast genes involved in coupling mitosis to completion of DNA replication. *Genes Dev.* **6**, 2035–2046 (1992). Li, R. & Murray, A. W. Feedback control of mitosis in
- budding yeast. Cell 66, 519-532 (1991).
- Hoyt, M. A., Totis, L. & Roberts, B. T. S. cerevisiae genes required for cell cycle arrest in response to loss of microtubule function. Cell 66, 507-517 (1991).
- Koshland, D., Kent, J. C. & Hartwell, I., H. Genetic analysis of the mitotic transmission of minichromosomes. Cell 40, 393-403 (1985).
- Hieter, P., Mann, C., Snyder, M. & Davis, R. W. Mitotic stability of yeast chromosomes: a colony color assay that measures nondisjunction and chromosome loss. Cell 40, 381–382 (1985).
- Rine, J. Gene overexpression in studies of Saccharomyces cerevisiae. Methods Enzymol. 194,
- Stevenson, L. F., Kennedy, B. K. & Harlow, E. A large-scale overexpression screen in Saccharomyces cerevisiae identifies previously uncharacterized cell cycle genes Proc. Natl Acad. Sci. USA 98, 3946-3951 (2001).
- Herskowitz, I. Functional inactivation of genes by dominant negative mutations. *Nature* **329**, 219–222 (1987). Prelich, G. Suppression mechanisms: themes from
- variations. Trends Genet. 15, 261-266 (1999).
- Guarente, L. Synthetic enhancement in gene interaction; a genetic tool come of age. Trends Genet. 9, 362-366
 - References 28 and 30-32 are classic reviews describing the main features of the screens covered in this article and providing some theoretical background.
- Booher, R. & Beach, D. Interaction between $cdc13^+$ and $cdc2^+$ in the control of mitosis in fission yeast: dissociation of the G1 and G2 roles of the cdc2+ protein kinase FMBO J 6 3441-3447 (1987)
- Sandrock, T. M., O'Dell, J. L. & Adams, A. E. Allele-specific

- suppression by formation of new protein-protein interactions in yeast. Genetics 147, 1635-1642 (1997).
- Tye, B. K. & Sawyer, S. The hexameric eukaryotic MCM helicase: building symmetry from nonidentical parts J. Biol. Chem. 275, 34833-34836 (2000).
- Hardy, C. F. J., Dryga, O., Seematter, S., Pahl, P. M. B. & Sclafani, R. A. *mcm5/cdc46-bob1* bypasses the requirement for the S phase activator Cdc7p. Proc. Natl Acad. Sci. USA 94, 3151-3155 (1997).
- Lee, M. G. & Nurse, P. Complementation used to clone a human homologue of the fission yeast cell cycle control gene cdc2. Nature 327, 31-35 (1987).
- Grallert, B. & Nurse, P. An approach to identify functional homologues and suppressors of genes in fission yeast. Curr. Genet. 32, 27-31 (1997).
- Wittenberg, C., Sugimoto, K. & Reed, S. I. G1-specific cyclins of S. cerevisiae: cell cycle periodicity, regulation by mating pheromone and association with the p34^{CL} protein kinase. Cell **62**, 225–237 (1990)
- Hennessy, K. M., Lee, A., Chen, E. & Botstein, D. A group of interacting yeast DNA replication genes. Genes Dev. 5,
- Kroll, E. S., Hyland, K. M., Hieter, P. & Li, J. J. Establishing genetic interactions by a synthetic dosage lethality phenotype. Genetics 143, 95-102 (1996).
- Pasion, S. G. & Forsburg, S. L. Nuclear localization of fission yeast Mcm2/Cdc19p requires MCM complex formation. Mol. Biol. Cell 10, 4043-4057 (1999).
- Clarke, L. Centromeres: proteins, protein complexes, and repeated domains at centromeres of simple eukarvotes Curr. Opin. Genet. Dev. 8, 212–218 (1998).
- Sikorski, R. S. & Boeke, J. D. *In vitro* mutagenesis and plasmid shuffling: from cloned gene to mutant yeast. Methods Enzymol. **194**, 302–318 (1991).
- Liang, D. T. & Forsburg, S. L. Characterization of *S. pombe* $mcm7^+$ and $cdc23^+$ (MCM10) and interactions with replication checkpoints. Genetics (in the press).
- Johnston, M. The yeast genome: on the road to the Golden Age. *Curr. Opin. Genet. Dev.* **10**, 617–623 (2000).
- Mullen, J. R., Kaliaraman, V., Ibrahim, S. S. & Brill, S. J. Requirement for three novel protein complexes in the absence of the Sgs1 DNA helicase in *Saccharomyces* cerevisiae. Genetics **157**, 103–118 (2001)

An excellent example of a synthetic lethal screen and plasmid shuffle analysis.

Acknowledgements

I thank L. Pillus for helpful comments on the manuscript. I am a scholar of the Leukemia Society of America. Support in my lab comes from the National Institutes of Health, the National Science Foundation and the American Cancer Society