

# Convex Sets

Kacper Kłos

April 4, 2025

## Exercise 2.1

Let  $C \subseteq \mathbf{R}^n$  be a convex set, with  $x_1, \dots, x_k \in C$ , and let  $\theta_1, \dots, \theta_k \in \mathbf{R}$  satisfy  $\theta_i \geq 0$ ,  $\theta_1 + \dots + \theta_k = 1$ . Show that  $\theta_1 x_1 + \dots + \theta_k x_k \in C$ . (The definition of convexity is that this holds for  $k = 2$ ; you must show it for arbitrary  $k$ .) *Hint.* Use induction on  $k$ .

**Solution.** We know that

$$\theta_1 + \dots + \theta_{k-2} + \tilde{\theta}_{k-1} + \tilde{\theta}_k = 1 \quad \text{and} \quad \theta_1 x_1 + \dots + \theta_{k-2} x_{k-2} + \tilde{\theta}_{k-1} \tilde{x}_{k-1} + \tilde{\theta}_k \tilde{x}_k \in C.$$

Next, define

$$\theta_{k-1} = \tilde{\theta}_{k-1} + \tilde{\theta}_k.$$

From this we see that

$$1 = \frac{\tilde{\theta}_{k-1} + \tilde{\theta}_k}{\theta_{k-1}},$$

so, using the fact that  $C$  is convex, we know that there is a  $x_{k-1}$  such that

$$\tilde{\theta}_{k-1} \tilde{x}_{k-1} + \tilde{\theta}_k \tilde{x}_k = \theta_{k-1} x_{k-1}.$$

Plugging this back into the original equations, we get

$$\theta_1 + \dots + \theta_{k-1} = 1 \quad \text{and} \quad \theta_1 x_1 + \dots + \theta_{k-1} x_{k-1} \in C,$$

thus reducing two  $x_i$ 's into one. Repeating this procedure recursively leaves us with a single point  $x_0$ , which must lie in  $C$  by its convexity.  $\square$