

ABE 721

Agricultural Systems Modelling and Simulation

Module 4: Model Validation and Advanced Analysis

Prepared by:

Kadeghe Fue, PhD, P.Eng(T)

Electronics and Precision Agriculture Laboratory (EPAL)

Sokoine University of Agriculture (SUA)

Morogoro, Tanzania

Contents

4 Model Validation and Analysis **3**

4.1 4.1 Epistemology and Metrics of Model Validation 3

4.1.1 Quantitative Statistical Metrics of Fit 4

4.2 4.2 Stochastic Models and Analytical Uncertainty 7

4.2.1 Random Variables in Agricultural Systems 7

4.2.2 Monte Carlo Simulation Methods 8

4.3 4.3 System Dynamics: Stocks, Flows, and Networks 10

4.3.1 The Fundamental Concepts of System Dynamics 10

4.3.2 Mathematical Integration of SD 10

4.3.3 Case Study: The Agricultural Soil Nitrogen Cycle 11

4.4 4.4 Causal Loop Diagrams (CLDs) and Archetypes 13

4.4.1 Components and Rules of a CLD 13

4.4.2 Feedback Loops 13

4.4.3 Case Study: Irrigation and Soil Salinity Paradox 14

4.5 4.5 Equilibrium, Stability, and Bifurcation Analysis 15

4.5.1 Mathematical Definition of Equilibrium 15

4.5.2 Stability and the Phase Plane 15

4.5.3 Multi-Variable Jacobian Stability Proof (Advanced) 16

4.6 4.6 Advanced Models of S-Shaped Growth 17

4.6.1 System Dynamics Interpretation of Sigmoidal Growth 17

4.6.2 Mathematical Formulations of Growth 18

4.7 4.7 Advanced Hydrological Networks: Water Flows and SPAC 20

4.7.1 The Soil-Plant-Atmosphere Continuum (SPAC) 20

4.7.2 The Electrical Resistance Analogy for Flow 20

4.7.3 Evapotranspiration: The Penman-Monteith Formulation . . . 21

Chapter 4

Model Validation and Analysis

Following the rigorous mathematical formulation detailed in Module 2 and the complex numerical execution and calibration frameworks established in Module 3, the agricultural engineering process must inevitably transition to rigorous evaluation and structural analysis. A mathematical model, no matter how theoretically sound, is practically useless if its predictive capabilities cannot be statistically verified against physical reality.

Module 4 explores the deep statistical validation of models, the critical incorporation of stochastic uncertainty to account for chaotic agricultural environments, and the powerful visual and mathematical paradigms of System Dynamics (Stocks, Flows, and Causal Loops) heavily utilized in environmental modelling (Ford, 1999). Finally, we apply these advanced concepts to the intricate hydrological flows within agricultural systems, specifically focusing on the Soil-Plant-Atmosphere Continuum (SPAC).

4.1 4.1 Epistemology and Metrics of Model Validation

Validation answers a fundamental epistemological question in engineering: *Does this mathematical abstraction accurately and reliably represent physical reality under novel, untested conditions?*

It is crucial to distinguish between **Verification** and **Validation**. Verification ensures that the computer code correctly solves the mathematical equations (i.e., ”Are we building the model right?”). Validation, however, ensures that the mathe-

matical equations actually represent the real world (i.e., "Are we building the right model?"). A calibrated model must *always* be validated against an independent, completely separate dataset that was not utilized during the parameter estimation phase. This is often achieved through data splitting techniques, such as reserving 70% of field data for calibration and 30% for validation.

4.1.1 Quantitative Statistical Metrics of Fit

Visual inspection of a simulated curve plotted alongside observed data points is highly subjective and scientifically insufficient. Agricultural engineers must utilize specific, standardized statistical indices to quantify the error and the goodness-of-fit. Let $Y_{obs,i}$ be the i-th observed value, $Y_{sim,i}$ be the i-th simulated value, $\bar{Y}_{obs}$ be the mean of observed values, and n be the total number of observations.

1. Root Mean Square Error (RMSE)

RMSE quantifies the absolute fit of the model to the data. It calculates the standard deviation of the residuals (prediction errors). It is measured in the exact same physical units as the state variable (e.g., kg/ha for crop yield, or m^3/m^3 for soil moisture).

$$RMSE = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\left(Y_{obs,i} - Y_{sim,i}\right)^2} \tag{4.1}$$

Because the errors are squared before they are averaged, the RMSE gives a relatively high weight to large errors. This means the RMSE is highly sensitive to outliers. An RMSE of 0 indicates a perfect fit.

2. Mean Absolute Error (MAE)

While RMSE squares the errors, the Mean Absolute Error simply measures the average magnitude of the errors without considering their direction. It is a linear score which means that all the individual differences are weighted equally in the average.

$$MAE = \frac{1}{n}\sum_{i=1}^{n}|Y_{obs,i} - Y_{sim,i}| \tag{4.2}$$

Comparing RMSE and MAE can provide insight into the distribution of error magnitudes. If $RMSE \gg MAE$, it indicates the presence of severe outliers in the simulation's predictions.

3. Nash-Sutcliffe Efficiency (NSE)

Crucial in hydrological, watershed, and agricultural modelling, the NSE normalizes the model's accuracy relative to the mean of the observed data. It indicates how well the plot of observed versus simulated data fits the $1:1$ line.

$$NSE = 1 - \frac{\sum_{i=1}^{n}(Y_{obs,i} - Y_{sim,i})^2}{\sum_{i=1}^{n}(Y_{obs,i} - \bar{Y}_{obs})^2} \tag{4.3}$$

The NSE provides a standardized metric of model utility:

- $NSE = 1.0$: Perfect, flawless match between the model and reality.
- $NSE = 0.0$: The model is only as accurate as simply guessing the historical average for every prediction.
- $NSE < 0.0$: The residual variance is larger than the data variance. The model is worse than simply using the historical average. The underlying mathematical structure is fundamentally flawed.

4. Percent Bias (PBIAS)

Percent bias measures the average tendency of the simulated data to be larger or smaller than their observed counterparts. It is an essential metric to determine if an agricultural model is systematically overestimating or underestimating a phenomenon (e.g., consistently under-predicting evaporation).

$$PBIAS = \left[\frac{\sum_{i=1}^{n}(Y_{obs,i} - Y_{sim,i}) \times 100}{\sum_{i=1}^{n} Y_{obs,i}}\right] \% \tag{4.4}$$

The optimal value of PBIAS is 0.0, indicating highly accurate model simulation. Positive values indicate model underestimation bias, and negative values indicate model overestimation bias.

5. Index of Agreement (d)

Developed by Willmott (1981), the index of agreement is a standardized measure of the degree of model prediction error which varies between 0 and 1.

$$d = 1 - \frac{\sum_{i=1}^{n}(Y_{obs,i} - Y_{sim,i})^2}{\sum_{i=1}^{n}\left(|Y_{sim,i} - \bar{Y}_{obs}| + |Y_{obs,i} - \bar{Y}_{obs}|\right)^2} \tag{4.5}$$

A value of 1 indicates a perfect match, and 0 indicates no agreement at all. The index of agreement can detect additive and proportional differences in the observed and simulated means and variances.

4.2 4.2 Stochastic Models and Analytical Uncertainty

Most agricultural processes are driven by the weather, biology, and human intervention, all of which are inherently chaotic. Deterministic models—which produce a single, exact numerical output for a given set of inputs—fail to capture the true risk reality of farming. Stochastic models incorporate probability distributions to generate a *range* of possible outcomes with associated confidence intervals, moving from absolute prediction to probabilistic forecasting.

4.2.1 Random Variables in Agricultural Systems

Inputs such as daily rainfall, solar radiation, wind speed, or machinery breakdown events are treated mathematically as random variables. For example, monthly rainfall in the Morogoro region cannot be represented as a constant scalar; it is often modelled using specific probability density functions (PDFs).

Common Probability Distributions

1. The Normal (Gaussian) Distribution: Used for symmetric phenomena, such as the measurement error of soil sensors or the distribution of crop heights in a genetically uniform field.

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \tag{4.6}$$

Where μ is the mean and σ^2 is the variance.

2. The Log-Normal Distribution: Hydrological variables, such as daily precipitation or streamflow, cannot be negative and are often heavily right-skewed. The log-normal distribution is ideal for this. If X is log-normally distributed, then $Y = \ln(X)$ is normally distributed.

$$f(x|\mu, \sigma) = \frac{1}{x\sigma\sqrt{2\pi}} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right) \quad \text{for } x > 0 \tag{4.7}$$

3. The Weibull Distribution: In precision agriculture and renewable energy de-

sign on farms, the Weibull distribution is the international standard for modelling stochastic wind speeds (v).

$$f(v|k,c) = \frac{k}{c}\left(\frac{v}{c}\right)^{k-1} e^{-(v/c)^k} \quad \text{for } v \geq 0 \tag{4.8}$$

Where k is the dimensionless shape parameter and c is the scale parameter (in m/s).




Figure 4.1: Probability Density Functions (PDFs). Note how the Log-Normal distribution restricts values to greater than zero and features a heavy right tail, making it ideal for hydrological modelling.

4.2.2 Monte Carlo Simulation Methods

The Monte Carlo method is the standard computational technique for stochastic risk analysis in agricultural systems. It relies on the law of large numbers to approximate complex, non-linear probability distributions that cannot be solved analytically.

The Standard Monte Carlo Algorithm:

1. Define the exact probability distributions for all uncertain exogenous inputs and parameters (e.g., $Rainfall \sim Lognormal(\mu, \sigma)$).
2. Use a pseudo-random number generator to sample one specific value from every defined distribution.
3. Run the deterministic mathematical simulation model (e.g., an RK4 crop growth

ODE solver) using these uniquely sampled values to generate one final output instance (e.g., $Yield_1$).

4. Repeat Steps 2 and 3 iteratively for thousands of iterations ($N \approx 10,000$).
5. Statistically analyze the resulting aggregate distribution of the outputs to determine means, variances, and 95% confidence intervals.

Latin Hypercube Sampling (LHS)

A major limitation of standard Monte Carlo simulation is that it requires a massive number of iterations to adequately sample the extreme "tails" of a probability distribution (e.g., 1-in-100 year flood events).

Latin Hypercube Sampling (LHS) is an advanced stratified sampling method used by engineers to massively reduce computational time. LHS divides the cumulative probability distribution into N equal intervals of probability ($1/N$). It then ensures that exactly one sample is randomly drawn from *every single interval*. This guarantees that the entire spectrum of possibilities—including extreme rare events—is tested uniformly without requiring millions of simulation runs.

Instead of stating deterministically: *"The farm will yield 4.2 tons/ha,"* a robust stochastic LHS model concludes: *"There is a 90% probability that the yield will fall between 3.8 and 4.6 tons/ha, with a statistically verified 5% risk of total crop failure due to extreme weather variables."* This probabilistic approach is absolutely crucial for modern agricultural financial and insurance risk analysis.

4.3 4.3 System Dynamics: Stocks, Flows, and Networks

System Dynamics (SD), initially pioneered by Jay Forrester at the Massachusetts Institute of Technology (MIT), provides a highly rigorous visual and mathematical language for framing complex, non-linear systems. As noted by Ford (1999), it is the premier methodology for environmental and agricultural systems modelling.

4.3.1 The Fundamental Concepts of System Dynamics

1. Stocks (Levels / Accumulations) Stocks represent the state variables of the system. They are physical or conceptual entities that accumulate or deplete over time. A critical rule of System Dynamics is that if time were to instantly freeze, stocks would still possess a measurable, physical value.

- *Examples:* Water volume in a reservoir (m^3), tractor diesel fuel ($Liters$), pest population ($Count$), soil nitrate concentration (kg/ha), or farm capital (TZS).
- *Visual Notation:* Represented conventionally as solid rectangles.

2. Flows (Rates / Derivatives) Flows represent the active rate of change of the stocks. They only exist dynamically over a period of time. If time freezes, a flow instantly becomes zero.

- *Examples:* Evaporation rate (m^3/day), fuel consumption ($Liters/hr$), pest birth rate ($Count/day$), nitrogen leaching rate ($kg/ha/day$).
- *Visual Notation:* Represented as pipes with directional arrows and butterfly valves. "Clouds" at the ends of pipes represent infinite sources or sinks located strictly outside the defined system boundary.

4.3.2 Mathematical Integration of SD

The visual representation of System Dynamics maps perfectly to integral calculus and differential equations. The absolute value of any Stock at time t is the mathematical integration of its Inflows minus its Outflows, plus the initial starting condition at time t_0:

$$Stock(t) = Stock(t_0) + \int_{t_0}^{t} \left(\sum Inflows(s) - \sum Outflows(s) \right) ds \tag{4.9}$$

By taking the derivative with respect to time, this yields the standard Ordinary Differential Equation (ODE) state-space formulation explored in Module 2:

$$\frac{d(Stock)}{dt} = \sum Inflows - \sum Outflows \tag{4.10}$$

4.3.3 Case Study: The Agricultural Soil Nitrogen Cycle

To demonstrate the power of System Dynamics in agriculture, consider the complex network of the soil nitrogen cycle. This cannot be modelled as a single equation; it is a networked system of interconnected stocks and flows.

We can define three primary Stocks: 1. N_{org}: Organic Nitrogen (in plant residue and humus). 2. NH_4^+: Ammonium (available to plants, relatively immobile). 3. NO_3^-: Nitrate (available to plants, highly mobile in water).

The physiological Flows connecting these stocks are governed by specific rate parameters:

- **Mineralization (F_{min}):** Conversion of N_{org} to NH_4^+ by soil microbes.
- **Nitrification (F_{nit}):** Conversion of NH_4^+ to NO_3^- by bacteria (Nitrosomonas and Nitrobacter).
- **Denitrification (F_{den}):** Loss of NO_3^- to the atmosphere as N_2 gas under anaerobic conditions.
- **Leaching (F_{lea}):** Loss of NO_3^- downwards out of the root zone due to water percolation.
- **Plant Uptake (F_{up1}, F_{up2}):** Removal of NH_4^+ and NO_3^- by crop roots.

This visual SD map directly translates into the following coupled system of ordinary




Figure 4.2: System Dynamics representation of the Soil Nitrogen Cycle. The rectangles represent the state variable stocks, while the double-lined arrows represent the rate flows governed by differential equations.

differential equations:

$$\frac{dN_{org}}{dt} = Inputs_{residue} - F_{min} \tag{4.11}$$

$$\frac{dNH_4^+}{dt} = Inputs_{fert} + F_{min} - F_{nit} - F_{up1} \tag{4.12}$$

$$\frac{dNO_3^-}{dt} = F_{nit} - F_{den} - F_{lea} - F_{up2} \tag{4.13}$$

By assigning mathematical rate equations to each flow (e.g., making F_{nit} a function of soil temperature, moisture, and NH_4^+ concentration), an engineer can simulate exact nutrient availability over a 120-day cropping season and optimize fertilizer application schedules to minimize groundwater pollution (leaching).

4.4 4.4 Causal Loop Diagrams (CLDs) and Archetypes

While Stock and Flow diagrams provide the strict quantitative mathematical architecture required for numerical simulation, **Causal Loop Diagrams (CLDs)** map the conceptual, qualitative feedback mechanisms that actually drive the system's behavioral dynamics. They are primarily used by engineers and policy-makers to identify systemic archetypes and unintended consequences before mathematically formulating the differential equations.

4.4.1 Components and Rules of a CLD

- **Variables:** Nodes representing quantities that can vary over time (e.g., Soil Moisture, Crop Stress, Farmer Income, Groundwater Salinity).
- **Causal Links (Arrows):** Directed arrows indicating a direct causal relationship from variable A to variable B.
- **Polarities (+ or -):** The fundamental heart of a CLD.
  - **Positive (+):** Indicates that changes move in the *same direction*. If Variable A increases, Variable B increases above what it would have been. If A decreases, B decreases.
  - **Negative (-):** Indicates that changes move in *opposite directions*. If Variable A increases, Variable B decreases. If A decreases, B increases.
- **Delays (//):** Time delays in a system are critical causes of instability and oscillation. A double hash mark across an arrow indicates a significant delay between the cause and the effect.

4.4.2 Feedback Loops

Agricultural systems are governed by closed, self-referential loops of causality.

1. **Reinforcing Loops (R or +):** A loop contains an *even* number of negative links (or zero). These loops compound change, leading to exponential growth, vicious cycles, or total collapse. (e.g., More pests create more offspring, which

create more pests).

2. **Balancing Loops (B or -):** A loop contains an *odd* number of negative links. These loops naturally seek equilibrium, stabilization, or goal-seeking behavior, actively resisting extreme changes. (e.g., High crop water stress triggers stomatal closure, which reduces water loss, thereby reducing crop water stress).

4.4.3 Case Study: Irrigation and Soil Salinity Paradox

Consider a classic agricultural paradox modelled via CLDs: The attempt to increase crop yield through heavy irrigation in arid regions often leads to long-term systemic collapse due to soil salinity.




Figure 4.3: Causal Loop Diagram of Irrigation-Induced Salinity. The farmer applies water to increase yield (Balancing Loop B1). However, heavy irrigation raises the water table, eventually drawing salts into the root zone via capillary action (indicated by the delay marks //). This increases salinity, decreasing yield, which prompts the farmer to apply *more* water, triggering a vicious Reinforcing Loop (R1) of environmental degradation.

Through CLD analysis, engineers can identify that the solution is not to manipulate the flow of irrigation water, but to physically break the causal link between the water table and soil salinity by installing subsurface tile drainage systems.

4.5 4.5 Equilibrium, Stability, and Bifurcation Analysis

Understanding the long-term fate of an agricultural system requires analyzing its **Equilibrium Points** (also referred to as steady states or fixed points). An equilibrium point exists where the system's state variables are no longer changing over time, regardless of how much time passes.

4.5.1 Mathematical Definition of Equilibrium

For a continuous, autonomous dynamic system defined by a set of differential equations $\frac{d\mathbf{x}}{dt} = f(\mathbf{x})$, an equilibrium state vector $\mathbf{x}^*$ exists mathematically if and only if:

$$f(\mathbf{x}^*) = 0 \tag{4.14}$$

At this precise coordinate in state-space, the sum of all inflows to every stock perfectly equals the sum of all outflows. The system is static.

4.5.2 Stability and the Phase Plane

Identifying an equilibrium point is insufficient; we must classify its stability. If an agricultural system is perturbed slightly away from an equilibrium point (e.g., a sudden temperature spike in a greenhouse, or a sudden influx of migratory pests):

- **Stable Equilibrium (Attractor / Sink):** The natural forces and balancing feedback loops of the system will pull the state variables back to $\mathbf{x}^*$. The system is resilient.
- **Unstable Equilibrium (Repeller / Source):** The system will accelerate away from $\mathbf{x}^*$ upon the slightest disturbance. Reinforcing feedback loops dominate the dynamics, causing system collapse or explosion.

4.5.3 Multi-Variable Jacobian Stability Proof (Advanced)

To mathematically prove the stability of a non-linear agricultural system at equilibrium point $\mathbf{x}^*$, engineers use Taylor series expansions to linearize the complex system around $\mathbf{x}^*$.

Consider the classic **Lotka-Volterra** predator-prey model, widely used to model biological pest control in agriculture. Let V be the pest (Victim) population, and P be the biocontrol agent (Predator) population.

$$\frac{dV}{dt} = \alpha V - \beta VP = f_1(V, P) \tag{4.15}$$

$$\frac{dP}{dt} = \delta VP - \gamma P = f_2(V, P) \tag{4.16}$$

Where $\alpha, \beta, \delta, \gamma$ are positive biological parameters. By setting the derivatives to zero, we find a non-trivial equilibrium point at $V^* = \frac{\gamma}{\delta}$ and $P^* = \frac{\alpha}{\beta}$.

To determine if this agricultural ecosystem is stable, we construct the **Jacobian Matrix** (**J**) of partial derivatives evaluated precisely at the equilibrium point:

$$\mathbf{J}(V^*, P^*) = \begin{bmatrix} \frac{\partial f_1}{\partial V} & \frac{\partial f_1}{\partial P} \\ \frac{\partial f_2}{\partial V} & \frac{\partial f_2}{\partial P} \end{bmatrix}_{(V^*, P^*)} = \begin{bmatrix} \alpha - \beta P^* & -\beta V^* \\ \delta P^* & \delta V^* - \gamma \end{bmatrix} = \begin{bmatrix} 0 & -\frac{\beta\gamma}{\delta} \\ \frac{\delta\alpha}{\beta} & 0 \end{bmatrix} \tag{4.17}$$

The Stability Theorem: If the real parts of *all* eigenvalues (λ) of the Jacobian Matrix **J** are strictly negative, the agricultural system is locally asymptotically stable. To find the eigenvalues, we solve $\det(\mathbf{J} - \lambda\mathbf{I}) = 0$:

$$\det \begin{bmatrix} -\lambda & -\frac{\beta\gamma}{\delta} \\ \frac{\delta\alpha}{\beta} & -\lambda \end{bmatrix} = \lambda^2 + \alpha\gamma = 0 \implies \lambda = \pm i\sqrt{\alpha\gamma} \tag{4.18}$$

Because the eigenvalues are purely imaginary (the real part is exactly zero), the Lotka-Volterra equilibrium is formally classified as a *Center*. The pest and predator populations will oscillate continuously in closed loop cycles indefinitely, neither stabilizing nor exploding. This mathematical proof informs agricultural engineers that introducing a predator will control, but *never permanently eradicate*, a pest population.




Figure 4.4: Phase plane analysis of a 1D system. The system is at equilibrium where $dx/dt = 0$ (crossing the x-axis). x_1^* is an unstable repeller because slight perturbations drive the system away (df/dx is positive). x_2^* is a stable attractor because perturbations are countered, pulling the system back (df/dx is negative).

4.6 4.6 Advanced Models of S-Shaped Growth

S-shaped (Sigmoidal) growth is the most ubiquitous behavioral archetype in biological, ecological, and agricultural systems. It accurately describes the accumulation of dry matter in a crop canopy, the epidemiological spread of a plant pathogen across a region, or the sociological adoption rate of precision agriculture technologies among rural farmers.

4.6.1 System Dynamics Interpretation of Sigmoidal Growth

S-shaped growth is not caused by a single isolated causal loop. Rather, it is the mathematical consequence of a **shift in feedback loop dominance** over time.

1. **Phase 1 (Exponential Growth):** The system is entirely dominated by a **Reinforcing (R) loop**. For crop growth, this means new leaves capture more solar radiation, which produces more assimilates, which are used to grow even more leaves. The growth rate $\frac{dN}{dt}$ is constantly accelerating. 2. **Phase 2 (The Inflection Point):** A critical mathematical threshold is reached where environmental limits (space, nutrients, water) begin to restrict the system. The acceleration stops. 3. **Phase 3 (Asymptotic Goal-Seeking):** A **Balancing (B) loop** seizes absolute dominance over the Reinforcing loop. For crops, leaves begin to heavily shade each other (canopy closure), and soil nutrients become severely limiting. The growth rate

dynamically decelerates toward zero as the system asymptotically approaches its environmental carrying capacity (K).

4.6.2 Mathematical Formulations of Growth

While the Logistic Equation (Module 2) is common, agricultural modellers frequently use alternative mathematical formulations depending on the biological asymmetry of the growth.

1. The Logistic Growth Model

The classic symmetric S-curve. The maximum growth rate occurs exactly at half of the carrying capacity.

$$\frac{dN}{dt} = rN\left(1 - \frac{N}{K}\right) \tag{4.19}$$

As proven in previous modules, setting the second derivative to zero ($\frac{d^2N}{dt^2} = 0$) verifies that the exact inflection point occurs at $N = \frac{K}{2}$.

2. The Gompertz Growth Model

Many biological phenomena in agriculture (such as individual animal weight gain or fruit expansion) are fundamentally asymmetric. They grow rapidly initially, but approach their maximum size very slowly. The Gompertz model captures this asymmetry by making the exponential growth rate decay exponentially over time.

$$\frac{dN}{dt} = rN \ln\left(\frac{K}{N}\right) \tag{4.20}$$

The analytical solution to this differential equation yields the Gompertz curve:

$$N(t) = K \exp\left(\ln\left(\frac{N_0}{K}\right) e^{-rt}\right) \tag{4.21}$$

By finding the second derivative, we can prove that the inflection point for the Gompertz curve occurs at $N = \frac{K}{e} \approx 0.37K$. The shift in loop dominance happens much earlier than in the Logistic model.

3. The Richards (Generalized Logistic) Curve

For ultimate flexibility in agricultural calibration, the Richards curve introduces a shape parameter (ν) that can shift the inflection point to any location, absorbing both Logistic ($\nu = 1$) and Gompertz ($\nu \to 0$) curves as special cases.

$$N(t) = K \left[1 + \left(\left(\frac{K}{N_0} \right)^{\nu} - 1 \right) e^{-rt} \right]^{-\frac{1}{\nu}} \tag{4.22}$$




Figure 4.5: Mathematical variations of S-Shaped growth. The Gompertz model is highly asymmetric, exhibiting a much earlier shift in loop dominance (inflection point) compared to the strictly symmetric Logistic model.

4.7 4.7 Advanced Hydrological Networks: Water Flows and SPAC

We now culminate Module 4 by applying the rigorous principles of System Dynamics, non-linear Differential Equations, and Causal Loops to one of the most critical and complex domains in agricultural engineering: Hydrology and Irrigation. As established by Kramer (2012) and Michael (2008), the movement of water through a farm cannot be modelled accurately as isolated puddles; it must be modelled as a continuous dynamic flow driven by strict thermodynamic potential energy gradients.

4.7.1 The Soil-Plant-Atmosphere Continuum (SPAC)

The SPAC paradigm mandates that water moves from the deep soil, up through the biological plant tissues, and out into the vast atmosphere as a single, unified, continuous hydraulic system. Based on the Second Law of Thermodynamics, water flows strictly from regions of high water potential (Ψ) to regions of low water potential.

$$\Psi_{soil} > \Psi_{root} > \Psi_{stem} > \Psi_{leaf} > \Psi_{atmosphere} \tag{4.23}$$

4.7.2 The Electrical Resistance Analogy for Flow

The transient volumetric flow of water (q) through the SPAC can be modelled mathematically identically to electrical current flowing through a network of resistors (Ohm's Law). The water potential difference ($\Delta\Psi$) acts as the driving voltage, and the various biological and physical impediments (R) act as electrical resistance.

$$q = \frac{\Delta\Psi}{R_{total}} = \frac{\Psi_{soil} - \Psi_{root}}{R_{soil}} = \frac{\Psi_{root} - \Psi_{leaf}}{R_{plant}} = \frac{\Psi_{leaf} - \Psi_{atm}}{R_{stomata} + R_{boundary}} \tag{4.24}$$

By treating the soil profile and the biological plant canopy as **Stocks** of water, and the transpiration sap flow (q) as a dynamically restricted **Outflow**, engineers

can build a comprehensive computational simulation capable of predicting precise crop water stress levels on an hourly basis.




Figure 4.6: System Dynamics representation of the Soil-Plant-Atmosphere Continuum (SPAC). Water flows continuously up the extreme thermodynamic potential gradient. If the leaf potential drops dangerously low, a biological balancing feedback loop triggers stomatal closure, radically increasing the resistance (r_s) and artificially halting the outflow to prevent catastrophic plant desiccation.

4.7.3 Evapotranspiration: The Penman-Monteith Formulation

The ultimate outflow driver of the entire SPAC system into the atmosphere is Evapotranspiration (ET). In advanced agricultural systems modelling, empirical formulas like Blaney-Criddle are discarded in favor of the physically rigorous **FAO-56 Penman-Monteith Equation**. This equation brilliantly combines a thermodynamic energy balance (accounting for incoming solar radiation) with an aerodynamic mass transfer theory (accounting for wind sweeping vapor away).

$$ET_0 = \frac{\Delta(R_n - G) + \rho_a c_p \frac{(e_s - e_a)}{r_a}}{\lambda\left(\Delta + \gamma\left(1 + \frac{r_s}{r_a}\right)\right)} \tag{4.25}$$

Where the deeply complex parameters are defined as:

- R_n: Net solar radiation at the crop surface ($MJ/m^2/day$). Calculated as $R_n = (1-\alpha)R_s - R_{nl}$, where α is crop albedo.
- G: Soil heat flux density, often negligible on a daily timescale.
- $\rho_a c_p$: The volumetric heat capacity of the air.
- $(e_s - e_a)$: The Vapor Pressure Deficit (VPD) of the air, representing the physical "drying power" of the atmosphere.
- Δ: The slope of the saturation vapor pressure curve at the current air temperature.
- γ: The psychrometric constant.
- λ: Latent heat of vaporization of water.
- $\mathbf{r_a}$: **Aerodynamic resistance**, a function of wind speed (u_2) and crop height.
- $\mathbf{r_s}$: **Bulk surface (stomatal) resistance**, which biologically links right back into the balancing feedback loop of the SPAC diagram.

By embedding the Penman-Monteith equation directly into the continuous numerical ODE solvers developed in Module 3, engineers can accurately simulate dynamic soil moisture depletion. This allows for the computational scheduling of irrigation pumps to precisely replenish the soil moisture stock only when necessary, optimizing water use efficiency on a regional scale.

End of Module 4.