

Intranasal neural protocol

NOVA-ELITE 2 cc via intranasal catheter · NOVA-MSC FLEX 1–2 cc deltoid IM

Intranasal administration through the olfactory and trigeminal pathways bypasses the blood–brain barrier. Exosomes may support reduction of neuroinflammation, neuronal repair signaling, and modulation.

TOTAL VOLUME

2 cc

1 cc per nostril

INSTILL RATE

10–20 sec

slow, per nostril

HEAD POSITION

30–45°

back-tilted, hold 8–10 min

DELTOID IM

1–2 cc

NOVA-MSC FLEX

Preparation

- Confirm no active sinus infection, severe congestion, or recent nasal surgery
- No nose blowing for 30 minutes before the procedure
- Review contraindications (pregnancy, active infection, recent chemotherapy)
- Obtain informed consent
- Seated in a treatment chair, head tilted back 30–45° to expose the olfactory cleft
- Draw 2 cc into sterile syringes (1 cc per nostril); remove air bubbles
- Attach syringes to the catheter per manufacturer instructions

POST-PROCEDURE INSTRUCTIONS

- No nose blowing for at least 60 minutes
- Avoid bending forward for 20–30 minutes
- Hydrate normally; avoid alcohol the same day
- Mild throat drainage sensation is normal

SAFETY

- Observe 5–10 minutes post-procedure
- Monitor for sinus discomfort or infection
- Consider ENT clearance for severe deviated septum

Administration sequence

- 1 Left nostril.** Insert catheter into position; instill 1 cc slowly over 10–20 s. Maintain head-back position and hold 8–10 minutes. Instruct the patient not to sniff heavily or swallow forcefully.
- 2 10-minute interval.** Keep the head back; wait 10 minutes before the second nostril.
- 3 Right nostril.** Instill 1 cc slowly over 10–20 s; maintain head-back position another 8–10 minutes.
- 4 Deltoid IM.** Administer 1–2 cc NOVA-MSC FLEX intramuscularly.

FOLLOW-UP

2 weeks

First review

6 weeks

Second review

4–6 wks

Repeat interval by severity

EXPECTED RESPONSES

- Reduced hyperarousal, intrusive thoughts, and nightmares
- Improved emotional regulation and cognitive clarity
- Typical onset 3–7 days; fuller change at 8–12 weeks

Today you received an exosome product given as a fine mist through the nose, plus one intramuscular cell injection in your upper arm. The products used are made from donated birth tissue and are not FDA-approved treatments. This page tells you what is normal in the days ahead, how to care for yourself, and when to call us.

DATE OF TREATMENT

SITE(S) TREATED

NEXT VISIT

WHAT TO EXPECT

Changes are gradual — do not judge the result in the first days

FIRST FEW HOURS

A drip at the back of the throat, an odd taste, mild congestion, or sneezing are all normal, along with soreness at the arm injection site.

FIRST WEEK

Nothing dramatic is expected. Keep your symptom diary daily — small notes beat big impressions.

WEEKS 4–8

We review your diary against baseline at follow-up and decide whether a repeat treatment makes sense.

DO

- Stay reclined at the clinic for the full observation period
- Keep the daily symptom diary we gave you
- Drink water normally through the day
- Take acetaminophen (Tylenol) for arm soreness if needed

AVOID

- Blowing your nose forcefully for 4–6 hours
- Nasal sprays for 24 hours unless prescribed
- Alcohol for 72 hours
- Anti-inflammatory pain relievers for 72 hours

CALL US RIGHT AWAY IF

For an emergency, call 911 first

- A nosebleed that does not stop with gentle pressure
- A severe headache or a stiff neck with fever
- Fever above 101 °F
- New confusion, or trouble breathing

CLINIC PHONE

AFTER-HOURS PHONE

NOTES FROM YOUR VISIT