

Hair & mesotherapy

NOVA-E2 2 cc meso gun · optional NOVA-M1 1-2 cc deltoid IM

Mesotherapy delivery places MSC-derived exosomes and signaling factors densely into dermal and superficial tissue layers to support localized regeneration, collagen remodeling, and tissue repair. Optional intramuscular NOVA-M1 adds systemic MSC signaling.

MESO VOLUME**2 cc**

NOVA-E2, evenly distributed

PER INJECTION**0.01-0.05 cc**

grid pattern, 0.5-1 cm spacing

DEPTH SETTING**0.5-2.0 mm**

0.5-1.0 dermal · 1.0-2.0 soft tissue

OPTIONAL IM**1-2 cc**

deltoid, 25-27 G

Preparation

- Confirm no active infection at the treatment site
- Avoid NSAIDs 24-48 h prior if possible
- Review contraindications (pregnancy, malignancy, etc.)
- Obtain informed consent
- Seated or supine by treatment area; expose and sterilize the region
- Draw 2 cc NOVA-E2 into the cartridge; clear air bubbles

POST-PROCEDURE INSTRUCTIONS

- Avoid strenuous activity 24-48 h
- Avoid NSAIDs 48 h
- Hydrate well
- Expect mild redness or swelling

SAFETY

- Observe 5-10 minutes post-procedure
- Monitor for infection or adverse reaction; maintain sterile technique

Administration

MESOTHERAPY

1. Set meso gun depth by indication.
2. Deliver microinjections in a grid-like pattern, 0.5-1 cm apart.
3. 0.01-0.05 cc per injection; 2 cc total evenly distributed.

DELTOID IM (OPTIONAL)

1. Inject 1-2 cc NOVA-M1 into the deltoid muscle.
2. 25-27 G needle; single or split injection sites.

SCHEDULE**Initial**

Every 2-4 weeks × 3 sessions

Maintenance

Every 3-6 months

EXPECTED RESPONSES

- Improved tissue quality and regeneration
- Enhanced collagen and elastin production
- Reduced inflammation and pain in MSK application
- Visible change typically within 1-4 weeks

PATIENT NAME	DATE OF BIRTH	PATIENT ID / MRN	DATE OF VISIT
EVALUATING PHYSICIAN	INITIAL · FOLLOW-UP VISIT TYPE — CIRCLE	REFERRING PROVIDER, IF ANY	TIME IN / OUT

S SUBJECTIVE

Hair thinning / loss —
 CHIEF COMPLAINT

ONSET / DURATION

SEVERITY — VAS 0-10

QUALITY / RADIATION

AGGRAVATING FACTORS

RELIEVING FACTORS

Minoxidil · finasteride · PRP · supplements · low-level laser
 PRIOR TREATMENT — CIRCLE ALL TRIED · ADD DATES

duration of thinning · family history · shedding pattern

fever · chills · weight loss · neuro changes — note positives

FUNCTIONAL IMPACT

PERTINENT ROS

MEDICATIONS — INCL. ANTICOAGULANTS, IMMUNOSUPPRESSANTS

ALLERGIES · RELEVANT PMH — CANCER, INFECTION, PREGNANCY

O OBJECTIVE

BP

HR

TEMP

WEIGHT

GENERAL APPEARANCE

Pattern — Norwood or Ludwig stage

Scalp — erythema, scarring, scale; rule out other dermatoses

Pull test — positive / negative

Density photos — frontal, mid, vertex, temporal, standardized angles

Part-width comparison

IMAGING / STUDIES REVIEWED — MODALITY, DATE, FINDINGS

BASELINE — Stage

BASELINE — Photos

BASELINE — Shed count

A ASSESSMENT

Androgenetic alopecia · diffuse hair thinning
 WORKING DIAGNOSIS — CONFIRM OR AMEND

ADDITIONAL DIAGNOSES / CODES

☐ Appropriate candidate — indications met, contraindications absent

☐ Alternatives discussed, including conservative care and doing nothing

☐ Non-FDA-approved status explained; no outcome promised

☐ Patient questions answered; understanding confirmed

P PLAN

Proposed treatment. Scalp mesotherapy: NOVA-E2 2-3 cc as intradermal microdepots across mapped zones, 30-32 G, meso gun or serial puncture, with optional NOVA-M1 IM. Series of 3 sessions, 4-6 weeks apart.

☐ Informed consent (CN-PC) reviewed and signed

☐ Procedure performed today — see CN-OP-HAIR · or scheduled (date right)

☐ Fees, non-coverage by insurance, and refund policy reviewed

☐ Follow-up: next session 4-6 wk; photos every visit

PHYSICIAN SIGNATURE	PRINTED NAME & LICENSE NUMBER	PROCEDURE DATE, IF SCHEDULED
---------------------	-------------------------------	------------------------------

PATIENT NAME

DATE OF BIRTH

PATIENT ID / MRN

DATE OF PROCEDURE

PERFORMING PHYSICIAN

Androgenetic alopecia · diffuse hair thinning

PRE-PROCEDURE DIAGNOSIS — CONFIRM OR AMEND

START TIME

ASSISTING STAFF

PRE-PROCEDURE CHECKLIST

Check each box before the first draw

- ☐ Informed consent (CN-PC) signed and on file; SOAP note (CN-SN-HAIR) complete
- ☐ Screening reviewed — no contraindications present
- ☐ Time-out performed — correct patient, site, product, route
- ☐ Five checks read aloud with second staff: ID · product & lot · expiry · volume & route · consent
- ☐ Baseline outcome measures recorded: Standardized photos · density · shed count
- ☐ Site marked and prepped — chlorhexidine / povidone-iodine (circle); chain-of-custody log initialed

BP · HR · TEMP

agent · concentration · volume — write NONE i...

BP · HR · TEMP

PRE-PROCEDURE VITALS

LOCAL ANESTHETIC

POST-OBSERVATION VITALS

PRODUCTS ADMINISTERED

Session ___ of 3 · series spaced 4-6 weeks apart

PRODUCT & TARGET	PLANNED	LOT	EXPIRY	GIVEN
NOVA-E2 (exosome) Intradermal, mapped thinning zones of the scalp	2-3 cc			
NOVA-M1 IM (optional) Deltoid IM, per treatment plan	per plan			

TECHNIQUE

Position: seated; scalp sections mapped and cleansed

- Intradermal microdepots across mapped zones via meso gun or serial puncture.
 - Shallow, even placement — intradermal, not subgaleal.
 - Standardized photographs taken before injection, same angles each session.
- ZONES: FRONTAL · MID-SCALP · VERTEX · TEMPORAL — circle all treated**

SITE / SIDE

MESO GUN · SERIAL PUNCTURE — circle

GUIDANCE

30-32 G

NEEDLE USED

target 15 min

OBSERVATION — MINUTES

NARRATIVE — FINDINGS, ASPIRATE, DEVIATIONS FROM PROTOCOL, PATIENT RESPONSE

POST-PROCEDURE & DISPOSITION

- Tolerated: WELL · WITH EVENTS — circle
- Discharged: STABLE · AMBULATORY · WITH ESCORT — circle
- ☐ Patient note CN-PN-HAIR reviewed and given at discharge

COMPLICATIONS / ADVERSE EVENTS — WRITE "NONE" IF NONE

Any adverse event also requires the adverse-event response steps in this book.

Follow-up plan. Next session in 4-6 weeks; photographic comparison at every visit across the 3-session series.

FOLLOW-UP SCHEDULED FOR

PHYSICIAN SIGNATURE

PRINTED NAME & LICENSE NUMBER

DATE & END TIME

Today you received a series of very small injections of an exosome product into the skin of your scalp. The products used are made from donated birth tissue and are not FDA-approved treatments. This page tells you what is normal in the days ahead, how to care for yourself, and when to call us.

DATE OF TREATMENT

SITE(S) TREATED

NEXT VISIT

WHAT TO EXPECT

Changes are gradual — do not judge the result in the first days

FIRST 24-48 HOURS

Pinpoint redness, tenderness, and tiny bumps at the injection points are normal, as are small spots of dried blood. These settle within a day or two.

WEEKS 1-4

A temporary increase in shedding can occur and is part of the normal hair cycle. No visible change is expected yet.

MONTHS 2-4

The earliest changes in density and texture appear here. Results are judged across the full series using your photos.

DO

- Sleep with your head slightly elevated tonight if tender
- Wash gently after 24 hours with a mild shampoo
- Take acetaminophen (Tylenol) for tenderness if needed
- Keep every session in the series — spacing matters

AVOID

- Washing or wetting the scalp for 24 hours
- Hair coloring or chemical treatments for 1 week
- Tight hats or helmets for 24 hours
- Heavy sweating exercise for 24 hours, and scratching the sites

CALL US RIGHT AWAY IF

For an emergency, call 911 first

- Spreading redness, warmth, or swelling of the scalp
- Pustules, crusting, or drainage at injection points
- Fever above 101 °F
- A severe or unusual headache

CLINIC PHONE

AFTER-HOURS PHONE

NOTES FROM YOUR VISIT