

Patient screening

Applies to every indication, in addition to protocol-specific criteria

Universal inclusion

- Diagnosis documented and consistent with the selected protocol
- Medically stable, no active systemic or local infection
- Realistic expectations discussed, including that outcomes vary and are not guaranteed
- Able to attend follow-up and complete outcome measures
- Written informed consent obtained before the procedure (page 16)

Absolute exclusion

- Active systemic infection, sepsis, or infection at the intended site
- Active or untreated malignancy
- Pregnancy or breastfeeding, unless approved by the Medical Director
- Known hypersensitivity to a product component or diluent
- Inability to give or have a proxy give informed consent

RELATIVE — DOCUMENT THE DECISION

Uncontrolled diabetes, active autoimmune flare, therapeutic anticoagulation, immunosuppression, recent corticosteroid injection at the site, or recent live vaccination.

Pre-procedure checklist

- | | |
|--|---|
| ☐ History, exam, and imaging reviewed for this indication | ☐ Anticoagulant or antiplatelet hold discussed with prescriber |
| ☐ Infection screen: no fever, no active skin lesion at site | ☐ NSAIDs and corticosteroids addressed per protocol |
| ☐ Pregnancy status confirmed where applicable | ☐ Allergy history reviewed, including anesthetic and diluent |
| ☐ Baseline outcome measures recorded (pain, function, photos) | ☐ Emergency medications and anaphylaxis kit checked in date |
| ☐ Consent signed, witnessed, and filed | ☐ Product, lot, expiry, and COA verified against the order |
| ☐ Post-procedure instructions printed and reviewed with patient | ☐ Transport home arranged where sedation or IN delivery is used |
| ☐ Guidance equipment available and tested where the protocol calls for it | ☐ Second staff member available for the five pre-dose checks (page 14) |
| ☐ Chain-of-custody line opened for each vial (page 18) | ☐ Follow-up appointment booked before the patient leaves |

Medication guidance

NSAIDS

Consider holding 5–7 days before and 2 weeks after, using acetaminophen for comfort where appropriate.

CORTICOSTEROIDS

Avoid injection at the same site within 4–6 weeks before or after, unless clinically necessary.

ANTICOAGULATION

Any hold is decided with the prescribing physician; never stop therapy on the day of the procedure alone.

SCREENING COMPLETED BY _____

DATE _____

PHYSICIAN APPROVAL _____