

Thaw & reconstitution

Bench sequence, diluent volumes, and the working window

THAW BATH**37 °C**

never above · 2–4 min typical

WORKING WINDOW**15–30 min**

thaw to administration

REFREEZE**Never**

single use · discard remainder

Thaw sequence — cryopreserved product

- 1 Confirm patient, indication, product, lot, and expiry against the order and the COA.
- 2 Prepare the field, syringes, needles, and guidance equipment before removing product.
- 3 Thaw in a 37 °C dry block or water bath, keeping the cap above the water line.
- 4 Remove as soon as the last ice crystal clears; swirl gently, never vortex or shake.
- 5 Inspect for particulates, discoloration, or a compromised seal — do not use if present.
- 6 Draw the full dose using aseptic technique; expel air without aerosolizing.
- 7 Note the thaw time on the record — administration is timed from this point.
- 8 Administer within the working window; discard any unused volume per policy.

Lyophilized reconstitution

VIAL	DILUENT	RESULT
Standard	1 mL sterile saline	1 cc working volume
Meso / topical	2–3 mL sterile saline	Dilute field coverage
Intranasal	Per protocol, page 08	Atomizer-ready volume

Introduce diluent slowly down the vial wall, then rotate gently until fully dissolved — typically under two minutes. The solution should be clear to faintly opalescent with no visible pellet.

Do not

- Vortex, shake, or sonicate — mechanical stress degrades the product.
- Warm above 37 °C or thaw in a microwave.
- Refreeze, pool lots, or hold a drawn syringe for a later patient.
- Filter through a 0.22 µm filter unless the label directs it.
- Co-mix with anesthetic, corticosteroid, or contrast in the same syringe unless directed.
- Use product with a missing or mismatched COA.