

Dosing quick reference

Every protocol in this book, summarized for the procedure room

INDICATION	PRODUCT	VOLUME	ROUTE / GUIDANCE	NEEDLE	REPEAT
Knee · 03	NOVA-MSC FLEX + NOVA-EXOS CORE	3-5 cc	Intra-articular · US optional	21-25 G	6-12 mo
Hip · 04	Perinatal MSC ± exosome	4-6 cc	IA + IM · US or fluoro	20-22 G	6-12 mo
Shoulder · 06	NOVA-CORE + NOVA-FLEX	2-4 cc	IA / peritendinous · US-guided	22-25 G	6-12 mo
Cervical spine · 07	NOVA-EXO+ + NOVA-FLEX	5 cc total	Bilateral trapezius IM + deltoid IM	23-25 G	6-12 mo
Hair / meso · 08	NOVA-ELITE ± NOVA-FLEX IM	2-3 cc	Intradermal · meso gun	30-32 G	4-6 wk × 3
Intranasal neural · 09	NOVA-ELITE + NOVA-MSC FLEX IM	1-2 cc IN	Intranasal catheter + IM	n/a · 23 G IM	4-8 wk
Dementia / MCI · 10	NOVA-ELITE + NOVA-MSC FLEX IM	2 cc IN	Intranasal catheter + deltoid IM	n/a · 23 G IM	4-6 wk
Spina bifida A · 11	EXM 2 cc + PV-MAX 3 cc	5 cc total	IM · gluteal preferred	22-25 G	4-8 wk
Spina bifida B · 11	NOVA-MSC FLEX + NOVA-CORE	7 cc total	IM · bilateral gluteal	22-25 G	4-8 wk

READ FIRST This page is a summary only and is not a substitute for the full protocol. Spina bifida regimens A and B are alternatives, not additive. Confirm every value against the protocol page and the lot-specific COA before drawing a dose.

VOLUME LIMITS BY IM SITE

- Gluteal: up to 3 cc per side in adults — preferred for larger volumes
- Deltoid: maximum 2 cc per side
- Vastus lateralis: pediatric and small-frame patients per weight
- Split any dose exceeding the site limit across sites, not into one site
- Aspirate before injecting and inject slowly at every IM site
- Observe 15-30 minutes after any systemic or intranasal administration

BEFORE EVERY DOSE — FIVE CHECKS

- PATIENT ID
- PRODUCT & LOT
- EXPIRY
- VOLUME & ROUTE
- CONSENT ON FILE

Read the five checks aloud with a second staff member present and initial the chain-of-custody log before injection.

IF ANYTHING DOES NOT MATCH

Stop, return the vial to storage, and resolve the discrepancy with the ordering physician before proceeding. A mismatch between order, label, and COA is never corrected at the bedside.