

IBD-associated arthritis

NOVA-FLOW + NOVA-FLEX (gluteal) + NOVA-EXO+ (deltoid) · IM · physician reference protocol

NOVA-FLOW**4 cc**

IM · divided gluteal

NOVA-FLEX**2 cc**

IM · gluteal

NOVA-EXO+**2 cc**

IM · deltoid

OBSERVATION**15-30 min**

post-injection

A regenerative IM approach for HLA-B27-associated inflammatory bowel disease and enteropathic arthritis: NOVA-FLOW with NOVA-FLEX (gluteal) and NOVA-EXO+ (deltoid) to support modulation of chronic autoimmune inflammation.

Indications

- Enteropathic arthritis with Crohn's disease or ulcerative colitis
- HLA-B27-associated inflammatory joint symptoms
- Peripheral or axial arthropathy with IBD
- Adjunct to a physician-directed treatment plan

Contraindications & cautions

- Active infection or sepsis
- Known hypersensitivity to any product (NOVA-FLOW, NOVA-FLEX, NOVA-EXO+)
- Uncontrolled coagulopathy or therapeutic anticoagulation — caution with IM injection
- Acute flare with systemic instability — stabilize before treatment
- Pregnancy or breastfeeding unless cleared by the treating physician

Procedure

PREPARATION

1. Review disease activity, medications (incl. anticoagulants / immunosuppressants), and allergies; obtain informed consent.
2. Confirm preparation, dose, and integrity of NOVA-FLOW, NOVA-FLEX, and NOVA-EXO+.
3. Prep gluteal and deltoid sites with chlorhexidine or povidone-iodine.
4. Select IM needle length for body habitus (e.g., 22-25 G, 1-1.5 in).
5. Have anaphylaxis / emergency supplies on hand.

POST — PATIENT INSTRUCTIONS

- Local soreness is expected; warm compress as needed
- Limit strenuous use of the injected limb 24-48 h
- Report any rash, wheezing, or swelling
- Hydrate well and resume activity as tolerated

IM INJECTION SEQUENCE

1. **NOVA-FLOW 4 cc** — IM, divided across gluteal sites (≤ 3 cc per site), slow injection.
2. **NOVA-FLEX 2 cc** — IM, gluteal.
3. **NOVA-EXO+ 2 cc** — IM, deltoid.

Aspirate before injecting per site practice, inject slowly, space/rotate sites, and monitor for local or systemic reaction.

FOLLOW-UP & OUTCOME MEASURES

Reassess disease activity at **4-6 weeks**. Repeat on a physician-directed cadence (e.g., **every 4-8 weeks**) based on response and tolerance.

SYMPTOM INDEX**CRP / CALPROTECTIN****ARTHRALGIA****QUALITY OF LIFE**

REGULATORY CelluNOVA products are supplied for use by licensed healthcare professionals. CelluNOVA does not direct, prescribe, or control the practice of medicine, and nothing here guarantees a clinical outcome.

PATIENT NAME

DATE OF BIRTH

PATIENT ID / MRN

DATE OF VISIT

PROVIDER

INITIAL · FOLLOW-UP

VISIT TYPE — CIRCLE

REFERRING PROVIDER, IF
ANY

TIME IN / OUT

S SUBJECTIVE

CHIEF COMPLAINT / GOALS

ONSET / DURATION

SEVERITY — VAS 0-10

PRIOR TREATMENTS TRIED

FUNCTIONAL IMPACT

MEDICATIONS — INCL. ANTICOAGULANTS /
IMMUNOSUPPRESSANTS

ALLERGIES · RELEVANT PMH

O OBJECTIVE

BP

HR

TEMP

WEIGHT

GENERAL APPEARANCE

Exam / relevant findings

BASELINE — VAS

BASELINE — CRP

BASELINE — CALPROTECTIN

BASELINE — SYMPTOM IDX

A ASSESSMENT**Enteropathic / IBD-associated arthritis (HLA-B27)**

WORKING DIAGNOSIS — CONFIRM OR AMEND

ADDITIONAL DIAGNOSES / CODES

☐ Appropriate candidate — indications met, contraindications absent☐ Alternatives discussed, including conservative care and doing nothing☐ Non-FDA-approved status explained; no outcome promised☐ Patient questions answered; informed consent obtained**P PLAN**

Administer **NOVA-FLOW 4 cc** IM (divided gluteal, ≤ 3 cc per site), **NOVA-FLEX 2 cc** IM gluteal, and **NOVA-EXO+ 2 cc** IM deltoid. Observe 15–30 min. Reassess disease activity at 4–6 weeks; repeat per physician cadence.

PROVIDER SIGNATURE

DATE

PATIENT NAME

DATE OF BIRTH

PATIENT ID / MRN

DATE OF VISIT

PROVIDER

INITIAL · FOLLOW-UP

VISIT TYPE — CIRCLE

REFERRING PROVIDER, IF
ANY

TIME IN / OUT

PRE-PROCEDURE CHECKLIST

- | | |
|---|--|
| ☐ Informed consent verified | ☐ Patient identity / timeout confirmed |
| ☐ Allergies and current medications reviewed | ☐ Product preparation, dose, and integrity verified |
| ☐ IV / IM access and site prep complete | ☐ Anaphylaxis / emergency supplies available |

PRODUCTS ADMINISTERED

PRODUCT	LOT #	EXP	VOLUME	SITE / ROUTE
NOVA-FLOW			4 cc	IM · gluteal (+ sites)
NOVA-FLEX			2 cc	IM · gluteal
NOVA-EXO+			2 cc	IM · deltoid

TECHNIQUE & TOLERANCE

Administration notes (approach, sequence, spacing)

TIME IN

TIME OUT

WELL TOLERATED · REACTION

TOLERANCE — CIRCLE

Adverse events / interventions (none, or describe)

DISPOSITION

- | | |
|--|---|
| ☐ Observed 15–30 min post-administration | ☐ No acute reaction on discharge |
| ☐ Post-care instructions given to patient | ☐ Follow-up scheduled / advised |

PROVIDER SIGNATURE

DATE

PATIENT NAME

DATE OF VISIT

PROVIDER

WHAT WE DID TODAY

You received a set of intramuscular injections as part of your regenerative treatment plan: NOVA-FLOW and NOVA-FLEX in the hip (gluteal) muscle, and NOVA-EXO+ in the shoulder (deltoid) muscle.

WHAT TO EXPECT

- Mild soreness, tightness, or bruising at the injection sites for a day or two
- Occasional temporary fatigue or low-grade achiness
- Gradual response — reassessed with your provider over the following weeks

AFTERCARE**DO**

- Use a warm compress on sore sites
- Stay hydrated and rest as needed
- Take acetaminophen if needed and appropriate
- Keep your follow-up appointment

AVOID

- Strenuous use of the injected limbs for 24-48 h
- NSAIDs for 48-72 h unless your provider approves
- Alcohol for 24 h
- Applying heat before checking with your provider if a site is red/hot

WHEN TO CALL US

Call the clinic — or seek urgent care — for: spreading rash or hives, wheezing or trouble breathing, facial or throat swelling, fever, or severe or worsening pain at an injection or IV site.

YOUR NEXT VISIT

FOLLOW-UP DATE

CLINIC CONTACT

PATIENT SIGNATURE — RECEIVED & UNDERSTOOD

DATE