

# Intranasal neural protocol

NOVA-ELITE 2 cc via intranasal catheter · NOVA-MSF FLEX 1-2 cc deltoid IM

Intranasal administration through the olfactory and trigeminal pathways bypasses the blood-brain barrier. Exosomes may support reduction of neuroinflammation, neuronal repair signaling, and modulation.

**TOTAL VOLUME****2 cc**

1 cc per nostril

**INSTILL RATE****10-20 sec**

slow, per nostril

**HEAD POSITION****30-45°**

back-tilted, hold 8-10 min

**DELTOID IM****1-2 cc**

NOVA-MSF FLEX

## Preparation

- Confirm no active sinus infection, severe congestion, or recent nasal surgery
- No nose blowing for 30 minutes before the procedure
- Review contraindications (pregnancy, active infection, recent chemotherapy)
- Obtain informed consent
- Seated in a treatment chair, head tilted back 30-45° to expose the olfactory cleft
- Draw 2 cc into sterile syringes (1 cc per nostril); remove air bubbles
- Attach syringes to the catheter per manufacturer instructions

**POST-PROCEDURE INSTRUCTIONS**

- No nose blowing for at least 60 minutes
- Avoid bending forward for 20-30 minutes
- Hydrate normally; avoid alcohol the same day
- Mild throat drainage sensation is normal

**SAFETY**

- Observe 5-10 minutes post-procedure
- Monitor for sinus discomfort or infection
- Consider ENT clearance for severe deviated septum

## Administration sequence

- 1 Left nostril.** Insert catheter into position; instill 1 cc slowly over 10-20 s. Maintain head-back position and hold 8-10 minutes. Instruct the patient not to sniff heavily or swallow forcefully.
- 2 10-minute interval.** Keep the head back; wait 10 minutes before the second nostril.
- 3 Right nostril.** Instill 1 cc slowly over 10-20 s; maintain head-back position another 8-10 minutes.
- 4 Deltoid IM.** Administer 1-2 cc NOVA-MSF FLEX intramuscularly.

**FOLLOW-UP****2 weeks**

First review

**6 weeks**

Second review

**4-6 wks**

Repeat interval by severity

**EXPECTED RESPONSES**

- Reduced hyperarousal, intrusive thoughts, and nightmares
- Improved emotional regulation and cognitive clarity
- Typical onset 3-7 days; fuller change at 8-12 weeks

CHART DOCUMENTATION · EVALUATION VISIT

|                      |                                            |                            |               |
|----------------------|--------------------------------------------|----------------------------|---------------|
| PATIENT NAME         | DATE OF BIRTH                              | PATIENT ID / MRN           | DATE OF VISIT |
| EVALUATING PHYSICIAN | INITIAL · FOLLOW-UP<br>VISIT TYPE — CIRCLE | REFERRING PROVIDER, IF ANY | TIME IN / OUT |

**S SUBJECTIVE**

**Neurologic symptoms per history —**

|                 |                  |                     |
|-----------------|------------------|---------------------|
| CHIEF COMPLAINT | ONSET / DURATION | SEVERITY — VAS 0-10 |
|-----------------|------------------|---------------------|

|                     |                     |                   |
|---------------------|---------------------|-------------------|
| QUALITY / RADIATION | AGGRAVATING FACTORS | RELIEVING FACTORS |
|---------------------|---------------------|-------------------|

**Current neurology care · medications reviewed · prior therapies — list**

PRIOR TREATMENT — CIRCLE ALL TRIED · ADD DATES

|                                                                         |                                                                                |
|-------------------------------------------------------------------------|--------------------------------------------------------------------------------|
| cognition · energy · mood · sleep · daily function<br>FUNCTIONAL IMPACT | fever · chills · weight loss · neuro changes — note positives<br>PERTINENT ROS |
|-------------------------------------------------------------------------|--------------------------------------------------------------------------------|

|                                                        |                                                         |
|--------------------------------------------------------|---------------------------------------------------------|
| MEDICATIONS — INCL. ANTICOAGULANTS, IMMUNOSUPPRESSANTS | ALLERGIES · RELEVANT PMH — CANCER, INFECTION, PREGNANCY |
|--------------------------------------------------------|---------------------------------------------------------|

**O OBJECTIVE**

|    |    |      |        |                    |
|----|----|------|--------|--------------------|
| BP | HR | TEMP | WEIGHT | GENERAL APPEARANCE |
|----|----|------|--------|--------------------|

**General and mental status**

**Cranial nerves**

**Motor and coordination**

**Gait and balance**

**Nasal passages — patency, no active sinusitis**

|                                                       |                       |                         |                        |
|-------------------------------------------------------|-----------------------|-------------------------|------------------------|
| IMAGING / STUDIES REVIEWED — MODALITY, DATE, FINDINGS | BASELINE — Cog screen | BASELINE — Diary issued | BASELINE — Baseline sx |
|-------------------------------------------------------|-----------------------|-------------------------|------------------------|

**A ASSESSMENT**

**Physician-directed neurologic support (per treatment plan)**

|                                      |                              |
|--------------------------------------|------------------------------|
| WORKING DIAGNOSIS — CONFIRM OR AMEND | ADDITIONAL DIAGNOSES / CODES |
|--------------------------------------|------------------------------|

|                                                                                            |                                                                                                |
|--------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|
| <input type="checkbox"/> Appropriate candidate — indications met, contraindications absent | <input type="checkbox"/> Alternatives discussed, including conservative care and doing nothing |
| <input type="checkbox"/> Non-FDA-approved status explained; no outcome promised            | <input type="checkbox"/> Patient questions answered; understanding confirmed                   |

**P PLAN**

**Proposed treatment.** Intranasal NOVA-ELITE 1-2 cc in divided aliquots per naris plus NOVA-MSC FLEX 2 cc deltoid IM, 23 G. Observation 15-30 min. Symptom diary issued and explained.

|                                                                                      |                                                                                                |
|--------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|
| <input type="checkbox"/> Informed consent (CN-PC) reviewed and signed                | <input type="checkbox"/> Procedure performed today — see CN-OP-INN · or scheduled (date right) |
| <input type="checkbox"/> Fees, non-coverage by insurance, and refund policy reviewed | <input type="checkbox"/> Follow-up: review at 4-8 wk with symptom diary                        |

|                     |                               |                              |
|---------------------|-------------------------------|------------------------------|
| PHYSICIAN SIGNATURE | PRINTED NAME & LICENSE NUMBER | PROCEDURE DATE, IF SCHEDULED |
|---------------------|-------------------------------|------------------------------|

PATIENT NAME

DATE OF BIRTH

PATIENT ID / MRN

DATE OF PROCEDURE

PERFORMING PHYSICIAN

Physician-directed neurologic support (per treatment...)

PRE-PROCEDURE DIAGNOSIS — CONFIRM OR AMEND

START TIME

ASSISTING

STAFF

## PRE-PROCEDURE CHECKLIST

Check each box before the first draw

- |                                                                                                                |                                                                                                                             |
|----------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|
| <input type="checkbox"/> Informed consent (CN-PC) signed and on file; SOAP note (CN-SN-INN) complete           | <input type="checkbox"/> Screening reviewed — no contraindications present                                                  |
| <input type="checkbox"/> Time-out performed — correct patient, site, product, route                            | <input type="checkbox"/> Five checks read aloud with second staff: ID · product & lot · expiry · volume & route · consent   |
| <input type="checkbox"/> Baseline outcome measures recorded: Symptom diary · cognitive screen · adverse events | <input type="checkbox"/> Site marked and prepped — chlorhexidine / povidone-iodine (circle); chain-of-custody log initiated |

BP · HR · TEMP

PRE-PROCEDURE VITALS

agent · concentration · volume — write NONE i...

LOCAL ANESTHETIC

BP · HR · TEMP

POST-OBSERVATION VITALS

## PRODUCTS ADMINISTERED

Divide intranasal volume between nares in small aliquots

| PRODUCT & TARGET                                                                    | PLANNED       | LOT | EXPIRY | GIVEN |
|-------------------------------------------------------------------------------------|---------------|-----|--------|-------|
| <b>NOVA-ELITE (exosome)</b><br>Intranasal via atomizer or soft catheter, both nares | <b>1-2 cc</b> |     |        |       |
| <b>NOVA-MSF FLEX</b><br>Deltoid IM                                                  | <b>2 cc</b>   |     |        |       |

## TECHNIQUE

Position: reclined, head tilted back

- Small aliquots per naris, alternating sides; keep patient reclined 10–15 minutes after the final aliquot.
- Deltoid IM component: aspirate before injecting, inject slowly.
- Nasal passages clear before starting; postpone if significant congestion or active sinus infection.

**BOTH NARES + DELTOID L · R — circle deltoid side**  
SITE / SIDE**ATOMIZER · SOFT CATHETER — circle**  
GUIDANCEn/a intranasal · 23 G IM  
NEEDLE USEDtarget 15–30 min  
OBSERVATION — MINUTES

NARRATIVE — FINDINGS, ASPIRATE, DEVIATIONS FROM PROTOCOL, PATIENT RESPONSE

## POST-PROCEDURE &amp; DISPOSITION

Tolerated: **WELL · WITH EVENTS — circle**Discharged: **STABLE · AMBULATORY · WITH ESCORT — circle**

- ☐
- Patient note CN-PN-INN reviewed and given at discharge

COMPLICATIONS / ADVERSE EVENTS — WRITE "NONE" IF NONE

Any adverse event also requires the adverse-event response steps in this book.

**Follow-up plan.** Repeat consideration at 4–8 weeks based on the symptom diary and clinical response.

FOLLOW-UP SCHEDULED FOR

PHYSICIAN SIGNATURE

PRINTED NAME &amp; LICENSE NUMBER

DATE &amp; END TIME

Today you received an exosome product given as a fine mist through the nose, plus one intramuscular cell injection in your upper arm. The products used are made from donated birth tissue and are not FDA-approved treatments. This page tells you what is normal in the days ahead, how to care for yourself, and when to call us.

DATE OF TREATMENT

SITE(S) TREATED

NEXT VISIT

**WHAT TO EXPECT**

Changes are gradual — do not judge the result in the first days

**FIRST FEW HOURS**

A drip at the back of the throat, an odd taste, mild congestion, or sneezing are all normal, along with soreness at the arm injection site.

**FIRST WEEK**

Nothing dramatic is expected. Keep your symptom diary daily — small notes beat big impressions.

**WEEKS 4-8**

We review your diary against baseline at follow-up and decide whether a repeat treatment makes sense.

**DO**

- Stay reclined at the clinic for the full observation period
- Keep the daily symptom diary we gave you
- Drink water normally through the day
- Take acetaminophen (Tylenol) for arm soreness if needed

**AVOID**

- Blowing your nose forcefully for 4–6 hours
- Nasal sprays for 24 hours unless prescribed
- Alcohol for 72 hours
- Anti-inflammatory pain relievers for 72 hours

**CALL US RIGHT AWAY IF**

For an emergency, call 911 first

- A nosebleed that does not stop with gentle pressure
- A severe headache or a stiff neck with fever
- Fever above 101 °F
- New confusion, or trouble breathing

CLINIC PHONE

AFTER-HOURS PHONE

**NOTES FROM YOUR VISIT**