

Rheumatoid arthritis

NOVA-FLOW + NOVA-FLEX (gluteal) + NOVA-EXO+ (deltoid) · IM · physician reference protocol

NOVA-FLOW**4 cc**

IM · divided gluteal

NOVA-FLEX**2 cc**

IM · gluteal

NOVA-EXO+**2 cc**

IM · deltoid

OBSERVATION**15-30 min**

post-injection

A regenerative IM approach for rheumatoid arthritis: NOVA-FLOW with NOVA-FLEX (gluteal) and NOVA-EXO+ (deltoid) to support modulation of chronic autoimmune joint inflammation.

Indications

- Seropositive or seronegative rheumatoid arthritis
- Chronic symmetric inflammatory polyarthritis
- Persistent disease activity despite standard therapy
- Adjunct to a physician-directed treatment plan

Contraindications & cautions

- Active infection or sepsis
- Known hypersensitivity to any product (NOVA-FLOW, NOVA-FLEX, NOVA-EXO+)
- Uncontrolled coagulopathy or therapeutic anticoagulation — caution with IM injection
- Acute flare with systemic instability — stabilize before treatment
- Pregnancy or breastfeeding unless cleared by the treating physician

Procedure

PREPARATION

1. Review disease activity, medications (incl. anticoagulants / immunosuppressants), and allergies; obtain informed consent.
2. Confirm preparation, dose, and integrity of NOVA-FLOW, NOVA-FLEX, and NOVA-EXO+.
3. Prep gluteal and deltoid sites with chlorhexidine or povidone-iodine.
4. Select IM needle length for body habitus (e.g., 22-25 G, 1-1.5 in).
5. Have anaphylaxis / emergency supplies on hand.

POST — PATIENT INSTRUCTIONS

- Local soreness is expected; warm compress as needed
- Limit strenuous use of the injected limb 24-48 h
- Report any rash, wheezing, or swelling
- Hydrate well and resume activity as tolerated

IM INJECTION SEQUENCE

1. **NOVA-FLOW 4 cc** — IM, divided across gluteal sites (≤ 3 cc per site), slow injection.
2. **NOVA-FLEX 2 cc** — IM, gluteal.
3. **NOVA-EXO+ 2 cc** — IM, deltoid.

Aspirate before injecting per site practice, inject slowly, space/rotate sites, and monitor for local or systemic reaction.

FOLLOW-UP & OUTCOME MEASURES

Reassess disease activity at **4-6 weeks**. Repeat on a physician-directed cadence (e.g., **every 4-8 weeks**) based on response and tolerance.

DAS28

CRP / ESR

MORNING STIFFNESS

JOINT COUNT

HAQ

REGULATORY CelluNOVA products are supplied for use by licensed healthcare professionals. CelluNOVA does not direct, prescribe, or control the practice of medicine, and nothing here guarantees a clinical outcome.

PATIENT NAME	DATE OF BIRTH	PATIENT ID / MRN	DATE OF VISIT
PROVIDER	INITIAL · FOLLOW-UP VISIT TYPE — CIRCLE	REFERRING PROVIDER, IF ANY	TIME IN / OUT

S SUBJECTIVE

CHIEF COMPLAINT / GOALS	ONSET / DURATION	SEVERITY — VAS 0-10
PRIOR TREATMENTS TRIED	FUNCTIONAL IMPACT	
MEDICATIONS — INCL. ANTICOAGULANTS / IMMUNOSUPPRESSANTS	ALLERGIES · RELEVANT PMH	

O OBJECTIVE

BP	HR	TEMP	WEIGHT	GENERAL APPEARANCE
Exam / relevant findings				
BASELINE — VAS	BASELINE — DAS28	BASELINE — CRP/ESR	BASELINE — JOINT COUNT	

A ASSESSMENT

Rheumatoid arthritis		
WORKING DIAGNOSIS — CONFIRM OR AMEND	ADDITIONAL DIAGNOSES / CODES	
☐ Appropriate candidate — indications met, contraindications absent	☐ Alternatives discussed, including conservative care and doing nothing	
☐ Non-FDA-approved status explained; no outcome promised	☐ Patient questions answered; informed consent obtained	

P PLAN

Administer **NOVA-FLOW 4 cc** IM (divided gluteal, ≤3 cc per site), **NOVA-FLEX 2 cc** IM gluteal, and **NOVA-EXO+ 2 cc** IM deltoid. Observe 15–30 min. Reassess disease activity at 4–6 weeks; repeat per physician cadence.

PROVIDER SIGNATURE	DATE
--------------------	------

PATIENT NAME

DATE OF BIRTH

PATIENT ID / MRN

DATE OF VISIT

PROVIDER

INITIAL · FOLLOW-UP
VISIT TYPE — CIRCLEREFERRING PROVIDER, IF
ANY

TIME IN / OUT

PRE-PROCEDURE CHECKLIST

- | | |
|---|--|
| ☐ Informed consent verified | ☐ Patient identity / timeout confirmed |
| ☐ Allergies and current medications reviewed | ☐ Product preparation, dose, and integrity verified |
| ☐ IV / IM access and site prep complete | ☐ Anaphylaxis / emergency supplies available |

PRODUCTS ADMINISTERED

PRODUCT	LOT #	EXP	VOLUME	SITE / ROUTE
NOVA-FLOW			4 cc	IM · gluteal (+ sites)
NOVA-FLEX			2 cc	IM · gluteal
NOVA-EXO+			2 cc	IM · deltoid

TECHNIQUE & TOLERANCE

Administration notes (approach, sequence, spacing)

TIME IN

TIME OUT

WELL TOLERATED · REACTION

TOLERANCE — CIRCLE

Adverse events / interventions (none, or describe)

DISPOSITION

- | | |
|--|---|
| ☐ Observed 15–30 min post-administration | ☐ No acute reaction on discharge |
| ☐ Post-care instructions given to patient | ☐ Follow-up scheduled / advised |

PROVIDER SIGNATURE

DATE

PATIENT NAME

DATE OF VISIT

PROVIDER

WHAT WE DID TODAY

You received a set of intramuscular injections as part of your regenerative treatment plan: NOVA-FLOW and NOVA-FLEX in the hip (gluteal) muscle, and NOVA-EXO+ in the shoulder (deltoid) muscle.

WHAT TO EXPECT

- Mild soreness, tightness, or bruising at the injection sites for a day or two
- Occasional temporary fatigue or low-grade achiness
- Gradual response — reassessed with your provider over the following weeks

AFTERCARE**DO**

- Use a warm compress on sore sites
- Stay hydrated and rest as needed
- Take acetaminophen if needed and appropriate
- Keep your follow-up appointment

AVOID

- Strenuous use of the injected limbs for 24-48 h
- NSAIDs for 48-72 h unless your provider approves
- Alcohol for 24 h
- Applying heat before checking with your provider if a site is red/hot

WHEN TO CALL US

Call the clinic — or seek urgent care — for: spreading rash or hives, wheezing or trouble breathing, facial or throat swelling, fever, or severe or worsening pain at an injection or IV site.

YOUR NEXT VISIT

FOLLOW-UP DATE

CLINIC CONTACT

PATIENT SIGNATURE — RECEIVED & UNDERSTOOD

DATE