

Hair & mesotherapy

NOVA-ELITE 2 cc meso gun · optional NOVA-FLEX 1–2 cc deltoid IM

Mesotherapy delivery places MSC-derived exosomes and signaling factors densely into dermal and superficial tissue layers to support localized regeneration, collagen remodeling, and tissue repair. Optional intramuscular NOVA-FLEX adds systemic MSC signaling.

MESO VOLUME

2 cc

NOVA-ELITE, evenly distributed

PER INJECTION

0.01–0.05 cc

grid pattern, 0.5–1 cm spacing

DEPTH SETTING

0.5–2.0 mm

0.5–1.0 dermal · 1.0–2.0 soft tissue

OPTIONAL IM

1–2 cc

deltoid, 25–27 G

Preparation

- Confirm no active infection at the treatment site
- Avoid NSAIDs 24–48 h prior if possible
- Review contraindications (pregnancy, malignancy, etc.)
- Obtain informed consent
- Seated or supine by treatment area; expose and sterilize the region
- Draw 2 cc NOVA-ELITE into the cartridge; clear air bubbles

POST-PROCEDURE INSTRUCTIONS

- Avoid strenuous activity 24–48 h
- Avoid NSAIDs 48 h
- Hydrate well
- Expect mild redness or swelling

SAFETY

- Observe 5–10 minutes post-procedure
- Monitor for infection or adverse reaction; maintain sterile technique

Administration

MESOTHERAPY

1. Set meso gun depth by indication.
2. Deliver microinjections in a grid-like pattern, 0.5–1 cm apart.
3. 0.01–0.05 cc per injection; 2 cc total evenly distributed.

DELTOID IM (OPTIONAL)

1. Inject 1–2 cc NOVA-FLEX into the deltoid muscle.
2. 25–27 G needle; single or split injection sites.

SCHEDULE

Initial

Every 2–4 weeks × 3 sessions

Maintenance

Every 3–6 months

EXPECTED RESPONSES

- Improved tissue quality and regeneration
- Enhanced collagen and elastin production
- Reduced inflammation and pain in MSK application
- Visible change typically within 1–4 weeks

Today you received a series of very small injections of an exosome product into the skin of your scalp. The products used are made from donated birth tissue and are not FDA-approved treatments. This page tells you what is normal in the days ahead, how to care for yourself, and when to call us.

DATE OF TREATMENT

SITE(S) TREATED

NEXT VISIT

WHAT TO EXPECT

Changes are gradual — do not judge the result in the first days

FIRST 24–48 HOURS

Pinpoint redness, tenderness, and tiny bumps at the injection points are normal, as are small spots of dried blood. These settle within a day or two.

WEEKS 1–4

A temporary increase in shedding can occur and is part of the normal hair cycle. No visible change is expected yet.

MONTHS 2–4

The earliest changes in density and texture appear here. Results are judged across the full series using your photos.

DO

- Sleep with your head slightly elevated tonight if tender
- Wash gently after 24 hours with a mild shampoo
- Take acetaminophen (Tylenol) for tenderness if needed
- Keep every session in the series — spacing matters

AVOID

- Washing or wetting the scalp for 24 hours
- Hair coloring or chemical treatments for 1 week
- Tight hats or helmets for 24 hours
- Heavy sweating exercise for 24 hours, and scratching the sites

CALL US RIGHT AWAY IF

For an emergency, call 911 first

- Spreading redness, warmth, or swelling of the scalp
- Pustules, crusting, or drainage at injection points
- Fever above 101 °F
- A severe or unusual headache

CLINIC PHONE

AFTER-HOURS PHONE

NOTES FROM YOUR VISIT