

$$x^{\log_y n} \Longrightarrow n^{\log_y x}$$

$$x^{0} + x^{1} + x^{2} + \dots + x^{n} = \frac{x^{n+1} - 1}{x - 1}$$
 for  $x \neq 1$ 

$$x^0 + x^1 + x^2 + \dots = \frac{1}{1-x}$$
 for  $|x| < 1$ 


$$T(n) = \begin{cases} 1 & n = 1\\ 2T\left(\frac{n}{2}\right) + n^2 & n > 1 \end{cases}$$

Solve the following recurrence using the Recurrence Tree Method.

Assumption: We assume that n is exact power of 2.


$$T(n) = 2 T\left(\frac{n}{2}\right) + n^2$$

$$T(n) = 2 T\left(\frac{n}{2}\right) + n^2$$




$$T(n) = 2 T\left(\frac{n}{2}\right) + n^2$$

$$T(\frac{n}{2}) = 2.T(\frac{n}{2^2}) + \frac{n^2}{2^2}$$



$$T(n) = 2 T\left(\frac{n}{2}\right) + n^2$$


$$T(\frac{n}{2}) = 2.T(\frac{n}{2^2}) + \frac{n^2}{2^2}$$



$$T(n) = 2 T\left(\frac{n}{2}\right) + n^2$$

$$T(\frac{n}{2}) = 2.T(\frac{n}{2^2}) + \frac{n^2}{2^2}$$


$$T(\frac{n}{2^2}) = 2.T(\frac{n}{2^3}) + \frac{n^2}{4^2}$$



$$T(n) = 2 T\left(\frac{n}{2}\right) + n^2$$

$$T(\frac{n}{2}) = 2.T(\frac{n}{2^2}) + \frac{n^2}{2^2}$$

$$T(\frac{n}{2^2}) = 2.T(\frac{n}{2^3}) + \frac{n^2}{4^2}$$




$$T(n) = 2T\left(\frac{n}{2}\right) + n^2$$

$$T(\frac{n}{2}) = 2.T(\frac{n}{2^2}) + \frac{n^2}{2^2}$$

$$T(\frac{n}{2^2}) = 2.T(\frac{n}{2^3}) + \frac{n^2}{4^2}$$

$$T(\frac{n}{2^k}) = T(1)$$




$$T(n) = 2 T\left(\frac{n}{2}\right) + n^2$$

$$T(\frac{n}{2}) = 2 \cdot T(\frac{n}{2^2}) + \frac{n^2}{2^2}$$

$$T(\frac{n}{2^2}) = 2 \cdot T(\frac{n}{2^3}) + \frac{n^2}{4^2}$$

$$T(\frac{n}{2^k}) = T(1)$$




$$T(n) = 2 T\left(\frac{n}{2}\right) + n^2$$
  
 $T(\frac{n}{2}) = 2 \cdot T(\frac{n}{2^2}) + \frac{n^2}{2^2}$ 

$$T(\frac{n}{2^2}) = 2.T(\frac{n}{2^3}) + \frac{n^2}{4^2}$$

$$T(\frac{n}{2^k}) = T(1)$$

$$n = 2^k \implies k = \lg n$$



$$T(n) = 2 T\left(\frac{n}{2}\right) + n^{2}$$

$$T\left(\frac{n}{2}\right) = 2.T\left(\frac{n}{2^{2}}\right) + \frac{n^{2}}{2^{2}}$$

$$T\left(\frac{n}{2^{2}}\right) = 2.T\left(\frac{n}{2^{3}}\right) + \frac{n^{2}}{4^{2}}$$

$$T\left(\frac{n}{2^{k}}\right) = T(1)$$

$$n = 2^{k} \implies k = \lg n$$

$$L_{c} = 2^{k} \implies 2^{\lg n} \implies n^{\lg 2} \implies n$$

$$T(n) = 2 T\left(\frac{n}{2^{2}}\right) + n^{2}$$

$$n^{2}/4 \qquad n^{2}/4$$

$$n^{2}/4 \qquad n^{2}/4$$

$$n^{2}/4 \qquad n^{2}/4$$

$$n^{2}/4 \qquad n^{2}/4$$

$$n^{2}/16 \qquad n^{2}/16 \qquad n^{2}/16 \qquad n^{2}/16$$

$$T(n/8) \qquad T(n/8) \qquad T(n/8)$$

$$T(n/8) \qquad T(n/8) \qquad T(n/8)$$

$$T(n) = 2 T\left(\frac{n}{2}\right) + n^{2}$$


$$T(\frac{n}{2}) = 2 \cdot T(\frac{n}{2^{2}}) + \frac{n^{2}}{2^{2}}$$

$$T(\frac{n}{2^{2}}) = 2 \cdot T(\frac{n}{2^{3}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{2^{k}}) = T(1)$$

$$n = 2^{k} \implies k = \lg n$$

$$L_{c} = 2^{k} \implies 2^{\lg n} \implies n^{\lg 2} \implies n$$



$$T(n) = 2 T\left(\frac{n}{2}\right) + n^{2}$$


$$T(\frac{n}{2}) = 2 \cdot T(\frac{n}{2^{2}}) + \frac{n^{2}}{2^{2}}$$

$$T(\frac{n}{2^{2}}) = 2 \cdot T(\frac{n}{2^{3}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{2^{k}}) = T(1)$$

$$n = 2^{k} \implies k = \lg n$$

$$L_{c} = 2^{k} \implies 2^{\lg n} \implies n^{\lg 2} \implies n$$



$$T(n) = 2 T\left(\frac{n}{2}\right) + n^{2}$$


$$T(\frac{n}{2}) = 2 \cdot T(\frac{n}{2^{2}}) + \frac{n^{2}}{2^{2}}$$

$$T(\frac{n}{2^{2}}) = 2 \cdot T(\frac{n}{2^{3}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{2^{k}}) = T(1)$$

$$n = 2^{k} \implies k = \lg n$$

$$L_{c} = 2^{k} \implies 2^{\lg n} \implies n^{\lg 2} \implies n$$



Total Cost =  $L_c + I_c$ 

$$T(n) = 2 T\left(\frac{n}{2}\right) + n^{2}$$


$$T(\frac{n}{2}) = 2 \cdot T(\frac{n}{2^{2}}) + \frac{n^{2}}{2^{2}}$$

$$T(\frac{n}{2^{2}}) = 2 \cdot T(\frac{n}{2^{3}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{2^{k}}) = T(1)$$

$$n = 2^{k} \implies k = \lg n$$

$$L_{c} = 2^{k} \implies 2^{\lg n} \implies n^{\lg 2} \implies n$$



Total Cost =  $L_c + I_c \implies n + 2n^2$ 

$$T(n) = 2 T\left(\frac{n}{2}\right) + n^{2}$$


$$T(\frac{n}{2}) = 2 \cdot T(\frac{n}{2^{2}}) + \frac{n^{2}}{2^{2}}$$

$$T(\frac{n}{2^{2}}) = 2 \cdot T(\frac{n}{2^{3}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{2^{k}}) = T(1)$$

$$n = 2^{k} \implies k = \lg n$$

$$L_{c} = 2^{k} \implies 2^{\lg n} \implies n^{\lg 2} \implies n$$



Total Cost =  $L_c + I_c$   $\implies n + 2n^2$ 


Hence:  $T(n) \in O(n^2)$ 

$$T(n) = \begin{cases} 1 & n = 1\\ 3 T\left(\frac{n}{4}\right) + n^2 & n > 1 \end{cases}$$

Solve the following recurrence using the Recurrence Tree Method.


$$T(n) = 3 T\left(\frac{n}{4}\right) + n^2$$

$$T(n) = 3 T\left(\frac{n}{4}\right) + n^2$$




$$T(n) = 3 T\left(\frac{n}{4}\right) + n^2$$

$$T(\frac{n}{4}) = 3.T(\frac{n}{4^2}) + \frac{n^2}{4^2}$$



$$T(n) = 3 T\left(\frac{n}{4}\right) + n^2$$


$$T(\frac{n}{4}) = 3.T(\frac{n}{4^2}) + \frac{n^2}{4^2}$$



$$T(n) = 3 T\left(\frac{n}{4}\right) + n^2$$

$$T(\frac{n}{4}) = 3.T(\frac{n}{4^2}) + \frac{n^2}{4^2}$$


$$T(\frac{n}{4^2}) = 3.T(\frac{n}{4^3}) + \frac{n^2}{16^2}$$



$$T(n) = 3 T\left(\frac{n}{4}\right) + n^2$$

$$T(\frac{n}{4}) = 3.T(\frac{n}{4^2}) + \frac{n^2}{4^2}$$

$$T(\frac{n}{4^2}) = 3.T(\frac{n}{4^3}) + \frac{n^2}{16^2}$$




$$T(n) = 3 T\left(\frac{n}{4}\right) + n^2$$

$$T(\frac{n}{4}) = 3.T(\frac{n}{4^2}) + \frac{n^2}{4^2}$$

$$T(\frac{n}{4^2}) = 3.T(\frac{n}{4^3}) + \frac{n^2}{16^2}$$


$$T(\frac{n}{4^k}) = T(1)$$



$$T(n) = 3 T\left(\frac{n}{4}\right) + n^2$$
  
 $T(\frac{n}{4}) = 3.T(\frac{n}{4^2}) + \frac{n^2}{4^2}$ 

$$T(\frac{n}{4^2}) = 3.T(\frac{n}{4^3}) + \frac{n^2}{16^2}$$

$$T(\frac{n}{4^k})=T(1)$$




$$T(n) = 3 T\left(\frac{n}{4}\right) + n^2$$
  
 $T(\frac{n}{4}) = 3 \cdot T(\frac{n}{4^2}) + \frac{n^2}{4^2}$ 

$$T(\frac{n}{4^2}) = 3.T(\frac{n}{4^3}) + \frac{n^2}{16^2}$$

$$T(\frac{n}{4^k})=T(1)$$

$$n=4^k$$



$$T(n) = 3 T \left(\frac{n}{4}\right) + n^{2}$$

$$T(\frac{n}{4}) = 3.T(\frac{n}{4^{2}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{4^{2}}) = 3.T(\frac{n}{4^{3}}) + \frac{n^{2}}{16^{2}}$$

$$T(\frac{n}{4^{2}}) = 3.T(\frac{n}{4^{3}}) + \frac{n^{2}}{16^{2}}$$

$$T(\frac{n}{4^{2}}) = T(1)$$

$$T(\frac$$

$$T(n) = 3 T \left(\frac{n}{4}\right) + n^{2}$$

$$T(\frac{n}{4}) = 3. T(\frac{n}{4^{2}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{4^{2}}) = 3. T(\frac{n}{4^{3}}) + \frac{n^{2}}{16^{2}}$$

$$\frac{n^{2}}{256} \frac{n^{2}}{256} \frac{n^{2}}{256}$$

$$T(n) = 3 T \left(\frac{n}{4}\right) + n^{2}$$

$$T(\frac{n}{4}) = 3.T(\frac{n}{4^{2}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{4^{2}}) = 3.T(\frac{n}{4^{3}}) + \frac{n^{2}}{16^{2}}$$

$$\frac{n^{2}}{256} \frac{n^{2}}{256} \frac{$$

$$T(n) = 3 T\left(\frac{n}{4}\right) + n^{2}$$

$$T(\frac{n}{4}) = 3.T(\frac{n}{4^{2}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{4^{2}}) = 3.T(\frac{n}{4^{3}}) + \frac{n^{2}}{16^{2}}$$

$$\frac{n^{2}}{256} \frac{n^{2}}{256} \frac{n$$

$$T(n) = 3 T\left(\frac{n}{4}\right) + n^{2}$$

$$T(\frac{n}{4}) = 3.T(\frac{n}{4^{2}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{4^{2}}) = 3.T(\frac{n}{4^{3}}) + \frac{n^{2}}{16^{2}}$$

$$\frac{n^{2}}{256} \frac{n^{2}}{256} \frac{n$$

Total Cost =  $L_c + I_c$ 

$$T(n) = 3T\left(\frac{n}{4}\right) + n^{2}$$

$$T(\frac{n}{4}) = 3.T(\frac{n}{4^{2}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{4^{2}}) = 3.T(\frac{n}{4^{3}}) + \frac{n^{2}}{16^{2}}$$

$$T(\frac{n}{4^{2}}) = 3.T(\frac{n}{4^{3}}) + \frac{n^{2}}{16^{2}}$$

$$T(\frac{n}{4^{2}}) = T(1)$$

$$T(\frac{n$$

$$T(n) = 3 T\left(\frac{n}{4}\right) + n^{2}$$

$$T(\frac{n}{4}) = 3. T(\frac{n}{4^{2}}) + \frac{n^{2}}{4^{2}}$$

$$T(\frac{n}{4^{2}}) = 3. T(\frac{n}{4^{3}}) + \frac{n^{2}}{16^{2}}$$

$$\frac{n^{2}}{256} \frac{n^{2}}{256} \frac$$

$$T(n) = \begin{cases} 1 & n = 1\\ T\left(\frac{n}{4}\right) + T\left(\frac{n}{2}\right) + n^2 & n > 1 \end{cases}$$

Solve the following recurrence using the Recurrence Tree Method.

$$T(n) = \begin{cases} 1 & n = 1\\ T\left(\frac{n}{3}\right) + T\left(\frac{2n}{3}\right) + n & n > 1 \end{cases}$$

Solve the following recurrence using the Recurrence Tree Method.