

2021.07.28

Lecture 12



서울대학교
SEOUL NATIONAL UNIVERSITY

Department of Chemistry

PChem 2

Physical Chemistry 2

Jiho Son

Physical Chemistry 2

The Last Lecture. Polyatomic molecules

July 28th, 2021

paradichlorobenzene1724@gmail.com

kadryjh1724@snu.ac.kr

"Please, do not distribute this lecture note without permission!"

Topics in Lecture 11

MO of heteronuclear diatomic molecules

Hybrid orbitals

π -electron approximation

Brief introduction to the group theory

In Atkins' *Physical Chemistry* (11th ed.),

9D MOT: heteronuclear diatomic molecules

9E MOT: polyatomic molecules

10A Shape and symmetry

2021.07.28

Lecture 12



서울대학교
SEOUL NATIONAL UNIVERSITY

Department of Chemistry

PChem 2

Physical Chemistry 2

Jiho Son

02 / 19



MO of heteronuclear diatomics

For heteronuclear diatomic molecules, difference in *electronegativity* occurs. Therefore, we cannot just set $c_A = c_B$ simply. Therefore, we need to find the optimizing value of coefficients.

Consider a heteronuclear diatomic molecule with atom A and B. Then, wavefunction is:

$$\psi = c_A\psi_a + c_B\psi_B$$

Then, energy calculated from this wavefunction is

$$E = \frac{\int \psi \hat{H} \psi d\tau}{\int \psi^2 d\tau} = E(c_A, c_B)$$

Denominator:

$$\int \psi^2 d\tau = \int (c_A\psi_A + c_B\psi_B)(c_A\psi_A + c_B\psi_B) d\tau = c_A^2 + c_B^2 + 2c_Ac_BS$$



MO of heteronuclear diatomics

Numerator:

$$\begin{aligned}
 \int \psi \hat{H} \psi d\tau &= \int (c_A \psi_A + c_B \psi_B) \hat{H} (c_A \psi_A + c_B \psi_B) d\tau \\
 &= c_A^2 \int \psi_A \hat{H} \psi_A d\tau + c_B^2 \int \psi_B \hat{H} \psi_B d\tau + c_A c_B \int \psi_A \hat{H} \psi_B d\tau + c_B c_A \int \psi_B \hat{H} \psi_A d\tau \\
 &\equiv c_A^2 H_{AA} + c_B^2 H_{BB} + 2c_A c_B H_{AB}
 \end{aligned}$$

Therefore, we have

$$E \cdot (c_A^2 + c_B^2 + 2c_A c_B S) = c_A^2 H_{AA} + c_B^2 H_{BB} + 2c_A c_B H_{AB}$$

Differentiate both side with respect to c_A and c_B .



MO of heteronuclear diatomics

Therefore

$$E \cdot (c_A^2 + c_B^2 + 2c_Ac_BS) = c_A^2H_{AA} + c_B^2H_{BB} + 2c_Ac_BH_{AB}$$

$$\frac{\partial E}{\partial c_A}(c_A^2 + c_B^2 + 2c_Ac_BS) + E(2c_A + 2c_BS) = 2c_AH_{AA} + 2c_BH_{AB} \implies \frac{\partial E}{\partial c_A} = 0, c_A(H_{AA} - E) + c_B(H_{AB} - ES) = 0$$

$$\frac{\partial E}{\partial c_B}(c_A^2 + c_B^2 + 2c_Ac_BS) + E(2c_B + 2c_AS) = 2c_BH_{BB} + 2c_AH_{AB} \implies \frac{\partial E}{\partial c_B} = 0, c_A(H_{AB} - ES) + c_B(H_{BB} - E) = 0$$

We yield *secular equations*.

$$c_A(H_{AA} - E) + c_B(H_{AB} - ES) = 0$$

$$c_A(H_{AB} - ES) + c_B(H_{BB} - E) = 0$$

Note that $c_A = c_B = 0$ is a trivial solution of secular equation. To require that secular equations have nontrivial solutions, we need

$$\begin{vmatrix} H_{AA} - E & H_{AB} - ES \\ H_{AB} - ES & H_{BB} - E \end{vmatrix} = 0$$



MO of heteronuclear diatomics

From the secular equation, we yield the energy eigenvalues.

For the simplification, let $S = 0$ (complete neglect of overlap). Then eigenvalues are


$$E_{\pm} = \frac{H_{AA} + H_{BB}}{2} \pm \frac{H_{AA} - H_{BB}}{2} \sqrt{1 + \left(\frac{2H_{AB}}{H_{AA} - H_{BB}} \right)^2}$$

Coefficients corresponding to each eigenvalues can be calculated. Do it yourself!




Hybrid orbitals

The ground state electron configuration of a carbon atom,



does not seem to lead to the tetrahedral bonding in methane and other saturated hydrocarbons. This problem is overcome with *promotion*.



After the promotion, *hybridization* follows.



sp orbitals

Consider first the linear molecule, BeH_2 . We know that this molecule has two equivalent Be-H bonds, with bond angle of 180° . However, BeH_2 has electronic configuration of $1s^2 2s^2$.

To counter this problem, we *mix* p_z orbital, to form *sp* hybrid orbital.

The two *sp* orbitals are equivalent, and are directed 180° from each other.

$$\psi_{sp} = \frac{1}{\sqrt{2}}(2s \pm 2p_z)$$




FIGURE 10.1

An illustration of the *sp* hybrid orbitals. The two *sp* hybrid orbitals are equivalent and are directed 180° from each other. The two remaining $2p$ orbitals are perpendicular to each other and to the line formed by the *sp* orbitals.

sp^2 orbitals

Same approach can be applied to BH_3 . To make three equivalent bonds directed 120° from each other, we need to mix p_x and p_y orbitals with s orbital.

$$\psi_1 = \frac{1}{\sqrt{3}}2s + \sqrt{\frac{2}{3}}2p_y$$

$$\psi_2 = \frac{1}{\sqrt{3}}2s + \frac{1}{\sqrt{2}}2p_x - \frac{1}{\sqrt{6}}2p_y$$

$$\psi_3 = \frac{1}{\sqrt{3}}2s - \frac{1}{\sqrt{2}}2p_x - \frac{1}{\sqrt{6}}2p_y$$




FIGURE 10.4

The geometry associated with sp^2 orbitals. The three sp^2 orbitals lie in a plane and point to the vertices of an equilateral triangle. The remaining $2p$ orbital is perpendicular to the plane formed by the three sp^2 orbitals.

These orbitals are not solutions of Schrödinger equation: hybrid orbitals are VBT concept, and they are custom-made to describe molecules we cannot deal in VBT.

sp^3 orbitals


For CH_4 , we need four orbitals pointing tetrahedral points.

$$\psi_1 = \frac{1}{2}(2s + 2p_x + 2p_y + 2p_z)$$

$$\psi_2 = \frac{1}{2}(2s - 2p_x - 2p_y + 2p_z)$$

$$\psi_3 = \frac{1}{2}(2s + 2p_x - 2p_y - 2p_z)$$

$$\psi_4 = \frac{1}{2}(2s - 2p_x + 2p_y - 2p_z)$$



We have learned that PES (PhotoElectron Spectroscopy) can blow a cover of electronic structure of certain molecule. However, PES result does not coincide with hybrid orbital results.

Therefore, we need MOT to describe polyatomic molecules qualitatively.

2021.07.28

Lecture 12



서울대학교
SEOUL NATIONAL UNIVERSITY

Department of Chemistry

PChem 2

Physical Chemistry 2

Jiho Son

sp^3 orbitals




FIGURE 10.15
A molecular-orbital energy-level diagram for the valence electrons in CH_4 .




FIGURE 10.16
The photoelectron spectrum of methane. The two lines observed in the photoelectron spectrum reflect the ionization of electrons from the $1t_2$ and $2a_1$ molecular orbitals. The energy difference between these two lines corresponds to the energy difference between the $1t_2$ and $2a_1$ molecular orbitals (see Figure 10.15). The lines are broad because ionization occurs from many different vibrational levels of the molecules.

2021.07.28

Lecture 12



서울대학교
SEOUL NATIONAL UNIVERSITY

Department of Chemistry

PChem 2


Physical Chemistry 2

Jiho Son



π -electron approximation

Unsaturated hydrocarbons can be treated simply, by separating σ bonds and π bonds.



σ bonds are relatively more localized between nuclei, than π bonds. Therefore, we ignore *σ -bond framework*, and only consider π electrons. This is π -electron approximation.

For example, consider the simplest conjugated π system, ethylene. Only remaining π contribution is from two $2p_z$ orbitals.

$$\psi_{\pi} = c_1 2p_{zA} + c_2 2p_{zB}$$

π -electron approximation

We can calculate secular determinant associated with that wavefunction.

$$\psi_{\pi} = c_1 2p_{zA} + c_2 2p_{zB}$$
$$\begin{vmatrix} H_{11} - ES_{11} & H_{12} - ES_{12} \\ H_{12} - ES_{12} & H_{22} - ES_{12} \end{vmatrix} = 0$$

where H_{11} and H_{22} defined same as previous problem. They are called *Coulomb integral*.
The off-diagonal element of Hamiltonian, H_{12} , is called *exchange integral*.

Here, we examine an approximation proposed by Erich Hückel in 1930.

2021.07.28

Lecture 12



서울대학교
SEOUL NATIONAL UNIVERSITY

Department of Chemistry

PChem 2

Physical Chemistry 2

Jiho Son



Hückel approximation

There are three simple assertions of Hückel molecular orbital theory.

[1] Overlap integrals, $S_{ij} = \delta_{ij}$. i.e., ignore all overlaps between different atoms.

[2] All Coulomb integrals are assumed to be the same for all carbon atoms, $H_{11} = H_{22} = \alpha$.

[3] The exchange integrals involving nearest-neighbor carbon atoms are assumed to be the same, and are denoted by $H_{12} = \beta$. If two carbon are not adjacent, $H_{ij} = 0$.

$$\begin{vmatrix} \alpha - E & \beta \\ \beta & \alpha - E \end{vmatrix} = 0 \implies E = \alpha \pm \beta$$

Since α approximates the energy of an electron in an isolated $2p$ orbital, we use it as a reference of energy. Note that $\beta < 0$ in general. Therefore, ground state is $E = \alpha + \beta$.



서울대학교
SEOUL NATIONAL UNIVERSITY

Department of Chemistry

PChem 2

Physical Chemistry 2

Jiho Son

Hückel approximation

For ethylene,




FIGURE 10.20

The ground-state electron configuration of the π electrons in ethene.

We can do similar thing with butadiene molecule.





FIGURE 10.21

A schematic representation of the $2p_z$ orbitals of each of the carbon atoms in a butadiene molecule.



서울대학교
SEOUL NATIONAL UNIVERSITY

Department of Chemistry

PChem 2

Physical Chemistry 2

Jiho Son

Hückel approximation

Write down the secular determinant.

$$\begin{vmatrix} \alpha - E & \beta & 0 & 0 \\ \beta & \alpha - E & \beta & 0 \\ 0 & \beta & \alpha - E & \beta \\ 0 & 0 & \beta & \alpha - E \end{vmatrix} = 0$$

By letting $x = (\alpha - E)/\beta$, we yield

$$\beta^4 \begin{vmatrix} x & 1 & 0 & 0 \\ 1 & x & 1 & 0 \\ 0 & 1 & x & 1 \\ 0 & 0 & 1 & x \end{vmatrix} = 0 \implies x = \pm 1.618, \pm 0.618$$




FIGURE 10.22

The ground-state electron configuration of the π electrons in butadiene.



Delocalization energy

Since butadiene has four π electrons, total energy of π electrons is

$$E_{\pi} = 2(\alpha + 1.618\beta) + 2(\alpha + 0.618\beta) = 4\alpha + 4.472\beta$$

Compared to energy of two ethylene molecules ($2E = 4\alpha + 4\beta$), energy of butadiene is lower.

$$E_{\text{deloc}} = E_{\pi}(\text{butadiene}) - 2E_{\pi}(\text{ethylene}) = 0.472\beta < 0$$

This extra stabilization is called *delocalization energy*.




FIGURE 10.23

A schematic representation of the π molecular orbitals of butadiene. Note that the corresponding energy increases with the number of nodes.

Hückel approximation

You can do the same thing with benzene molecule.

$$\begin{aligned}
 \psi_1 &= \frac{1}{\sqrt{6}}(2p_{z1} + 2p_{z2} + 2p_{z3} + 2p_{z4} + 2p_{z5} + 2p_{z6}) & E_1 &= \alpha + 2\beta \\
 \psi_2 &= \frac{1}{\sqrt{4}}(2p_{z2} + 2p_{z3} - 2p_{z5} - 2p_{z6}) & E_2 &= \alpha + \beta \\
 \psi_3 &= \frac{1}{\sqrt{3}}(2p_{z1} + \frac{1}{2}2p_{z2} - \frac{1}{2}2p_{z3} - 2p_{z4} - \frac{1}{2}2p_{z5} + \frac{1}{2}2p_{z6}) & E_3 &= \alpha + \beta \\
 \psi_4 &= \frac{1}{\sqrt{4}}(2p_{z2} - 2p_{z3} + 2p_{z5} - 2p_{z6}) & E_4 &= \alpha - \beta \\
 \psi_5 &= \frac{1}{\sqrt{3}}(2p_{z1} - \frac{1}{2}2p_{z2} - \frac{1}{2}2p_{z3} + 2p_{z4} - \frac{1}{2}2p_{z5} - \frac{1}{2}2p_{z6}) & E_5 &= \alpha - \beta \\
 \psi_6 &= \frac{1}{\sqrt{6}}(2p_{z1} - 2p_{z2} + 2p_{z3} - 2p_{z4} + 2p_{z5} - 2p_{z6}) & E_6 &= \alpha - 2\beta
 \end{aligned} \tag{10.31}$$





FIGURE 10.24

The ground-state electron configuration of the π electrons in benzene.



서울대학교
SEOUL NATIONAL UNIVERSITY

Department of Chemistry

PChem 2

Physical Chemistry 2

Jiho Son

Group theory

For the small molecules, considering symmetry can help you solving electronic problems.

If you are interested in symmetry and its interesting consequences, attend *Inorganic Chemistry 1 & 2* courses.

This is the end of **SNU CNS summer major school 2021, Physical Chemistry 2 course.**

Stay safe at home, wash your hands, and avoid social contacts.

I hope this course can help you in your fall semester.

2021.07.28

Lecture 12



서울대학교
SEOUL NATIONAL UNIVERSITY

Department of Chemistry

PChem 2

Physical Chemistry 2

Jiho Son