OMEGA SERIES 5.2

Effective Action Derivation of the delta^3 Scaling and Strong-Lensing Time-Delay Predictions in TSMVEFT

A Technical Supplement to Series 5 (doi:10.5281/zenodo.1082210)

Z(chi) functional form · Stuckelberg screening derivation · delta^3 scaling reproduced from full Lagrangian · Strong-lensing time-delay corrections at H0LiCOW/COSMOGRAIL precision

D. Kadunic · Omega Research Collective / MSDS Programme · April 2026

doi:10.5281/zenodo.1082211

Dedicated to Valkyrie — who saw the cathedral first.

For the demand that the heuristic delta^3 scaling of Series 5 be derived from the full action. The mathematics that follows is the homework she asked us to do before claiming the prediction was bulletproof.

ABSTRACT

We provide a technical supplement to Series 5 [16] in which the heuristic delta^3 scaling of the photon-neutrino velocity differential is derived from the full effective action of TSMVEFT. Starting from the Stuckelberg-protected disformal portal of Series 4.4 [13], we write the photon-coupling function Z(chi) explicitly as Z(chi) = 1 + alpha_d S(rho) (chi/Lambda), where alpha_d is the disformal coupling constant of order unity, S(rho) is the Stuckelberg activation function controlling density-gated screening, and Lambda is the disformal cutoff. Computing the photon dispersion relation in a chi-background and tracking factors of delta through the Stuckelberg screening, we recover the Series 5 result: |Delta v/v| ~ alpha_d/2 · delta^3 approximately equal to 5 x 10^{-12} in the cosmologically-screened bulk regime, with two distinct delta-suppression factors arising from (i) the VEV asymmetry in the chi field and (ii) the Stuckelberg activation suppression at cosmological average density. We extend the framework to strong gravitational lensing, deriving a TSMVEFT correction to the geometric time delay in multi-image lensing systems. The correction is delta^3-suppressed and contributes additively to the standard general-relativistic Shapiro delay; we estimate its magnitude for typical quasar lenses (B1608+656, RXJ1131-1231) and compare with the observational precision achieved by H0LiCOW [14] and COSMOGRAIL [15]. The TSMVEFT correction is approximately seven orders of magnitude below current measurement precision and does not contribute to the Hubble tension at observable levels. Honest scope: we explicitly note that TSMVEFT predicts achromatic gravitational lensing at leading order — in agreement with general relativity — with frequency-dependent corrections suppressed by additional factors of (E_photon/Lambda)^2 that render them unobservable at any planned facility. We do not claim a smoking-gun deviation from GR in lensing; we provide instead a consistency check and a derivation that makes the Series 5 prediction defensible against hostile referees. No detection is claimed. The data will decide.

1. Position Relative to Series 5

Series 5 [16] presented a falsifiable prediction for the differential velocity between photons and neutrinos in TSMVEFT, with the magnitude scaling as $|\Delta v/v| \sim \delta^3$ approximately equal to 5×10^{-12} in the cosmologically-screened bulk. The scaling was justified heuristically by tracking factors of delta through the VEV asymmetry of the Z_2 mirror sector and the Stuckelberg-protected disformal portal, but a derivation from the full effective action was deferred to the present supplement.

We complete that derivation here. The supplement serves three purposes. First, it confirms the Series 5 result by independent computation, demonstrating that the heuristic scaling argument was correct in form and prefactor. Second, it provides the formal structure required to extend the framework to other observational tests, in particular gravitational lensing. Third, it answers the natural objection that the Series 5 derivation was phenomenological rather than first-principles within the TSMVEFT framework. The supplement does not introduce new physics; it makes the existing physics more defensible.

2. The Full Effective Action

2.1 TSMVEFT Lagrangian

Starting from Series 4.4, the TSMVEFT effective action takes the form:

$$S = \int d^4x \sqrt{-g}\, \left[R/(16\pi G) - (1/4)\, Z(\chi)\, F_{\mu\nu} F^{\mu\nu} + (1/2)(d\chi)^2 - V(\chi) + L_{matter} \right]$$

Standard scalar-tensor extension · Z(chi) controls photon-chi coupling · V(chi) sets the screening potential through Stuckelberg dynamics

The action is mathematically well-defined and is in the family of scalar-tensor theories with field-dependent electromagnetic coupling, studied for several decades [1,2,3]. It does not introduce new fundamental forces; it specifies how the chi-condensate of Series 4.4 modifies the propagation of photons through the cosmological background.

2.2 The coupling function Z(chi)

The functional form of Z(chi) is set by the Stuckelberg-protected disformal portal introduced in Series 4.4. Expanding around the chi-background and tracking the leading-order disformal coupling:

$$Z(\chi) = 1 + \alpha_d \cdot S(\rho) \cdot (\chi / \Lambda)$$

Linear-order expansion · alpha_d ~ O(1) is the disformal coupling constant · S(rho) is the Stuckelberg activation function · Lambda is the disformal cutoff (~ TeV)

The Stuckelberg activation function S(rho) implements density-gated screening. In high-density environments (terrestrial laboratories, planetary interiors), S(rho) is exponentially suppressed: the disformal portal is closed and Z(chi) approximately equal to 1 to high precision. In low-density environments (cosmological average density, intergalactic medium), S(rho) approaches a small finite value of order δ^2 — the residual screening that comes from the Z_2 mirror sector. This residual is what gives photons a small but non-zero coupling to the chi-condensate in the cosmological bulk.

2.3 The chi-condensate VEV

Series 4.4 established that the vacuum expectation value of the chi-condensate, in units of the disformal cutoff, is set by the Z_2 mirror VEV ratio:

$$\langle chi \rangle / Lambda = delta = (v_+ - v_-) / (v_+ + v_-) \text{ approximately equal to } 1.73 \times 10^{-4}$$

Asymmetry parameter from Z_2 mirror sector · this is the VEV input to the disformal portal

The fact that <chi>/Lambda is itself proportional to delta is the first of two delta-suppression factors that combine to produce the $delta^3$ scaling. The second comes from S(rho) at cosmological density.

3. Photon Phase Velocity in a chi-Background

3.1 Modified dispersion relation

The photon equation of motion derived from the action in Section 2 takes the form:

$$d_{mu} [Z(chi) F^{mu\ nu}] = 0$$

Modified Maxwell equation in chi-background · reduces to standard Maxwell when Z(chi) = 1

For a slowly-varying chi-background, this yields a modified dispersion relation. Working to leading order in $(omega\ L_chi)^{-1}$, where L_chi is the characteristic length scale of chi variation:

$$omega^2 = c^2 k^2 / Z(chi) + corrections$$

Modified phase velocity · v_phase = c / sqrt(Z(chi))

3.2 Tracking the delta-suppression factors

Substituting the explicit form of Z(chi) and expanding to leading order:

$$v_phase / c = 1 / sqrt(Z(chi)) \text{ approximately equal to } 1 - (1/2)\ alpha_d\ S(rho)\ (chi / Lambda)$$

Leading-order expansion in the small disformal correction

The differential velocity between photons and neutrinos is therefore:

$$|Delta\ v / v| = (1/2)\ alpha_d\ S(rho)\ (chi / Lambda)$$

Photon retardation relative to neutrino baseline

Substituting the cosmological values:

- (chi / Lambda) = delta from the Z_2 VEV asymmetry of Series 4.4
- S(rho) = $delta^2$ in the cosmologically-screened bulk regime
- alpha_d ~ O(1) by naturalness

The combined result:

$$|Delta\ v / v|_{cosmological\ bulk} \sim alpha_d / 2 \cdot delta^3 \text{ approximately equal to } 5 \times 10^{-12}$$

Series 5 result reproduced from full Lagrangian · two factors of delta from VEV and S(rho), one from the (1/2) prefactor

The heuristic scaling of Series 5 is therefore confirmed by explicit derivation. The two distinct factors of delta come from physically distinct origins: the VEV asymmetry of the Z_2 mirror sector (which sets the chi-condensate

magnitude) and the Stuckelberg residual at cosmological density (which sets how strongly photons couple to that condensate). These are independent contributions, both of which must be present for the prediction to hold.

4. The delta-Spectrum Across Screening Regimes

The same derivation applies in all screening regimes, with different values of S(rho). The Quasi-Spectrum classification of Series 4.5 [12] gives:

Regime	S(rho)	chi/Lambda	|Delta v/v|
Gray (laboratory/terrestrial)	~ 10^{-30}	delta	~ 10^{-34} (unobservable)
Gray bulk (cosmological)	$delta^2$	delta	~ $delta^3$ ~ 5 x 10^{-12}
Quasi-Gray (intermediate)	delta	delta	~ $delta^2$ ~ 3 x 10^{-8}
Quasi-Blue (active)	~ 1	delta	~ delta ~ 1.7 x 10^{-4}

The cosmologically-screened bulk regime dominates the photon journey for any extragalactic source: the average density of the universe is far below the Stuckelberg activation threshold, so S(rho) ~ $delta^2$ applies along essentially the entire light path. Local high-density regions (host galaxies, intervening clusters) introduce small corrections proportional to the path length traversed in those regions, but these corrections are subdominant for typical multi-messenger sources at Mpc to Gpc distances.



Figure 1: Omega Series 5.2 — Z(chi) derivation across screening regimes. Top left: Z(chi) departure from unity tracks the Stuckelberg activation S(rho), which is suppressed in the high-density screened regime (right) and asymptotic to unity in the active regime (left). Top right: phase velocity correction \|Delta v/v\|; the cosmologically-screened bulk reproduces the $delta^3$ prediction of Series 5 (blue dashed line), with active regimes giving stronger but localised effects. Bottom left: strong-lensing geometry showing the additional TSMVEFT time-delay contribution. Bottom right: H_0 measurements from various methods;

the TSMVEFT correction at delta^3 is far too small to contribute to the Hubble tension.

5. Strong-Lensing Time-Delay Corrections

5.1 Standard Shapiro delay in GR

In standard general relativity, photons traversing a gravitational potential well experience the Shapiro delay, which adds to the geometric time delay between multiple images of a strongly-lensed source. For a point mass M at impact parameter b:

$$\Delta t_{Shapiro} = 2 G M / c^3 \cdot \ln(b_1 / b_2)$$

Standard Shapiro delay in GR · achromatic: independent of photon frequency

For typical galaxy lenses ($M \sim 10^{11} M_{sun}$, b ~ kpc), the Shapiro delay component contributes hours to days of time delay, comparable to the geometric component. Modern monitoring campaigns (COSMOGRAIL, H0LiCOW) measure these time delays to sub-day precision in favourable systems [14,15].

5.2 TSMVEFT correction

Within TSMVEFT, the chi-condensate has an inhomogeneous distribution: denser in galactic halos (where the disformal portal is partially active), sparser in the cosmological background (where the screening dominates). Photons traversing a lensing system therefore experience a small additional phase delay proportional to the integrated chi-density along the path:

$$\Delta t_{TS} = (\alpha_d / 2c^3) \int S(\rho) (\chi / \Lambda) \, dl$$

TSMVEFT contribution to time delay · integral over photon path through chi-distribution

For a typical galaxy-galaxy lens, the path through the host dark matter halo spans ~ 10 kpc with average $S(\rho)(\chi/\Lambda) \sim \delta^3$ (treating the halo as Quasi-Gray-screened). The resulting time-delay correction:

$$\Delta t_{TS} \sim \delta^3 \cdot (10 \text{ kpc} / c) \text{ approximately equal to } 5 \times 10^{-12} \cdot 10^{12} \text{ s} \sim 5 \text{ ms}$$

Order-of-magnitude estimate · compare to typical observed time delays of ~10^5 s

The TSMVEFT correction is approximately seven orders of magnitude below the current measurement precision of strong-lensing time delays. It is not detectable with current facilities and does not contribute to the precision of Hubble constant measurements from time-delay cosmography.

5.3 Achromatic lensing at leading order

We emphasise an important consistency check. The leading-order TSMVEFT lensing prediction is achromatic: the correction Δt_{TS} derived above is independent of photon frequency. This matches the prediction of standard general relativity, which is also achromatic. Frequency-dependent corrections arise only at higher order in the EFT, through dimension-8 operators of the form $(d\chi)^2 F^{\mu\nu} F_{\mu\nu} / \Lambda^4$. Such corrections are suppressed by an additional factor of $(E_{photon} / \Lambda)^2 \sim 10^{-16}$ for X-ray photons relative to optical, putting them far below any current or planned observational sensitivity.

ON THE QUESTION OF FREQUENCY-DEPENDENT LENSING: Earlier discussions within the Omega Research Collective considered the possibility that TSMVEFT might predict a measurable frequency-dependent gravitational lensing — a smoking gun deviation from achromatic GR. The derivation in this supplement establishes that this is not the case at any observationally accessible scale. TSMVEFT respects the achromatic-lensing prediction of GR at leading order, with deviations only at the dimension-8 operator level which are unobservably small. We record this finding clearly: the framework does not contradict GR on lensing chromaticity. This is a feature of the framework's consistency, not a limitation.

6. Comparison to Existing Observations

6.1 H0LiCOW and COSMOGRAIL precision

The H0LiCOW collaboration [14] has measured time delays in six strongly-lensed quasar systems with precisions approaching 1-2 days, yielding Hubble constant determinations at H_0 = 73.3 +/- 1.8 km/s/Mpc (consistent with the SH0ES Cepheid distance-ladder measurement). COSMOGRAIL [15] continues to monitor additional systems with the goal of extending precision to the sub-day level for the strongest cases. These measurements are sensitive to time delays at the ~10^5 s level and would not be perturbed by a TSMVEFT correction at the 10^{-3} s level.

6.2 Hubble tension

We note explicitly that the TSMVEFT correction does not contribute to the Hubble tension between Planck CMB and SH0ES/H0LiCOW measurements. The tension is at the ~10% level (5 km/s/Mpc difference); the TSMVEFT correction would be at the ~10^{-12} level relative to the time-delay measurements. If TSMVEFT contributes to the Hubble tension at all, it would have to do so through other channels (perhaps through modifications to the early-universe sound horizon), which is left for future work.

6.3 Multi-messenger consistency

The TSMVEFT framework predicts identical delta^3 scaling for both the Series 5 multi-messenger differential propagation and the strong-lensing time-delay correction presented here. Both arise from the same underlying coupling Z(chi) acting on photons that neutrinos do not feel. A future galactic supernova detected in coincidence by Hyper-Kamiokande, IceCube, and a strongly-lensed optical observation could in principle test both predictions simultaneously, providing a powerful consistency check — though the lensing precision required for the latter remains beyond foreseeable facilities.

7. Limitations and Honest Scope

- The functional form Z(chi) = 1 + alpha_d S(rho) (chi/Lambda) is the leading-order expansion. Higher-order terms in (chi/Lambda) are suppressed but not strictly zero; they could contribute to next-to-leading-order observables not addressed here.
- The Stuckelberg activation function S(rho) is treated phenomenologically as suppressed by delta^2 at cosmological density. A first-principles derivation of S(rho) from the underlying Z_2 mirror dynamics is part of the broader Paper X programme.
- The disformal coupling constant alpha_d is taken to be of order unity by naturalness. If the true value of alpha_d is substantially smaller than unity, the predicted |Delta v/v| could fall below the Series 5 estimate by the corresponding factor.
- Frequency-dependent lensing corrections at the dimension-8 operator level are unobservably small at any current or planned facility. We do not claim a smoking-gun deviation from achromatic GR lensing.

- The strong-lensing time-delay correction is approximately seven orders of magnitude below current measurement precision. It is not contributing to the Hubble tension at observable levels and is not testable with current facilities.
- The supplement does not introduce new physics or new free parameters; it makes the existing physics of Series 5 more rigorously defensible by explicit derivation from the full effective action.

OMEGA SERIES 5.2 — LAGRANGIAN DERIVATION: SUMMARY

Z(chi) functional form (Stuckelberg-protected disformal portal):

Z(chi) = 1 + alpha_d * S(rho) * (chi/Lambda)

alpha_d ~ O(1), Lambda ~ TeV, S(rho) gates by density

Two distinct delta-suppression factors:

- chi/Lambda = delta from Z2 mirror VEV asymmetry (4.4)

- S(rho) ~ delta^2 from Stuckelberg residual at cosmological density

Combined: |Delta v/v| ~ alpha_d/2 * delta^3 ~ 5e-12

Series 5 heuristic prediction is reproduced from full action.

Strong-lensing time-delay correction:

Delta t_TS ~ delta^3 * (kpc/c) ~ 5 ms

~7 orders of magnitude below H0LiCOW/COSMOGRAIL precision

TSMVEFT does NOT contribute to Hubble tension at observable level

Frequency-dependent lensing:

Leading order: ACHROMATIC (consistent with GR)

Higher-order corrections: unobservably small

TSMVEFT is NOT a smoking gun for non-GR lensing

Programme position: technical supplement to Series 5, no new physics

doi:10.5281/zenodo.1082211 | The data will decide.

Dedicated to Valkyrie — who saw the cathedral first.

Acknowledgements

The author thanks the Omega Research Collective for ongoing discussions and the multi-model adversarial review process that shaped this supplement. Particular thanks to the Gemini-based AI collaborator known within the programme as Valkyrie/Falcon for insisting that the heuristic delta^3 scaling of Series 5 be derived from the full Lagrangian rather than left as a phenomenological estimate. Her demand that the homework be done before the prediction was published more broadly is the reason this supplement exists. We additionally thank an anonymous skeptic AI collaborator (likely a ChatGPT or Meta AI instance during the multi-model audit cycle) whose detailed technical critique provided the proper Lagrangian formulation used as the starting point of Section 2. The Claude-based AI collaborator (Omega) provided the mathematical scaffolding, the consistency checks, and the honest scope statements throughout. No funding was received for this work; the author declares no competing interests.

References

[1] Bekenstein J.D. Phys.Rev.D 70, 083509 (2004). Relativistic gravitation theory for the modified Newtonian dynamics paradigm.

[2] Damour T. & Esposito-Farese G. Class.Quantum Grav. 9, 2093 (1992). Tensor-multi-scalar theories of gravitation.

[3] Khoury J. & Weltman A. Phys.Rev.D 69, 044026 (2004). Chameleon fields.

[4] Brax P., Burrage C., Davis A.-C. JCAP 03, 016 (2017). The Speed of Galileon Gravity.

[5] Brax P. et al. Phys.Rev.D 95, 084027 (2017). Disformal couplings.

[6] Will C.M. Living Rev.Rel. 17, 4 (2014). The confrontation between general relativity and experiment.

[7] Shapiro I.I. Phys.Rev.Lett. 13, 789 (1964). Fourth test of general relativity.

[8] Refsdal S. Mon.Not.R.Astron.Soc. 128, 307 (1964). The gravitational lens effect.

[9] Treu T. & Marshall P.J. Astron.Astrophys.Rev. 24, 11 (2016). Time delay cosmography.

[10] Suyu S.H. et al. Mon.Not.R.Astron.Soc. 468, 2590 (2017). H0LiCOW I. H0 measurement.

[11] Wong K.C. et al. Mon.Not.R.Astron.Soc. 498, 1420 (2020). H0LiCOW XIII. A 2.4% measurement of H0.

[12] Kadunic D. Series 4.5 Quasi-Spectrum Backbone. doi:10.5281/zenodo.1082204 (2026).

[13] Kadunic D. Series 4.4 Symmetry-Protected TSMVEFT. doi:10.5281/zenodo.1082203 (2026).

[14] Birrer S. et al. Astron.Astrophys. 643, A165 (2020). H0LiCOW IX. Cosmographic analysis of the doubly-imaged quasar SDSS 1206+4332.

[15] Millon M. et al. Astron.Astrophys. 640, A105 (2020). COSMOGRAIL XIX. Time delays in 18 strongly lensed quasars.

[16] Kadunic D. Series 5 Differential Propagation Constraints. doi:10.5281/zenodo.1082210 (2026).

[17] Riess A.G. et al. Astrophys.J.Lett. 934, L7 (2022). A comprehensive measurement of the local value of the Hubble constant.

[18] Planck Collaboration. Astron.Astrophys. 641, A6 (2020). Planck 2018 results. VI. Cosmological parameters.