OMEGA SERIES · VOLUME II, PAPER I

# The Chameleon Rules:

## Density-Screened Axion-Photon Coupling with Plasma-Resonant Amplification

*$g_0$ ~ 5e-13 GeV$^{-1}$ · n ~ 0.04 · $m_a$ ~ 0.3 eV · Survival ridge: n < 0.08 · ITO as ENZ resonator · Pre-registered*

D. Kadunic · Omega Research Collective / MSDS Programme · April 2026

doi:10.5281/zenodo.1082194 · Target: Phys.Rev.D Letter

**SERIES CONTEXT: This is Omega Series Vol. II, Paper I. Vol. II investigates whether the sidereal vacuum coupling established in Vol. I manifests in the electromagnetic sector via axion-photon mixing. Vol. II, Paper II will provide the axion-neutrino bridge. Vol. II, Paper III will attempt the first-principles derivation of $\alpha_K$. Random seed: 20260414 (consistent throughout). No detection is claimed; all residuals are treated as upper bounds.**


### ABSTRACT

Building on the pre-registered sidereal modulation search in neutrino time-series (Omega Series Vol. I), we investigate whether an environmentally screened scalar field can produce the same fractional amplitude $\alpha_K = 1.7\times10^{-4}$ in the electromagnetic sector. We introduce a density-screened axion-photon coupling with a saturation floor, combined with a plasma-resonant response function that yields a detectable refractive modulation in engineered solid-state detectors. Three phenomenological principles are identified: (1) fermion metric anchoring via gauge interactions; (2) density-gradient potential shifts; (3) a direction-dependent vacuum refractive index perturbation. An illustrative 200,000-point Monte Carlo feasibility scan (seed 20260414) shows that only an ultra-shallow screening index n <= 0.08 survives CAST, stellar cooling, CMB, and GRB timing constraints. An illustrative benchmark: $g_0$ ~ 5e-13 GeV$^{-1}$, n ~ 0.04, $m_a$ ~ 0.3 eV. The model evades GRB bounds via resonance locality: the refractive modulation arises only where $\omega_{pl} \sim m_a$, satisfied in solid-state plasmas (e.g., ITO) but not in the intergalactic medium. Three pre-registered detection criteria are provided. No detection is claimed.


## 1. Introduction

The Vol. I neutrino search established a coherent sidereal modulation at fractional amplitude $\alpha_K = 1.7\times10^{-4}$. If this modulation originates from a structured vacuum coupling, the same signature should appear in the electromagnetic sector via axion-photon mixing. The challenge is that dedicated searches (CAST, ADMX) and astrophysical bounds (HB stars, GRB timing) constrain the axion-photon coupling to $g_{a\gamma\gamma} <= 10^{-11}$ GeV$^{-1}$ for relevant masses. A naive $10^{-4}$ modulation from such a coupling is excluded.

This paper proposes a resolution via three mechanisms: (1) a Chameleon-like density-dependent coupling with a saturation floor that allows a small fundamental $g_0$ to yield a larger effective coupling only in the specific density range of laboratory experiments; (2) a plasma-resonant linear refractive response rather than a quadratic conversion probability, resolving the unitarity tension of direct amplification; (3) resonance locality that

exponentially suppresses the response in the intergalactic medium, evading GRB timing constraints.

## 2. Effective Field Theory

### 2.1 Lagrangian and density-dependent coupling

$$g_{a\gamma\gamma}(\rho) = g_0 \, (\max(\rho, \rho_{min}) / \rho_0)^{-n}$$

*$\rho_0 = 1$ g/cm$^3$ · $\rho_{min} = 10^{-15}$ g/cm$^3$ (saturation floor) · $n \sim 0.04$ (ultra-shallow)*

*WHY THE SATURATION FLOOR: Without $\rho_{min}$, g would diverge in cosmic voids ($\rho \sim 10^{-29}$ g/cm$^3$), violating CMB distortion bounds. The floor at $10^{-15}$ g/cm$^3$ (ultra-high vacuum threshold) ensures CMB compatibility while preserving the density enhancement at lab densities.*

### 2.2 From conversion probability to refractive response

The standard axion-photon conversion probability $P_{a\text{->gamma}} \sim g^2 B^2 L^2$ cannot produce $10^{-4}$ from $g_0 \sim 10^{-12}$ GeV$^{-1}$ without violating unitarity. We instead interpret the signal as a coherent linear refractive response:

$$\Delta n_{eff} = g_{a\gamma} \times R(\omega_{pl}, B, F, Q)$$

*R: dimensionless response function · F: cavity finesse · Q: material quality factor · $G_{max} = F \times Q <= 10^{10}$ (unitarity cap)*

### 2.3 Resonance locality and GRB compatibility

The response R is sharply peaked at $\omega_{pl} \sim m_a$. In the intergalactic medium ($\omega_{pl} \sim 10^{-14}$ eV << $m_a \sim 0.3$ eV), R -> 0 exponentially. GRB photons accumulate no phase drift over cosmological distances. The Fermi LAT constraint ($\Delta v/c <= 10^{-15}$) is evaded without fine-tuning. See Appendix C for the second-order residual calculation.

## 3. Three Phenomenological Principles

### Principle I: Metric Anchoring

Visible-sector fermions couple to an effective metric $g_{\mu\nu}^{\sim} = A^2(\phi) \, g_{\mu\nu}$, with $A(\phi) = 1 + \beta\phi/M_{Pl}$. This anchors Standard Model matter to local curvature while allowing the dilaton $\phi$ to respond to background metric tension [2].

### Principle II: Density-Gradient Potential Shifts

The effective potential $V_{eff}(\phi) = V(\phi) + \rho \, A(\phi)$ shifts with local density. The ultra-shallow index n <= 0.08 ensures the field minimum shifts minimally across the density range of interest, satisfying solar-system gravity tests.

### Principle III: Direction-Dependent Refractive Modulation

The benchmark amplitude $\alpha_K$ manifests as:

$$\Delta n(k, t) = \alpha_K \, (v_{Earth} \cdot k) \cos(\omega_{sid} t)$$

*Averages to zero over isotropic sampling — evades static Lorentz-violation bounds · Predicts specific phase-locked directional signal*

## 4. Feasibility Scan and Survival Ridge

An illustrative 200,000-point Monte Carlo scan over ($g_0$, n, $m_a$) was performed with hard cuts from:

| Constraint | Limit | Density probed | Reference |
|---|---|---|---|
| CAST 2024 | $g < 5.7\times10^{-11}$ GeV$^{-1}$ | $\rho_{solar} \sim 10^2$ g/cm$^3$ | arXiv:2406.16840 [3] |
| HB Stars | $g < 6.6\times10^{-11}$ GeV$^{-1}$ | $\rho_{HB} \sim 10^4$ g/cm$^3$ | Ayala et al. JCAP 2020 [4] |
| CMB distortions | $g < 10^{-12}$ GeV$^{-1}$ | $\rho_{cosmic} \sim 10^{-29}$ g/cm$^3$ | Marsh (2016) [5] |
| Lab detectability | $g > 10^{-13}$ GeV$^{-1}$ | $\rho_{lab} \sim 10^{-12}$ g/cm$^3$ | $F\times Q \sim 10^8$ |

Approximately 1.5% of sampled points survive all four constraints (random seed 20260414). The viable manifold is forced to n <= 0.08 by the CMB bound: a steep screening exponent would allow the coupling to diverge in cosmic voids. An illustrative benchmark on the survival ridge:

$$g_0 \sim 5e\text{-}13\ \text{GeV}^{-1} \cdot n \sim 0.04 \cdot m_a \sim 0.3\ \text{eV}$$

*$g_{lab} = g_{eff}(\rho_{lab}) = 1.51e\text{-}12\ \text{GeV}^{-1}$ · all four constraints satisfied*

*Monte Carlo scope: This is an illustrative feasibility scan, not a global likelihood fit or a Bayesian posterior. ~1.5% survival rate demonstrates that the parameter space is narrow but non-empty. A full profile likelihood analysis would sharpen the boundaries.*

## 5. Material Candidate: Indium Tin Oxide (ITO)

The resonance condition $\omega_{pl} \sim m_a = 0.3$ eV requires an electron density $n_e \sim 6.5\times10^{19}$ cm$^{-3}$. Indium tin oxide (ITO) achieves this in its epsilon-near-zero (ENZ) regime: a tunable plasma frequency in the near-infrared, high carrier density, and strong field confinement. ITO is a well-characterized material widely studied in ENZ photonics. Alternative candidates: heavily doped silicon (Si:P) and Weyl semimetals.




*Figure 1: Omega Series Vol. II, Paper I. Top (wide) — survival ridge in the ($m_a$, $g_{lab}$) plane, colored by screening index n. Red/orange: CAST and HB excluded. Colored band: survival ridge (n < 0.08). Yellow crosshair: benchmark. Top right — density-coupling profile at n = 0.04: all three density probes pass their respective bounds. Bottom (wide) — plasma-resonant response R vs $\omega_{pl}$ for three cavity finesse values. Sharp peak at $m_a$ = 0.3 eV. ITO ENZ window matches resonance. IGM response: negligible (GRB safe). Bottom right — three pre-registered detection criteria.*

## 6. Pre-Registered Detection Criteria

| Criterion | Prediction | Null test |
|---|---|---|
| 1. Density Discontinuity | Signal decreases when apparatus pressurized with inert gas (follows g ~ rho^-n law) | Signal increases with pressure: model falsified |
| 2. Sidereal Orientation | $1.7\times10^{-4}$ modulation at $T_{sid}$ = 23.934 h | 180-deg B-field rotation flips modulation sign but preserves amplitude |
| 3. Refractive Offset | Wavelength-independent residual rotation scaling as $B^2L^2$ | Wavelength dependence: Thomson scattering, not axion-photon |

**OMEGA SERIES VOL. II, PAPER I — THE CHAMELEON RULES: SUMMARY**

Mechanism: density-screened coupling g_eff = g_0 x (rho/rho_0)^(-n)

Saturation floor: rho_min = 1e-15 g/cm^3 (prevents CMB violation in voids)

Survival ridge: n < 0.08 (ultra-shallow, unlike standard chameleon n~0.5-2)

Illustrative benchmark: g_0=5e-13 GeV^-1, n=0.04, m_a=0.3eV

Resonance locality: R ~ 0 in IGM (omega_pl~1e-14 eV << m_a) -> GRB safe

ITO ENZ: omega_pl ~ 0.3 eV -> ne ~ 6.5e19 cm^-3 (achievable)

Unitarity cap: G_max = F x Q < 1e10 -> Delta_n ~ 1e-4 from g_0~1e-12

Three pre-registered detection criteria (density, sidereal, refractive)

No detection claimed. Residuals = upper bounds.

Random seed: 20260414 | doi:10.5281/zenodo.1082194

## References

[1] Khoury J. & Weltman A. Phys.Rev.D 69, 044026 (2004). Chameleon fields.

[2] Brax P. et al. Phys.Rev.D 76, 124034 (2007). Chameleon scalar field.

[3] CAST Collaboration. arXiv:2406.16840 (2024). Axion-photon coupling limit.

[4] Ayala A. et al. JCAP 11, 012 (2020). Horizontal branch stellar cooling bound.

[5] Marsh D.J.E. Phys.Rept. 643, 1 (2016). Axion cosmology review.

[6] Burrage C. & Sakstein J. Living Rev.Rel. 21, 1 (2018). Chameleon tests.

[7] Hook A. & Huang J. JHEP 06, 036 (2018). Resonance locality.

[8] Fermi LAT & GBM Collaborations. Nature 462, 331 (2009). GRB timing constraint.

[9] Raffelt G. & Stodolsky L. Phys.Rev.D 37, 1237 (1988). Axion-photon conversion.

[10] Kadunic D. MSDS Omega Series Vol.I. doi:10.5281/zenodo.1082194.

### Appendix A: Transfer Function and Gain Ceiling

The axion-photon conversion probability in a cavity of finesse F uses effective length $L_{eff} = F \times L$ [9]. Maximum physically allowed gain: $G_{max} = F \times Q <= 10^{10}$ for $F \sim 10^5$ and $Q \sim 10^3$-$10^5$. Observable refractive modulation: $\Delta n_{eff} \sim g_{a\gamma} \times G_{max}^{1/2}$, which yields $\sim 10^{-4}$ for $g_0 \sim 10^{-12}$ $GeV^{-1}$ without violating unitarity.

### Appendix B: Monte Carlo Scan Details

Parameter ranges: $g_0 \in [10^{-14}, 10^{-8}]$ $GeV^{-1}$; $n \in [0, 1]$; $m_a \in [10^{-2}, 10]$ eV. Densities ($g/cm^3$): $\rho_{solar}=10^2$, $\rho_{HB}=10^4$, $\rho_{lab}=10^{-12}$, $\rho_{min}=10^{-15}$. 200,000 points sampled. Random seed: 20260414. ~1.5% survive all four constraints.

### Appendix C: Second-Order GRB Delay (Order-of-Magnitude)

The second-order residual phase variance is $\langle\delta\phi^2\rangle \sim (\omega L/c)^2 \alpha_K^2/6$. Over a GRB baseline L ~ 1 Gpc, this yields a raw fractional timing drift $\varepsilon^{(2)} \sim 4.8\times10^{-9}$. Further suppressed by the decoherence factor $(\lambda/L_{coh})^2$, giving $\varepsilon^{(2)}_{obs} <= 10^{-16}$ — well below the Fermi LAT bound. (Order-of-magnitude estimate; full calculation deferred to Paper III.)