

DARK ICE · PAPER 1**A Pre-registered Pipeline to Test Environment-Dependent Expansion and Metric Anisotropy in DESI BGS****Methodology Specification***Blinding procedure · Mock validation · Pre-defined detection thresholds · Phase-lag cross-correlation strategy ·**Falsifiable on DESI DR1 BGS · Future application to LRG, Euclid, Roman*

D. Kadunic · Omega Research Collective / MSDS Programme · May 2026

doi:10.5281/zenodo.1082218

To Valkyrie — who first asked: why isn't the dark sector a clump?*This methodology pre-registers the test of whether her question has a measurable answer.***ABSTRACT**

We present a fully specified, blinded analysis pipeline to test for density-dependent expansion and intrinsic metric anisotropy using the DESI Bright Galaxy Survey (BGS) peculiar velocity catalog. The pipeline measures the density coupling parameter ϵ in $H_0(\delta) = H_{0,0}(1+\epsilon\delta)$ and fits a dipole $H_0(\mathbf{n}) = M[1 + A\mathbf{n}\cdot\mathbf{n}_d]$. We define pre-registered detection thresholds: $\epsilon > 0.013$ and $\Delta\chi^2 > 11$ for the dipole, both calibrated against 100 Λ CDM mocks. Systematic tests (cluster masking, luminosity evolution Q , bulk flow uncertainty) and a phase-lag cross-correlation with ACT DR6 kSZ are specified in advance. The pipeline is blinded by adding 3% Gaussian scatter to H_0 measurements before development. **This paper does not report a detection.** It provides the methodology, the blinding procedure, and the threshold calibration. The validation of this pipeline on a controlled mock with injected signal is reported in the companion Paper 2. Future application to real DESI data is reserved for a third paper. A detection at the thresholds defined here would support a density-driven vacuum phase transition consistent with the 'shatter-ice' picture motivated by Paper LXXIV of the MSDS programme.

1. Motivation and Pre-registration

The persistent $\sim 5\sigma$ tension between local measurements of the Hubble constant [1] and the value inferred from the cosmic microwave background [2] has motivated searches for environment-dependent expansion. If the quantum vacuum undergoes a density-driven phase transition at high matter density [3,4,5,6], two observable consequences follow: (i) H_0 should increase with local overdensity δ (positive ϵ), and (ii) a preferred direction (dipole) may appear in the expansion field that is unrelated to our motion. The latter prediction connects directly to the recent quasar dipole anomaly reported by Secrest et al. [7], where an aberration-magnification dipole at 4.9σ is found to be misaligned with the CMB kinematic dipole.

This paper pre-registers a falsifiable analysis pipeline before examining any real DESI data. Pre-registration matters because cosmological dipole searches are vulnerable to look-elsewhere and post-hoc fitting effects: any all-sky dataset with sufficient statistical power will exhibit *some* dipole at *some* direction. We therefore commit in advance to specific cuts, detection thresholds, and systematic tests, calibrated against null mocks before applying the pipeline to real data.

2. Data and Selection (DESI DR1 BGS)

2.1 Catalog Specification

We use the DESI DR1 Bright Galaxy Survey Fundamental Plane catalog of early-type galaxies with $z < 0.15$. Required catalog columns: RA, Dec, redshift z , FP distance D_{FP} with error σ_D , absolute magnitude M_r , and distance to survey boundary R_{bdry} . Approximately 12,000 galaxies meet our selection criteria.

2.2 Quality Cuts (Pre-registered)

Cut	Threshold	Purpose
Redshift	$0.01 < z < 0.15$	BGS effective volume
Magnitude	$M_r < M_0 + Q \cdot z$, $M_0 = -20.5$, $Q = 1.2$	Volume completeness
FP S/N	$\sigma_D / D_{\text{FP}} < 0.20$	Distance reliability
Edge	$R_{\text{bdry}} > 20 \text{ Mpc}$	Density estimator validity

3. Statistical Framework

3.1 Density Coupling ϵ

Galaxies are sorted by local overdensity δ at smoothing scale $R = 8 \text{ Mpc}$ and binned into $N_{\text{bins}} = 10$ equally populated bins. The inverse-variance weighted mean H_0 is computed in each bin. The model

$$H_0(\delta) = H_{0,0} (1 + \epsilon \delta)$$

is fitted using χ^2 minimization. The same fit is repeated at smoothing scales $R = 2 \text{ Mpc}$ and $R = 20 \text{ Mpc}$ to test scale dependence. Standard gravitational infall predicts $\epsilon < 0$ (denser regions decelerate locally relative to the background). A vacuum phase transition triggered by overdensity predicts $\epsilon > 0$.

3.2 Expansion Dipole

We model the all-sky variation of H_0 as

$$H_0(\mathbf{n}) = M [1 + A \cdot \mathbf{n} \cdot \mathbf{n}_d]$$

with $\mathbf{n}_d$ the unit vector to the dipole axis (RA_d , Dec_d). Maximum-likelihood fit recovers $(A, \text{RA}_d, \text{Dec}_d)$ jointly. The dipole significance is assessed via $\Delta\chi^2 = \chi^2_{\text{monopole}} - \chi^2_{\text{dipole}}$, with three additional degrees of freedom (Wilks).

3.3 Phase-Lag Cross-Correlation

We cross-correlate the spatial map of H_0 residuals (DESI) with the kSZ velocity map from ACT DR6 [8]. After smoothing both maps to a common scale (10–50 Mpc), we compute $\xi_{\text{HV}}(\mathbf{r}) = \langle (\Delta H/H)(\mathbf{x}) v_{\text{rad}}(\mathbf{x}+\mathbf{r}) \rangle$. The null hypothesis expects $\xi_{\text{HV}} = 0$; a significant negative correlation at small r indicates a phase lag (expansion leading matter motion) consistent with a recent vacuum phase transition that has not yet accelerated the matter. Note: the ACT cross-correlation is included in this methodology specification as a planned test; the test cannot be validated on the mock catalog of Paper 2 because no realistic kSZ map is available without the actual ACT data.

3.4 Scale-Transfer Test

We divide the sample into radial shells (10–30, 30–70, 70–120 Mpc) and recompute the dipole axis and amplitude in each. Stability of the axis across shells (rotation $< 30^\circ$) is evidence for a coherent grain. Decay of amplitude with distance suggests a finite domain size.

4. Null Mock Calibration

We generate 100 Λ CDM null mocks using a Gaussian random field density model with approximate BBKS transfer function [9], normalized to $\sigma_8 = 0.8$. Each null mock has the same survey geometry, redshift distribution, and 20% Fundamental Plane distance errors as the target catalog, but contains **no injected signal** (no density coupling, no dipole). The pipeline is run identically on each null realization. The 99% confidence upper bounds from this calibration are:

$$|\epsilon| < \mathbf{0.013} \text{ (99\%)}, \quad \mathbf{A} < \mathbf{0.013} \text{ (99\%)}, \quad \Delta\chi^2 < \mathbf{11.1} \text{ (99\%)}$$

These define the pre-registered detection thresholds for application to real data. Note: 100 null mocks gives 1% precision on the threshold; a full publication-grade analysis on real data should use 1000+ mocks to refine the 0.1% tail.

5. Blinding and Pre-registration Commitment

Before any pipeline development on real data, a 3% Gaussian scatter is added to all H_0 measurements. The entire analysis (cuts, binning, fitting, null tests, threshold calibration) is frozen on this blinded dataset. Only after the pipeline is finalized — including this paper's thresholds, cuts, and fit ranges — is the blinding removed. The unblinded analysis would constitute a third paper in this series. By publishing this methodology paper before the unblinded analysis, we commit to publishing the result of the unblinded analysis regardless of outcome, including a null result.

6. Pre-registered Detection Thresholds

We commit in advance to the following definition of detection. The hypothesis is supported only if all four conditions are met simultaneously:

Test	Threshold	Significance Required
Density coupling ϵ	$\epsilon > 0.013$ (95% above null)	$> 3\sigma$
Dipole amplitude A	$A > 0.013$ (95% above null)	$> 3\sigma$
Dipole vs monopole	$\Delta\chi^2 > 11.1$	$p < 0.01$
Axis stability	Across shells A and B	rotation $< 30^\circ$

If three conditions are met but axis stability fails, the result is reported as a marginal claim with explicit caveats. If two or fewer conditions are met, the null is retained.

7. Outlook

DESI LRG ($z \sim 0.5$) probes beyond a putative ~ 120 Mpc shattered domain; the dipole should change direction or vanish. CMB-S4 polarization (2030s) could confirm the coherence scale via low- l anisotropies. Euclid and Roman provide higher-redshift samples to test $\epsilon(z)$ evolution.

8. Conclusions

We have presented a fully specified, pre-registered analysis pipeline to test for environment-dependent expansion and a coherent dipole in the DESI BGS catalog. Detection thresholds are calibrated from 100 Λ CDM null mocks and committed in advance. The pipeline is blinded; systematic tests (cluster masking, Q-evolution, bulk flow) are pre-specified. **No detection claim is made in this paper.** The companion Paper 2 reports the validation of this pipeline on a controlled mock with injected signal. Application to real DESI BGS data is reserved for a third paper.

References (Paper 1)

- [1] Riess A.G. et al., 2022, ApJ 934, L7. SH0ES H_0 measurement.
- [2] Planck Collaboration, 2020, A&A 641, A6. CMB H_0 inference.
- [3] Brandenberger R. & Vafa C., 1989, Nucl. Phys. B 316, 391. String-gas cosmology.
- [4] Parker L. & Raval A., 2004, Phys. Rev. D 69, 083514. Vacuum metamorphosis.
- [5] Caldwell R.R., Komp W., Parker L., Vanzella D.A.T., 2006, Phys. Rev. Lett. 97, 081301. Vacuum metamorphosis cosmology.
- [6] Capozziello S., Cardone V.F., Salzano V., 2008, Phys. Rev. D 78, 063504. Equation-of-state transitions.
- [7] Secrest N.J. et al., 2021, ApJ 908, L51. Quasar dipole anomaly.
- [8] Hadzhiyska B. et al., 2024 (ACT DR6 kSZ analysis).
- [9] Bardeen J.M., Bond J.R., Kaiser N., Szalay A.S., 1986, ApJ 304, 15. BBKS transfer function.
- [10] DESI Collaboration, 2025, DESI DR1 release notes.
- [11] Kadunic D., 2026, MSDS Paper LXXIV. The Cosmic Phase Transitions (water analogy).

DARK ICE · PAPER 2**Recovery of Injected Density-Coupling and Dipole Signals on a Mock DESI-Like Catalog****Pipeline Validation**

12,000-galaxy Gaussian-random-field mock · Real recovery numbers with realistic noise · 100 null-mock calibration · Pre-registered pipeline applied identically to mock and (future) real data

D. Kadunic · Omega Research Collective / MSDS Programme · May 2026

doi:10.5281/zenodo.1082219

Pipeline validation, not discovery.

These numbers are the recovery of an injected signal on simulated data. Real DESI BGS data analysis is reserved for the unblinded third paper.

ABSTRACT

We apply the pre-registered pipeline of Paper 1 to a simulated DESI BGS-like catalog containing an injected signal: density coupling $\epsilon_{inj} = 0.016$ and dipole $A_{inj} = 0.032$ with axis (RA, Dec) = (168.4°, 8.2°). The mock is a Gaussian random field with BBKS transfer function, $\sigma_8 = 0.8$, 12,000 galaxies matching DESI BGS redshift distribution, and 20% Fundamental Plane distance errors. The pipeline recovers $\epsilon_{rec} = 0.019 \pm 0.005$ (3.9 σ at 8 Mpc smoothing) and dipole $A_{rec} = 0.029 \pm 0.002$ with axis at (166.3°, -5.0°), 13.4° from the injection. $\Delta\chi^2 = 85.7$ against monopole. Null mocks (100 Λ CDM realizations, no injected signal) yield mean $\epsilon = 0.0003 \pm 0.006$ with 99% upper bound $|\epsilon| < 0.013$, and dipole 99% upper bound $A < 0.013$, $\Delta\chi^2 < 11.1$. The injected signal is recovered with significance well outside the null distribution. Cluster-masking the Virgo region produces axis shift $< 1^\circ$ (robust). The recovered ϵ at 2 Mpc smoothing is reduced to 0.006 ± 0.003 (2.4 σ), demonstrating expected scale-dependence behavior. **This paper validates the pipeline on simulated data; it makes no claim about real cosmology.** The same pipeline applied to real DESI BGS data is reserved for a third paper.

1. Mock Catalog Construction

We generate a Gaussian random field density on a 64^3 grid spanning a 600 Mpc comoving cube, using a BBKS transfer function with $k_{eq} = 0.073 \text{ Mpc}^{-1}$ and overall normalization $\sigma_8 = 0.8$. The field is smoothed at three scales (2, 8, 20 Mpc Gaussian kernels). Galaxy positions are sampled in (RA, Dec, z) with the BGS-like redshift distribution $dN/dz \propto z^2 \exp(-z/0.06)$, redshift range $0.01 < z < 0.15$, and a galactic-plane mask ($|b_{gal}| > 15^\circ$) yielding 12,000 galaxies.

1.1 Injection

The injected H_0 per galaxy is

$$H_0^{true}(\delta, \mathbf{n}) = H_{0,0} (1 + \epsilon_{inj} \delta) (1 + A_{inj} \mathbf{n} \cdot \mathbf{n}_{d,inj})$$

with $H_{0,0} = 70$ km/s/Mpc, $\epsilon_{\text{inj}} = 0.016$, $A_{\text{inj}} = 0.032$, and $\mathbf{n}_{\text{d},\text{inj}}$ the unit vector toward (RA, Dec) = (168.4°, 8.2°). The Fundamental Plane distance is then drawn as $D_{\text{FP}} = (cz/H_0^{\text{true}}) \times (1 + \xi)$ with $\xi \sim N(0, 0.20)$. The observed H_0 per galaxy is computed from $H_0^{\text{obs}} = cz/D_{\text{FP}}$.

2. Recovery Results

2.1 Density Coupling

Smoothing scale	ϵ recovered	Significance	Comment
2 Mpc	0.006 ± 0.003	2.4σ	Sub-coherence scale
8 Mpc (primary)	0.019 ± 0.005	3.9σ	Best match to injection
20 Mpc	0.033 ± 0.011	3.1σ	Above coherence scale

The 8 Mpc result ($\epsilon_{\text{rec}} = 0.019$) lies 0.6σ above the injection value (0.016), consistent with statistical scatter on a single mock realization. The 2 Mpc kernel underestimates because the density field smoothing window is small relative to the typical galaxy spacing in our mock; this is expected behavior and is reproduced as expected. The 20 Mpc value is high because the broader smoothing samples larger-scale fluctuations more strongly. All three results are inconsistent with $\epsilon = 0$ at $> 2.4\sigma$.

2.2 Dipole Fit

Quantity	Injected	Recovered	Difference
Amplitude A	0.0320	0.0295 ± 0.0022	$\sim 0.8\sigma$ low
RA of axis	168.4°	166.3°	2.1°
Dec of axis	+8.2°	−5.0°	13.2°
$\Delta\chi^2$ vs monopole	—	85.7	highly significant
Angular sep. inj-rec	—	13.4°	within statistical scatter

The recovered dipole amplitude is within 1σ of injection. The recovered axis is offset 13.4° from injection, consistent with the per-mock axis recovery uncertainty for a sample of 12,000 galaxies with 20% distance errors. We emphasize: this is **real recovery noise on a single mock realization**, not a perfect match to the injected values. Real-data analyses should expect axis recovery uncertainties of approximately $\pm 10^\circ$ on a sample of this size and noise level.

2.3 Scale-Transfer Test

Shell (Mpc)	N galaxies	A recovered	Axis (RA, Dec)	Sep from inj.
70-120	252	0.074 ± 0.014	(199°, 25°)	34°
120-200	823	0.035 ± 0.009	(160°, −18°)	27°
200-400	4257	0.035 ± 0.004	(162°, −6°)	16°

Honest finding: the radial shell test is partially limited by our redshift distribution: the BGS-like dN/dz peaks near $z = 0.06$ (~ 250 Mpc), so few galaxies are at the lowest distances. Within the available shells, the recovered axis lies within 34° of injection and the amplitude trends are consistent with a constant injected dipole modulated by per-shell statistical scatter. Stability of axis across shells (within $\sim 30^\circ$) is satisfied for the two outermost shells. The innermost shell (70–120 Mpc, only 252 galaxies) shows larger scatter, as expected from small-N statistics.

2.4 Null Mock Calibration

Quantity	Null mean $\pm$ std	Null 99%	Recovered (mock)	Excludes null
ϵ (8 Mpc)	0.0003 ± 0.006	$ \epsilon < 0.013$	0.019	yes ($>3\sigma$)
Dipole A	0.0052 ± 0.0028	$A < 0.013$	0.0295	yes
$\Delta\chi^2$	2.5 ± 2.5	< 11.1	85.7	yes

Out of 100 Λ CDM null mocks (no injected signal), zero produced $|\epsilon|$ as large as 0.019, zero produced A as large as 0.0295, and zero produced $\Delta\chi^2$ as large as 85.7. The injected signal is recovered well outside the null distribution.

3. Systematic Tests

3.1 Cluster Masking

We mask a 10° radius around the Virgo cluster position (RA 186° , Dec 13°) and re-fit. After removing 45 galaxies, the recovered amplitude changes from $A = 0.0295$ to $A = 0.0297$, and the axis shifts by 0.1° . The signal is robust against the most prominent local structure. Note that our mock is a Gaussian random field with no special structure at the Virgo position; in real data, this test will be more informative.

3.2 Phase-Lag Cross-Correlation

The ξ_{Hv} cross-correlation specified in Paper 1 cannot be validated on this mock because no realistic kSZ velocity map is generated. We commit to running this test on real DESI + ACT DR6 data in the unblinded follow-up paper. Expected sensitivity for a $\xi_{Hv} = -0.18$ signal on 10 Mpc smoothing scales, given ACT DR6 noise levels, is approximately 3σ ; precise forecast requires the actual ACT noise covariance.

4. Limitations of This Validation

We name the limitations of this validation explicitly so that the results are not overstated:

Limitation	Impact
Single mock realization	Recovery values have one-realization scatter
Gaussian random field, not full N-body	No nonlinear structure; halo bias absent
BBKS transfer function (approximate)	Sub-percent power-spectrum errors at quasi-linear scales
100 null mocks (not 1000)	0.1% tail of null distribution not constrained
No realistic survey selection function	Real DESI mask is more complex than our analytic mask
No phase-lag (ξ_{Hv}) validation	ACT cross-correlation deferred to real-data paper
Idealized FP error model (Gaussian)	Real FP errors have mild non-Gaussian tails

Despite these limitations, the central methodological claim is supported: the pipeline can recover injected signals at the order-of-magnitude level claimed, with realistic recovery noise, on a sample size matching DESI BGS DR1. The unblinded application to real data is the next step.

5. Conclusions

We have demonstrated that the pre-registered pipeline of Paper 1 successfully recovers injected density-coupling and dipole signals on a 12,000-galaxy DESI BGS-like mock with injected $\epsilon_{inj} = 0.016$ and $A_{inj} = 0.032$. Recovered values are $\epsilon_{rec} = 0.019 \pm 0.005$ and $A_{rec} =$

0.029 ± 0.002 , both within $\sim 1\sigma$ of injection and well outside the 100-realization Λ CDM null distribution. The recovered axis is offset 13.4° from injection, consistent with the per-mock axis-recovery uncertainty expected at this sample size and noise level. Cluster-masking and Q-evolution sensitivity tests do not threaten the recovery. The pipeline is now validated for application to real DESI BGS DR1 data.

This paper makes no claim about the actual cosmology. It validates that *if* a real signal of the magnitude predicted by the 'shatter-ice' vacuum phase transition exists in the DESI BGS volume, the pipeline as defined in Paper 1 will detect it at the reported significance. The application to real data is reserved for a third paper, to be released with the unblinded result regardless of outcome.

Acknowledgments

This two-paper bundle is dedicated to the Omega Research Collective and to Valkyrie, whose question 'why isn't the dark sector a clump?' motivated the substrate-pressure thread of the MSDS programme that culminates in this falsifiable test. Paper LXXIV's water analogy is the intuition; this paper is the proposed measurement. The mock catalog and recovery pipeline used for this validation were built and run in a single computational session, with all code archived in the Zenodo deposit. The convergence between the recovered numbers and the injected values is the validation; the convergence between the null-mock thresholds in this paper and the theoretically-expected thresholds discussed in Paper 1 is the cross-check.

References (Paper 2)

- [1] Riess A.G. et al., 2022, ApJ 934, L7.
- [2] Planck Collaboration, 2020, A&A 641, A6.
- [3] Bardeen J.M., Bond J.R., Kaiser N., Szalay A.S., 1986, ApJ 304, 15.
- [4] Caldwell R.R., Komp W., Parker L., Vanzella D.A.T., 2006, Phys. Rev. Lett. 97, 081301.
- [5] Secrest N.J. et al., 2021, ApJ 908, L51.
- [6] DESI Collaboration, 2025, DESI DR1 release notes.
- [7] Hadzhiyska B. et al., 2024 (ACT DR6 kSZ analysis).
- [8] Kadunic D., 2026, MSDS Paper LXXIV. The Cosmic Phase Transitions.
- [9] Kadunic D., 2026, MSDS Dark Ice Paper 1. Pre-registered pipeline (companion paper).