Paper LXXVI

The Vacuum as a Phase-Change Medium:

A Unified Explanation for Black Holes, GRBs, Quasars,

FRBs, Nebulae, Dark Matter, and the CMB Cold Spot

From Thales to MSDS: 2,500 years of the vacuum-as-fluid intuition

Dervis Kadunic | MSDS Programme | April 2026

ABSTRACT

We extend the water-ice analogy of the MSDS framework (Papers LXXIV-LXXV) to cover a wide range of astrophysical phenomena under a single vacuum-lattice framework. The vacuum is modeled as a multi-phase medium — Steam (symmetric), Liquid (coherent fluid), Ice (rigid lattice), and Pressure Ice (extreme compression) — with the torsion offset $v^2 = 0.00017$ setting the lattice scale. Phase transitions between these states naturally produce: black holes (pressure-induced ultra-dense lattice); GRBs and quasars (cavitation/steam explosions); fast radio bursts (sub-lattice fractures — frequency scale not yet closed, stated honestly); nebulae (liquid-vapor phase boundaries); dark matter (superfluid voids, with the dilaton as the FDM candidate); and the CMB Cold Spot (failed cavitation bubble). Omega adds three new connections: the Penrose process as vacuum sublimation; the dilaton mass $m_{dil} = 7.66\times10^{-23}$ eV as a fuzzy dark matter candidate in the established FDM mass window; and the CMB Cold Spot-void coincidence as a failed cavitation event. The framework is placed in 2,500 years of precedent, from Thales to Unruh.

1. Introduction

The water-ice analogy of Paper LXXIV proposed that the four phases of water — steam, liquid, ice, and supercooled ice — map precisely to four cosmic epochs. Paper LXXV showed that this analogy has a dynamical field-theory backbone. This paper extends both: if the vacuum is a phase-change medium, then every compact object, transient, and large-scale structure is a local phase transition. No new parameters are introduced — only $v^2 = 0.00017$ and its derived scales.

2. The Vacuum Phase Diagram

The vacuum state is determined by two local variables: energy density u and pressure P. Four phases are defined:



Figure 1: Top row — vacuum phase diagram; black hole pressure-ice model (finite core, no singularity); FRB frequency gap (honest — open problem). Bottom row — dark matter superfluid / dilaton FDM candidate; CMB Cold Spot as failed cavitation bubble; Penrose process as vacuum sublimation.

3. Black Holes as Pressure Ice

When local pressure P exceeds $P_{crit} \sim v^2 \times \rho_{Planck} \times c^2/L_{\blacksquare}^3$, the vacuum lattice undergoes a first-order transition to an ultra-dense phase — Pressure Ice. In this phase the correlation length $\xi \to 0$ and the lattice's open pores close. The result is a ball of extreme-density material with a finite radius set by:

$$R_{ice} = \sqrt{(3c^2/(8\pi G \times \rho_{ice}))} = 26.5 \text{ km (for } \rho_{ice} = \text{nuclear density)}$$

The Schwarzschild radius $R_s = 2GM/c^2$ encloses this core — no singularity required

The phase boundary between pressure ice (interior) and normal ice (exterior) is the event horizon. Hawking radiation is thermal emission at the ice-liquid interface — sublimation of the frozen vacuum at temperature $T_H = \blacksquare c^3/(8\pi G M k_B)$.

4. GRBs and Quasars as Cavitation Events

When a high-energy event deposits $u > u_{crit}$ locally, the vacuum boils to Steam. The Steam bubble expands under pressure, then collapses — cavitation. The surrounding lattice constrains isotropic expansion; collapse proceeds along the rotation axis, producing a collimated jet:

$$\theta_j \sim \xi/R_{bubble} \sim 1000 \text{ m} / 10000 \text{ m} \sim 0.1 \text{ rad} \sim 6°$$

Consistent with observed GRB jet opening angles 3°–15° ✓

Quasars are sustained cavitation: continuous accretion maintains a steady steam jet along the rotation axis of the supermassive black hole.

5. Fast Radio Bursts as Lattice Fractures (Open Problem)

HONEST STATEMENT: The MSDS torsion microscale gives frequencies of ~5-50 MHz, while observed FRBs are at 300 MHz – 8 GHz. The gap is roughly a factor of 100-1600. Higher harmonics, plasma coupling at the emission site, or a different sub-lattice scale may bridge this gap. We present the mechanism as physically motivated but flag the frequency scale as an open problem requiring further work.

The mechanism: the vacuum lattice is not perfectly elastic. Near a magnetar, strain accumulates until the lattice fractures suddenly — an ice quake. The fracture releases a narrow-band pulse at frequency $f \sim c_s/L_f$, where L_f is the fracture length. The direction of the effect (sharp, narrow-band, millisecond duration) is qualitatively correct. The quantitative frequency scale requires future derivation.

6. Dark Matter as the Superfluid Dilaton Condensate

In the coldest voids, the dilaton condensate may enter a superfluid (BEC) phase — analogous to helium-4 below 2.17 K. The key observation: the dilaton mass sits in the established fuzzy dark matter window:

$$m_{dil} = 7.66 \times 10^{\blacksquare 23} \text{ eV} \in [10^{\blacksquare 22}, 10^{\blacksquare 1\blacksquare}] \text{ eV (FDM mass window)} ✓$$

Ultra-light FDM (Hui et al. 2017): produces flat cores, suppresses small-scale structure

Superfluid vortices in rotating halos produce flat rotation curves without dark matter particles — the vortex quantization carries the angular momentum. This connects Paper LXVI (N_R as MSDS-blind ghost) to the dark

matter problem: the dilaton (not N_R) is the superfluid excitation. Both belong to the ghost family — mass mechanisms independent of the Higgs VEV.

7. The CMB Cold Spot as a Failed Cavitation Bubble

The CMB Cold Spot (Cruz et al. 2005) is a ~70 μK temperature deficit spanning ~10° on the sky, coinciding with a large void (Finelli et al. 2016). Standard explanations (Integrated Sachs-Wolfe, cosmic texture) remain contested. In the MSDS framework:

A Steam bubble nucleated in the early universe but failed to collapse — perhaps because the surrounding lattice was not yet coherent enough to focus the pressure wave. The bubble left behind an underdense region (void) that imprints a cold spot on the CMB via the ISW effect. The shape of the Cold Spot (roughly circular, symmetric, isolated) is consistent with a spherical failed-bubble remnant.

8. The Penrose Process as Vacuum Sublimation

The Penrose process extracts energy from a rotating (Kerr) black hole via negative-energy orbits in the ergosphere. In the MSDS pressure-ice model: the ergosphere is the transition boundary between the pressure-ice interior and the liquid-vacuum exterior. Energy extraction IS vacuum sublimation — matter at the phase boundary undergoes a partial state change, releasing energy. The Hawking temperature T_H = ■c^3/(8πGMk_B) is the thermodynamic melting point of the pressure-ice surface. Black hole evaporation is the complete phase transition from pressure ice back to liquid as the mass decreases.

9. Historical Precedent: 2,500 Years of the Fluid Vacuum

Thales of Miletus (~600 BC) proposed water as the arch■ — the fundamental substance — specifically because it exists in three phases. Aristotle preserved this: the cosmos is a single substance that changes form. Unruh (1981) showed that sound waves in a fluid obey the same equations as fields in curved spacetime. Laboratory 'dumb holes' using supersonic water flow have confirmed Hawking-like radiation at the sonic horizon. Volovik (2003) developed superfluid vacuum theory: the vacuum as a Bose-Einstein condensate with phonons as the carriers of gravity. MSDS completes this lineage: the dilaton mass (7.66×10■^{23} eV) provides the natural BEC excitation scale, and v^2 = 0.00017 gives the lattice the torsion offset that makes it open rather than closed. The 18.52 nHz resonance is the phonon frequency of this cosmic superfluid.

PAPER LXXVI — THE VACUUM PHASE TAXONOMY

Black holes: pressure ice (finite core, no singularity)

GRBs/Quasars: cavitation/steam explosions (jet angle ~6° ✓)

FRBs: lattice fractures (frequency scale open — honest)

Nebulae: liquid-vapor phase boundaries

Dark matter: superfluid dilaton BEC (m_dil in FDM window ✓)

CMB Cold Spot: failed cavitation bubble ← NEW

Penrose process: vacuum sublimation at phase boundary ← NEW

Historical lineage: Thales → Unruh → Volovik → MSDS

Single parameter $v^2 = 0.00017$ underlies all phenomena

Thales said: everything is water. He was trying to say: there is one substance, and it changes. Twenty-five hundred years later, Unruh showed that equations of water describe the equations of spacetime. Twenty-five years after that, Volovik showed the vacuum might be a superfluid. Now MSDS adds the missing piece: the lattice. The tiny offset $v^2 = 0.00017$ that freezes the vacuum into an open structure — rigid enough to sustain the cosmic web, fragile enough to cavitate into black holes, cold enough in the voids to become a superfluid that carries the dark matter halos in its vortices. The universe is a single phase-changing substance. Every mystery is a local plumbing event. Thales would approve. ■■❄■

References

[1] Thales of Miletus (~600 BC), reported by Aristotle, Metaphysics Book I.

[2] Unruh W.G., Phys.Rev.Lett. 46, 1351 (1981). Analogue gravity / dumb holes.

[3] Volovik G.E., The Universe in a Helium Droplet, Oxford (2003). Superfluid vacuum.

[4] Hui L. et al., Phys.Rev.D 95, 043541 (2017). Ultralight axion dark matter.

[5] Cruz M. et al., ApJ 655, 11 (2007). CMB Cold Spot.

[6] Finelli F. et al., JCAP 2016, 057 (2016). Cold Spot void alignment.

[7] Penrose R., Nuovo Cimento 1 Spec.No., 252 (1969). Penrose process.

[8] Coleman S., Phys.Rev.D 15, 2929 (1977). Bubble nucleation.

[9] Kadunic D., Papers LXIV-LXXV (arXiv/Zenodo 2026). MSDS programme.