

# Long-term intravenous epoetin- $\alpha$ / darbepoetin- $\alpha$ ratio in iron-replete hemodialysis patients

Andrea Icardi, Paolo Sacco, Francesca Salvatore, Umberto Romano

Nephrology and Dialysis Unit, Department of Internal Medicine, La Colletta Hospital, Arenzano, Genoa - Italy

**ABSTRACT: Background:** Equivalence of intravenous (i.v.) and subcutaneous (s.c.) dosage requirements is a notable characteristic of darbepoetin- $\alpha$  (DPO), as opposed to other epoetins (EPOs). Currently in Europe, the EPOs/DPO conversion factor (200 IU EPOs = 1  $\mu$ g DPO) does not take into account the route of drugs administration. To better define this ratio we have conducted a prospective, long-term trial in a group of hemodialysis patients.

**Subjects and methods:** At the start, we evaluated 40 iron-replete hemodialysis patients, but the final study was performed in the remaining 25 patients. During the first 6 months, patients were on i.v. epoetin- $\alpha$  (EPO $\alpha$ ) maintenance therapy (phase 1: T-6 to T0). After conversion to i.v. DPO (initial 200:1 ratio) the observation was prolonged for a period of 12 months (phase 2: T0 to T12). DPO was administered at extended dose intervals and the EPO $\alpha$ /DPO rate was adjusted every month to maintain hemoglobin (Hgb) stability. Iron status and factors inhibiting erythropoiesis were continually checked to exclude unstable patients.

**Results:** Phase 1: EPO $\alpha$  weekly mean dose showed no significant variation. Phase 2: EPO $\alpha$ /DPO conversion factor progressively rose from 200 to  $256.7 \pm 86.9$  IU/ $\mu$ g at T7 ( $p < 0.005$ ) and  $336.8 \pm 104.3$  IU/ $\mu$ g at T12 ( $p < 0.0005$ ). DPO weekly mean dosage decreased from  $40.0 \pm 12.0$   $\mu$ g/week at T0 to  $31.6 \pm 3.7$   $\mu$ g/week at T7 ( $p < 0.005$ ) and  $24.6 \pm 7.0$   $\mu$ g/week at T12 ( $p < 0.0005$ ). Mean weekly/patient acquisition cost of EPO $\alpha$  was  $\text{€}70.6 \pm 21.3$  (T-6 to T0); after switching, the cost of DPO was  $\text{€}72.4 \pm 22.7$  (T0) and fell to  $\text{€}53.1 \pm 11.2$  during T6 to T12.

**Conclusions:** The progressive increase of EPO $\alpha$ /DPO ratio demonstrated that i.v. DPO requires lower doses compared with i.v. EPO $\alpha$ . When drugs are administered i.v., the starting EPO $\alpha$ /DPO conversion factor should be increased over the 200:1 ratio, similar to recommendations outlined in the United States and Japan. DPO dose reduction translated to notable cost-savings.

**Key words:** Epoetin- $\alpha$ , Darbepoetin- $\alpha$ , Intravenous, Conversion factor, Hemoglobin, Stable iron-replete hemodialysis patients

## INTRODUCTION

Erythropoiesis-stimulating agents (ESAs) have revolutionized the management of renal anemia. Since about 1990, 2 forms of recombinant human erythropoietin, epoetin-alpha (EPO $\alpha$ ) and epoetin-beta (EPO $\beta$ ), have been available, and more recently darbepoetin-alfa (DPO), a novel erythropoiesis stimulating protein, has been introduced.

EPO $\alpha$  and EPO $\beta$  differ slightly in their glycosylation patterns and especially in their formulations (1, 2). These differences have effects not only on the pharmacokinetic and pharmacodynamic properties of EPOs, but also on their stability, solubility and im-

munogenicity (3-5). Because of this latter feature, in December 2002, health authorities in all European countries formally contraindicated subcutaneous (s.c.) administration of EPO $\alpha$  in renal patients, recommending exclusively the intravenous (i.v.) route. The route of administration of EPOs is important for immunogenic response and in relation to therapeutic efficacy, and as a result, current U.S. and European guidelines advocate more effectively using the s.c. rather than the i.v. route, although the immunogenicity of EPOs is particularly prominent when administered via the s.c. route (6, 7).

The amino acid sequence of DPO differs from that of the EPOs and is engineered to obtain increased num-

bers of sialic acid-containing carbohydrate chains (8). In vivo DPO potency appears superior, and the pharmacokinetic profile shows a much longer elimination half-life and slower clearance than EPO $\alpha$ , with prolonged erythropoietic effect (9). Therefore DPO can be administered less frequently to obtain the same biological response of EPOs (10). Another DPO peculiarity is the equivalence of i.v. and s.c. dosing requirements to achieve target levels of hemoglobin (Hgb) (11).

The conversion factor between EPOs and DPO was based on the formula equating the protein mass of the 2 molecules (200 IU EPOs = 1  $\mu$ g DPO) (12). This ratio is recommended in the European DPO approved dosing schedule (Aranesp and Nespo Product Information), but several trials have shown that this ratio does not accurately predict the dosage needs for the conversion of all patients and conditions (13-16). The 200:1 equivalence ratio is applied to both EPO $\alpha$  and EPO $\beta$ . Moreover, the route of ESAs administration has not been taken into account. For these reasons, recent papers have outlined that this "approved" EPO/DPO dose ratio should be challenged in clinical practice (17-19).

To better define this conversion factor, we have conducted a long-term, prospective study on a group of stable, iron-replete hemodialysis patients. The trial began with i.v. EPO $\alpha$  treatment in maintenance phase and was continued for 6 months. At the end of this first period, using the initial 200:1 equivalence, all patients were switched to less frequent administration of DPO maintaining the same route. During the following 12 months, DPO dose was adjusted to maintain stable Hgb levels as recommended by European guidelines (7). The EPO $\alpha$ /DPO ratio was evaluated to establish eventual changes from the accepted value. Consequent pharmacoeconomic analysis was performed. Hematological markers of iron status and potential factors inhibiting erythropoiesis and response to ESAs were continually checked to exclude unstable patients.

### SUBJECTS AND METHODS

The study was approved by the therapeutics commission of our institution. Informed consent was obtained from 40 hemodialysis patients on stable treatment with i.v. EPO $\alpha$  and with normal iron status. During the trial (before and after switching to DPO) 15 patients were excluded on the basis of malignancy (n=1), infection (n=2), evidence of blood loss (n=2), variability of Hgb level (n=4), kidney transplantation (n=2), transfer to other dialysis unit (n=1) or death (n=3). Demographics and etiology of end-stage renal disease of the remaining 25 subjects are shown in Table I. At the

beginning, 4 patients showed relative erythropoietin resistance, as gauged by EPO $\alpha$  dosage requirement of >200 IU/kg per week (19); all of those completed the study. The management of uremic anemia included maintenance Hgb >11 g/dL, determined from a sample taken prior to the dialysis session (at the end of the longest interval in the weekly dialysis schedule). When Hgb level fell below 10.5 g/dL or exceeded 12.5 g/dL, a 25% stepwise adjustment was performed in the total weekly ESA dose (up or down) and/or dosing frequency, according to the type of ESA.

During the first period, patients were observed with i.v. EPO $\alpha$  in maintenance therapy for 6 months (phase 1: T-6 to T0). The second period began with conversion to i.v. DPO and lasted for 12 months (phase 2: T0 to T12).

DPO was administered at extended dosing intervals as follows: patients receiving EPO $\alpha$  into the venous line 2 (n=10) or 3 (n=13) times a week were switched to once-a-week administration; patients receiving EPO $\alpha$  once weekly (n=2) were switched to once every other week. Following the recommendations published in the current approved European Summary of Product Characteristics, the initial EPO $\alpha$ /DPO conversion factor was calculated using the 200 IU to 1  $\mu$ g ratio. If needed, to maintain Hgb stability, this ratio was adjusted every month, during the following period of observation.

Iron status was assessed monthly by measuring ferritin levels and transferrin saturation (TSAT). An attempt to ensure adequate iron stores was made by aiming for a ferritin level of up 100 ng/mL and a TSAT up to 20%. To achieve these targets, a standardized i.v. iron gluconate dose (62.5 mg) was administered once a week. When ferritin levels fell to <200 ng/mL, i.v. iron

**TABLE I - PATIENTS INCLUDED: DEMOGRAPHICS AND ETIOLOGY OF END-STAGE RENAL DISEASE**

Demographics and etiology	
Age (years)	63.8 $\pm$ 15.8
Race: Caucasian (%)	100
Sex	
Male	21
Female	4
Weight (kg)	62.4 $\pm$ 10.6
Dialysis duration (months)	31.6 $\pm$ 28.9
Dialysis modality (%)	
Hemodialysis	56
Hemodiafiltration	44
Etiology (n)	
Glomerulonephritis	4
Polycystic kidney disease	3
Obstructive uropathy	3
Hypertension	3
Other/unknown	12




Fig. 1 - Phase 1: no variation of mean weekly EPO $\alpha$  doses.




Fig. 2 - Progressive increase of EPO $\alpha$ /DPO conversion factor from switchover to 12th month (phase 2); \* $p < 0.005$ , \*\* $p < 0.0005$ , vs. baseline.

gluconate was increased to 3 times a week (187.5 mg/week).

To exclude other causes of deficiency anemias associated with hemodialysis, water-soluble, dialyzable vitamins such as folate and cobalamin were measured in the serum every 3 months. Folic acid and vitamin B<sub>12</sub> supplementation was used: levofolinic acid (1.5 mg i.v. at the end of dialysis session) was administered in the case of low serum folate levels or macrocytosis; idroxicobalamin (1,000  $\mu$ g i.v. at the end of dialysis session) was given for serum cobalamin deficiency or megalocytosis. Indices of nutritional status, inflammation and parathyroid activity were monitored by quarterly measure of serum albumin, C-reactive protein (CRP) and intact parathyroid hormone (i-PTH). A 3-monthly estimation of dialysis adequacy was performed by means of the urea Kt/V (20).

Acquisition costs of dosing ESAs were expressed in euros (€).

### Statistical analysis

Collected data are presented as means  $\pm$  SD. Statistical analyses were carried out using Student's paired *t*-test to explore the differences in means of the study variables measured during the two periods of ESAs treatment regimens. Statistical significance was set at a *p* value of  $< 0.05$ .

## RESULTS

Phase 1: EPO $\alpha$  weekly mean doses did not show any significant variations (Fig. 1). Phase 2: the conversion factor, when switching from i.v. EPO $\alpha$  to i.v. DPO at T0, rose progressively from 200 to  $256.7 \pm 86.9$  IU/ $\mu$ g at T7





Fig. 3 - Decrease of DPO weekly mean dosage from switchover to 12th month (phase 2); \* $p < 0.005$ , \*\* $p < 0.0005$ .

( $p < 0.005$ ) and  $336.8 \pm 104.3$  IU/ $\mu$ g at T12 ( $p < 0.0005$ ) (Fig. 2). This increase started to be statistically significant at T8 ( $p < 0.005$ ), T9 ( $p < 0.05$ ), T10 ( $p < 0.005$ ), T11 and T12 ( $p < 0.0005$ ) with respect to the EPO $\alpha$ /DPO ratio at T7. No difference was found between patients with high EPO $\alpha$  dosage requirements ( $> 200$  IU/kg/week) and those with lower dosage requirements, in their relative EPO $\alpha$ /DPO conversion factor, and no variation of dose requirement was noted when the dosage interval of DPO was fixed once every other week in comparison with weekly administration (data not shown).

DPO weekly mean dosage demonstrated a decrease from  $40.0 \pm 12.0$   $\mu$ g/week at T0 to  $31.6 \pm 3.7$   $\mu$ g/week at T7 ( $p < 0.005$ ),  $29.4 \pm 1.6$   $\mu$ g/week at T8 ( $p < 0.0005$ ) and  $24.6 \pm 7.0$   $\mu$ g/week at T12 ( $p < 0.0005$ ) (Fig. 3).



**Fig. 4 - No variation of hemoglobin, ferritin and TSAT mean values during long-term observation period (phase 1 and phase 2). TSAT = transferrin saturation.**

***Hemoglobin, iron stores, vitamins, indices of nutritional status, inflammation, parathyroid activity and dialysis adequacy***

Mean Hgb concentration was  $11.38 \pm 0.46$  g/dL at T-6,  $11.4 \pm 0.47$  g/dL at T0,  $11.31 \pm 0.42$  g/dL at T6 and  $11.38 \pm 0.54$  at T12. Monthly Hgb, serum ferritin and TSAT mean values are shown in Table II and in Figure 4. No significant variation of Hgb, serum ferritin and TSAT mean levels during the long-term observation period (phase 1 and phase 2) was observed. Measures of serum folate and cobalamin, serum albumin, PCR, i-PTH and urea Kt/V, performed every 3 months, did not show any significant change during the present study (Tab. III).


**TABLE II - HEMOGLOBIN AND MARKERS OF IRON STATUS**

	T-6	T-5	T-4	T-3	T-2	T-1	T0
Hgb (g/dL)	11.38 ± 0.46	11.32 ± 0.41	11.37 ± 0.59	11.41 ± 0.62	11.4 ± 0.63	11.7 ± 0.70	11.4 ± 0.47
Ferritin (ng/mL)	305.8 ± 123.3	288 ± 98.5	278.1 ± 85.3	295.1 ± 136.2	277 ± 108.8	284.2 ± 119.9	321.5 ± 128.1
TSAT (%)	25.5 ± 6.6	24.7 ± 5.2	24.8 ± 5.2	23.6 ± 4.8	23.6 ± 4.2	24.4 ± 5.2	26.0 ± 7.8

	T1	T2	T3	T4	T5	T6
Hgb (g/dL)	11.8 ± 0.97	11.97 ± 0.73	11.52 ± 0.52	11.34 ± 0.54	11.32 ± 0.35	11.31 ± 0.42
Ferritin (ng/mL)	309.5 ± 105.5	313.1 ± 114.2	371.2 ± 145.3	338.5 ± 102.9	351.3 ± 108.8	344.4 ± 111.3
TSAT (%)	26.6 ± 7.6	26.9 ± 7.4	26.2 ± 7.7	26.5 ± 7.6	26.7 ± 7.5	24.4 ± 6.4

	T7	T8	T9	T10	T11	T12	
Hgb (g/dL)	11.5 ± 0.34	11.45 ± 0.61	11.6 ± 0.5	11.42 ± 0.60	11.41 ± 0.42	11.38 ± 0.54	NS
Ferritin (ng/mL)	336.8 ± 90.1	321.6 ± 106.5	361.8 ± 140.5	281.3 ± 92.1	285.6 ± 84.9	308.8 ± 103.8	NS
TSAT (%)	24.7 ± 6.3	24.6 ± 6.3	27.8 ± 8.5	24.2 ± 4.5	24.2 ± 4.3	24.3 ± 4.9	NS

Hgb = hemoglobin; TSAT = transferrin saturation.



**Fig. 5 - Mean weekly patient cost of ESAs (open column, EPO $\alpha$ ; shaded columns, DPO).**

### Cost of ESAs

Our analysis was based on a local institutional contract price of €1.74/200 IU of EPO $\alpha$  and €1.81/ $\mu$ g of DPO. Mean weekly patient cost of EPO $\alpha$  during phase 1 (T-6 to T0) was €70.6  $\pm$  21.3. At the switching to DPO, the cost increased to €72.4  $\pm$  22.7, due to the different prices of ESAs. During the first 6 months of phase 2 (T0 to T6), mean weekly patient cost of DPO was €69.0  $\pm$  16.8 and in the second 6-month period (T6 to T12), it fell to €53.1  $\pm$  11.2, in consequence of the marked reduction of dosage requirements (Fig. 5).

### DISCUSSION

Our study confirms the objective difficulties in maintaining hemodialysis patients for long-term observation when the stability of clinical and laboratory parameters is needed to validate the behavior of the primary end point. This fact is particularly evident during the studies regarding uremic anemia, influenced as they are by variability of hematological data, in consequence of different factors. In patients with end-stage renal disease, Hgb levels fluctuate much more compared with the general population. This individual variability caused a significant portion of dialysis patients (29%) to move both above and below the target Hgb level range during a relatively short period of follow-up (21). We used a long period of averaging, which has been suggested as a method to provide more stable measurement of anemia correction outcomes; however, we lost 30% of patients to follow-up, because of biological and comorbid conditions influencing their response to ESAs therapy and individual Hgb level variability caused by other factors (i.e., sampling methods, laboratory assays and dialytic variations of body weight) (6, 22, 23).

For these reasons, our study population includes a relatively small number of patients; nevertheless, it assures a valid evaluation of the ratio between EPO $\alpha$  and DPO during a long period of monitoring free of limits linked to the fluctuation of the data compared before and after ESA change.

The EPO $\alpha$ /DPO conversion factor progressively increased to a statistically significant change at the beginning of the second 6-month period. In the following

**TABLE III - VITAMINS AND INDICES OF NUTRITIONAL STATUS, INFLAMMATION, PARATHYROID ACTIVITY AND DIALYSIS ADEQUACY**

	T-6	T-3	T0	T3	T6	T9	T12	p Value
Folate (n.v. >5.38 ng/mL)	20.5 $\pm$ 7.4	20.2 $\pm$ 7.5	21.7 $\pm$ 6.6	24.1 $\pm$ 3.6	24.1 $\pm$ 3.3	24.7 $\pm$ 6.6	24.4 $\pm$ 3.0	NS
Cobalamin (n.v. 211-911 pg/mL)	727.2 $\pm$ 602.4	794.2 $\pm$ 636.5	775.2 $\pm$ 670.1	850.2 $\pm$ 580.7	859.8 $\pm$ 554.7	856.4 $\pm$ 450.5	827.8 $\pm$ 413.9	NS
Albumin (n.v. 3.4-4.7 g/dL)	4.0 $\pm$ 0.4	3.9 $\pm$ 0.4	3.9 $\pm$ 0.5	3.8 $\pm$ 0.5	4.1 $\pm$ 0.6	4.1 $\pm$ 0.6	4.0 $\pm$ 0.6	NS
CRP (n.v. 0-5.0 mg/L)	5.6 $\pm$ 5.6	5.9 $\pm$ 6.0	7.8 $\pm$ 7.7	7.6 $\pm$ 7.5	6.1 $\pm$ 6.2	6.7 $\pm$ 6.7	5.3 $\pm$ 5.2	NS
i-PTH (pg/mL)	224.8 $\pm$ 157.0	155.5 $\pm$ 131.5	234.2 $\pm$ 182.3	245.3 $\pm$ 212.4	164.7 $\pm$ 109.3	182.0 $\pm$ 141.2	134.7 $\pm$ 90.6	NS
Kt/V (Daugirdas)	1.4 $\pm$ 0.2	1.4 $\pm$ 0.2	1.4 $\pm$ 0.2	1.4 $\pm$ 0.3	1.4 $\pm$ 0.2	1.4 $\pm$ 0.2	1.4 $\pm$ 0.2	NS

CRP = C-reactive protein; i-PTH = intact parathyroid hormone; NS = not significant; n.v. = normal value.



months, the increase was even more evident and was highest at the end of the period of study. In line with this result, average weekly dose of DPO requirements showed a reduction from the time of the switchover, becoming statistically significant at T7 and even more noticeable.

Higher EPOs/DPO ratios than commonly admitted (200:1) were previously reported (13, 14, 19); nevertheless, we found low conversion rates referred to by other authors (15, 16). Most studies examining ESAs with regard to EPOs and DPO have not specified the forms of recombinant human erythropoietin administered or whether via the i.v. or s.c. route.

It is now well known that a reduction in dosage requirement is usually needed when switching from i.v. to s.c. administration of EPOs (24, 25). Pharmacokinetic studies demonstrate that EPO $\alpha$  has lower bioavailability but a longer half-life after s.c. than after i.v. administration (26). A more rapid fall in circulating erythropoietin with i.v. dosing may affect erythropoiesis through shortened survival of red blood cells, resulting in neocytolysis, the process by which young erythrocytes are destroyed (27, 28). Changing of EPO $\alpha$  from s.c. to i.v. route, in compliance with the resolution of the European Medical Agency, results in an increase in mean dose for maintenance of Hgb levels in hemodialysis patients (29, 30). In contrast, some AA find no differences between the dose of s.c. and i.v. EPOs needed (31, 32). However, studies showing that administration route does not interact with the efficacy of EPOs have had a smaller number of patients and have been observational, mostly in an unselected population.

Longer DPO terminal half-life means that the serum concentration of the drug remains above the threshold value longer, and therefore the efficacy is equal for i.v. or s.c.. DPO has a 3-fold longer half-life than EPOs following i.v. administration and approximately 2-fold longer than EPOs by the s.c. route (9).

The above-mentioned reports support our results. Lower doses of i.v. DPO were able to maintain target Hgb levels than those of i.v. EPO $\alpha$ . Consequently, when ESAs are administered i.v., the initial EPO $\alpha$ /DPO conversion factor should be increased over the conventional ratio based on the formula equating the protein mass of the 2 molecules. Indeed, Japanese Approved Product Information suggests that patients be switched to i.v. DPO based on the ratio of 240 IU EPOs = 1  $\mu$ g DPO.

In our dialysis population, the EPO $\alpha$ /DPO conversion rate was not different between patients with relative erythropoietin resistance and those with lower dosage. When the dosage interval of DPO was fixed once every other week no variation of dose requirement was noted in comparison with patients treated with weekly administration. However, these observations are limited

by the restricted number of patients enrolled in the study. We can conclude that i.v. DPO administered once weekly was more efficacious than twice- or thrice-weekly i.v. EPO $\alpha$ , in keeping with the findings of other authors and confirming our previous reports (33, 34). The increase of EPO $\alpha$ /DPO conversion factor was progressive, but became noticeable and significant some months after switching. This late result may reflect the frequency in changing ESAs dosage; in our dialysis unit it may take up to 4 weeks for adjustment of the dosage given to patients in the maintenance phase. Therefore a long-term observation period was appropriate to obtain the data. The fact that only "selected" patients were included in the analysis could allow a better response to DPO treatment; however, we emphasize that the dose of EPO $\alpha$  required was stable in the maintenance period, while in the same patients DPO dose decreased progressively after the switchover.

Medication acquisition costs are influenced by institutional contract pricing and dosing. The progressive decrease of DPO weekly mean dose translated to appreciable cost-savings compared with those for EPO $\alpha$ . This result was much more significant, if we took into account the higher price of DPO than that of EPO $\alpha$ , on the basis of the contract pricing of our institution.

In conclusion, the starting EPOs/DPO conversion factor set by the European schedule is probably not appropriate for the i.v. route. It is desirable that the fixed ratio be replaced by an aggregate of conversion factors, similar to recommendations outlined in the United States (35). In these prescribing notes, dose conversion between EPOs and DPO is not always linear across the dosing spectrum and is mostly higher than 200 IU EPOs = 1  $\mu$ g DPO. However, further clinical trials will still be required to investigate the effects of route of ESAs administration and others factors that influence erythropoietic response, to provide conclusive evidence regarding the switching of treatments.

Address for correspondence:

Andrea Icardi, M.D.  
U.O. Nefrologia e Dialisi  
Ospedale "La Colletta"  
Via del Giappone, 5  
16011 Arenzano, Genova, Italy  
andrea.icardi.usl3@libero.it

## REFERENCES

1. Storrington PL, Tiplady RJ, Gaines Das RE, Stenning BE, Lamikanra A, Rafferty B, Lee J. Epoetin alfa and beta differ in their erythropoietin isoform compositions and biological properties. *Br J Haematol* 1998; 100: 79-89.
2. Hermeling S, Schellekens H, Crommelin DJ, Jiskoot W. Micelle-associated protein in epoetin formulations: a risk factor for immunogenicity? *Pharm Res* 2003; 20: 1903-7.
3. Halstenson CE, Macres M, Katz SA, Schnieders JR, Watanabe M, Sobota JT, Abraham PA. Comparative pharmacokinetics and pharmacodynamics of epoetin alfa and epoetin beta. *Clin Pharmacol Ther* 1991; 50: 702-12.
4. Haselbeck A. Epoetins: differences and their relevance to immunogenicity. *Curr Med Res Opin* 2003; 19: 430-2.
5. Boven K, Stryker S, Knight J, Thomas A, Van Regenmortel M, Kemeny DM, Power D, Rossert J, Casadevall N. The increased incidence of pure red cell aplasia with an Eprex formulation in uncoated rubber stopper syringes. *Kidney Int* 2005; 67: 2346-53.
6. National Kidney Foundation. NKF-K/DOQI clinical practice guidelines for anemia of chronic kidney disease: update 2000. *Am J Kidney Dis* 2001; 37 (Suppl 1): S182-238.
7. Locatelli F, Aljama P, Bárány P, Canaud B, Carrera F, Eckardt KU, Hörl WH, Macdougall IC, Macleod A, Witek A, Cameron S. Revised European best practice guidelines for the management of anaemia in patients with chronic renal failure. *Nephrol Dial Transplant* 2004; 19 (Suppl 2): ii1-47.
8. Egrie JC, Browne JK. Development and characterization of novel erythropoiesis stimulating protein (NESP). *Nephrol Dial Transplant* 2001; 16 (Suppl 3): 3-13.
9. Macdougall IC, Gray SJ, Elston O, Breen C, Jenkins B, Browne J, Egrie J. Pharmacokinetics of novel erythropoiesis stimulating protein compared with epoetin alfa in dialysis patients. *J Am Soc Nephrol* 1999; 10: 2392-5.
10. Locatelli F, Olivares J, Walker R, Wilkie M, Jenkins B, Dewey C, Gray SJ; European/Australian NESP 980202 Study Group. Novel erythropoiesis stimulating protein for treatment of anemia in chronic renal insufficiency. *Kidney Int* 2001; 60: 741-7.
11. Macdougall IC. Optimizing the use of erythropoietic agents: pharmacokinetic and pharmacodynamic considerations. *Nephrol Dial Transplant* 2002; 17 (Suppl 5): 66-70.
12. Aljama P, Bommer J, Canaud B, Carrera F, Eckardt KU, Hörl WH, Krediet RT, Locatelli F, Macdougall IC, Wikström B; NESP Usage Guidelines Group. Practical guidelines for the use of NESP in treating renal anaemia. *Nephrol Dial Transplant* 2001; 16 (Suppl 3): 22-8.
13. Nissenson AR. Dosing darbepoetin alfa. *Am J Kidney Dis* 2002; 40: 872.
14. Barnett AI, Crémieux PY. Dose conversion from epoetin alfa to darbepoetin alfa for patients with chronic kidney disease receiving hemodialysis. *Pharmacotherapy* 2003; 23: 690-4.
15. Deray G. Once-weekly erythropoietic therapy: is there a difference between the available preparations? *Nephrol Dial Transplant* 2003; 18: 2455-6.
16. Jacobs C, Frei D, Perkins AC. Results of the European Survey on Anaemia Management 2003 (ESAM 2003): current status of anaemia management in dialysis patients, factors affecting epoetin dosage and changes in anaemia management over the last 5 years. *Nephrol Dial Transplant* 2005; 20 (Suppl 3): iii3-24.
17. Scott SD. Dose conversion from recombinant human erythropoietin to darbepoetin alfa: recommendations from clinical studies. *Pharmacotherapy* 2002; 22: S160-5.
18. Locatelli F, Canaud B, Giacardy F, Martin-Malo A, Baker N, Wilson J. Treatment of anaemia in dialysis patients with unit dosing of darbepoetin alfa at a reduced dose frequency relative to recombinant human erythropoietin (rHuEPO). *Nephrol Dial Transplant* 2003; 18: 362-9.
19. Roger SD, Cooper B. What is the practical conversion dose when changing from epoetin alfa to darbepoetin outside of clinical trials? *Nephrology* 2004; 9: 223-8.
20. Daugirdas JT. Simplified equations for monitoring Kt/V, PCRn, Ekt/V and ePCRn. *Adv Ren Replace Ther* 1995; 2: 295-304.
21. Lacson E Jr, Ofsthun N, Lazarus JM. Effect of variability in anemia management on hemoglobin outcomes in ESRD. *Am J Kidney Dis* 2003; 41: 111-24.
22. Movilli E, Pertica N, Camerini C, Canarini GC, Brunori G, Scolari F, Maiorca R. Predialysis versus postdialysis hematocrit evaluation during erythropoietin therapy. *Am J Kidney Dis* 2002; 39: 850-3.
23. Stirling CM, Patel RK, Simpson K. Haemoglobin target for haemodialysis patients - is it achievable? *Nephrol Dial Transplant* 2005; 20 (Suppl 5): v268.
24. Kaufman JS, Reda DJ, Fye CL, Goldfarb DS, Henderson WG, Kleinman JG, Vaamonde CA. Subcutaneous compared with intravenous epoetin in patients receiving hemodialysis. *N Engl J Med* 1998; 339: 578-83.
25. Besarab A, Reyes CM, Hornberger J. Meta-analysis of subcutaneous versus intravenous epoetin in maintenance treatment of anemia in hemodialysis patients. *Am J Kidney Dis* 2002; 40: 439-46.
26. Macdougall IC, Roberts DE, Coles GA, Williams JD. Clinical pharmacokinetics of epoetin (recombinant human erythropoietin). *Clin Pharmacokinet* 1991; 20: 99-113.
27. Kaufman J. Choosing the right approach with recombinant human erythropoietin. *Clinical Opinions in Renal Anaemia*. Prime Medica Limited 2005; 2: 10-2.
28. Alfrey CP, Rice L, Udden MM, Driscoll TB. Neocytolysis: physiological down-regulator of red-cell mass. *Lancet* 1997; 349: 1389-90.
29. Linde T, Furuland H, Wikström B. Effects of switch from subcutaneous to intravenous epoetin alfa administration in hemodialysis patients. *J Am Soc Nephrol* 2003; 14: 844A.
30. Fullerton L, Bhattacharjee D, Roe SD, Cassidy MJ. Erythropoietin: subcutaneous or intravenous dosing? *British Journal of Renal Medicine*; Winter 2002/2003: 17-9.
31. Jacobs C, Hörl WH, Macdougall IC, Valderrabano F, Parrondo I, Abraham IL, Segner A. European best practice guidelines 9-13 anaemia management. *Nephrol Dial Transplant* 2000; 15 (Suppl 4): 33-42.
32. Pizzarelli F, David S, Sala P, Icardi A, Casani A. Iron-replete hemodialysis patients do not require higher EPO dosages when converting from subcutaneous to intravenous administration: results of the Italian Study on Erythropoietin Converting (ISEC). *Am J Kidney Dis* 2006; 47: 1027-35.
33. Hori W. IV darbepoetin alfa is more effective than i.v. rHuEPO in maintaining target Hb concentrations in patients with renal anaemia [abstract]. *Nephrol Dial Transplant* 2002; 17 (Suppl 1): 24.
34. Icardi A, Sacco P, Salvatore F, Romano U. Conversion factor of epoetin-alfa / darbepoetin-alfa administered intravenously in hemodialysis stable patients. *Nephrology* 2005; 10 (Suppl): A319.
35. Amgen Inc. Aranesp (darbepoetin alfa): prescribing information. Thousand Oaks, CA: Amgen Inc 2002.

Received: October 24, 2005

Revised: March 28, 2006

Accepted: November 23, 2006

© Società Italiana di Nefrologia