Minimum Pizzas

Each pizza consists of 4 slices. There are $\it N$ friends and each friend needs exactly $\it X$ slices.

Find the **minimum** number of pizzas they should order to satisfy their appetite.

Input Format

- ullet The first line of input will contain a single integer \it{T} , denoting the number of test cases.
- ullet Each test case consists of two integers N and X, the number of friends and the number of slices each friend wants respectively.

Output Format

For each test case, output the **minimum** number of pizzas required.

Constraints

- $1 \le T \le 100$
- $1 \leq N, X \leq 10$

Sample 1:	
Input	Coutput
4 15 26 43 35	2 3 3 4
Explanation:	
Test case 1: There is only 1 friend who requires 5 slices. If he orders 1 pizza, he will get only 4 slices. Thus, at least 2 pizzas should be ordered to have required number of slices.	
$\textbf{Test case } 2 \textbf{:} \ \textbf{There are } 2 \textbf{ friends who require } 6 \textbf{ slices each. Thus, total } 12 \textbf{ slices are required. To get } 12 \textbf{ slices, they should order } 3 \textbf{ pizzas.}$	
$\textbf{Test case 3:} \ \text{There are 4 friends who require 3 slices each. Thus, total } \ 12 \ \text{slices are required.} \ \textbf{To get } \ 12 \ \text{slices, they should order 3 pizzas.}$	
Test case 4: There are 3 friends who require 5 slices each. Thus, total 15 slices are required. To get 15 slices, they should order at least 4 pizzas.	