

P-adic Numbers

Kevin Guillen

UC Santa Cruz

March 10th 2022

Number Systems

A review of what we know:

► $\mathbb{N} = \{1, 2, \dots\}$

Number Systems

A review of what we know:

- ▶ $\mathbb{N} = \{1, 2, \dots\}$
- ▶ $\mathbb{Z} = \{-2, -1, 0, 1, 2\}$ through —

Number Systems

A review of what we know:

- ▶ $\mathbb{N} = \{1, 2, \dots\}$
- ▶ $\mathbb{Z} = \{-2, -1, 0, 1, 2\}$ through $-$
- ▶ $\mathbb{Q} = \{\dots, -\frac{4}{7}, \frac{1}{2}, \dots\}$ through $/$

Number Systems

What now? We need new machinery. Bring in limits of sequences (x_n) where $x_n \in \mathbb{Q}$ for all n .

► $\forall \varepsilon > 0$

Number Systems

What now? We need new machinery. Bring in limits of sequences (x_n) where $x_n \in \mathbb{Q}$ for all n .

► $\forall \varepsilon > 0$

► $\exists N$

Number Systems

What now? We need new machinery. Bring in limits of sequences (x_n) where $x_n \in \mathbb{Q}$ for all n .

► $\forall \varepsilon > 0$

► $\exists N$

► $\forall n > N$

Number Systems

What now? We need new machinery. Bring in limits of sequences (x_n) where $x_n \in \mathbb{Q}$ for all n .

- ▶ $\forall \varepsilon > 0$
- ▶ $\exists N$
- ▶ $\forall n > N$
- ▶ $|x_n - x| < \varepsilon$

Number Systems

This leads to the idea of Cauchy Sequences.

Number Systems

This leads to the idea of Cauchy Sequences.

Which is when terms in a sequence become arbitrarily close to one another and in a sense settle on something. [4]

Number Systems

This leads to the idea of Cauchy Sequences.

Which is when terms in a sequence become arbitrarily close to one another and in a sense settle on something. [4]

When we consider a sequence like:

$$(x_n) = \left(1 + \frac{1}{n}\right)^n$$

Number Systems

This leads to the idea of Cauchy Sequences.

Which is when terms in a sequence become arbitrarily close to one another and in a sense settle on something. [4]

When we consider a sequence like:

$$(x_n) = \left(1 + \frac{1}{n}\right)^n$$

Settles on something that is not in \mathbb{Q}

Number Systems

This leads to the idea of Cauchy Sequences.

Which is when terms in a sequence become arbitrarily close to one another and in a sense settle on something. [4]

When we consider a sequence like:

$$(x_n) = \left(1 + \frac{1}{n}\right)^n$$

Settles on something that is not in \mathbb{Q} , it goes to e .

Which then gives us \mathbb{R} .

Distance (Absolute Value)

Limit was dependent on $|\cdot|$, which is our measure of distance.

Distance (Absolute Value)

Limit was dependent on $|\cdot|$, which is our measure of distance.

An **absolute value** on a field \mathbb{F} is a function $|\cdot|$ from \mathbb{F} to $\mathbb{R}_{\geq 0}$ that satisfies the following properties for all $a, b \in \mathbb{F}$:

- (1) Positive-definiteness: $|a| = 0 \iff a = 0$
- (2) Multiplicativity: $|ab| = |a| |b|$
- (3) Triangle Inequality: $|a + b| \leq |a| + |b|$

Metric/Cauchy Space

A **Metric Space** is a non-empty set equipped with a metric (distance function).[5]

Metric/Cauchy Space

A **Metric Space** is a non-empty set equipped with a metric (distance function). [5]

Previous examples our metric space was $(\mathbb{Q}, |\cdot|)$.

Metric/Cauchy Space

A **Metric Space** is a non-empty set equipped with a metric (distance function). [5]

Previous examples our metric space was $(\mathbb{Q}, |\cdot|)$.

A metric space is **Complete** if every Cauchy Sequence's limit point is in metric space, also called a **Cauchy Space**. [5]

Metric/Cauchy Space

A **Metric Space** is a non-empty set equipped with a metric (distance function). [5]

Previous examples our metric space was $(\mathbb{Q}, |\cdot|)$.

A metric space is **Complete** if every Cauchy Sequence's limit point is in metric space, also called a **Cauchy Space**. [5]

\mathbb{R} completes \mathbb{Q} .

p -adic valuation

Defined by $v_p : \mathbb{Q} \rightarrow \mathbb{Z} \cup \{\infty\}$

p -adic valuation

Defined by $v_p : \mathbb{Q} \rightarrow \mathbb{Z} \cup \{\infty\}$

$x \in \mathbb{Z}$ then $v_p(x)$ is the unique positive integer satisfying,

$$x = p^{v_p(x)} x' \text{ where } p \nmid x'$$

p -adic valuation

Defined by $v_p : \mathbb{Q} \rightarrow \mathbb{Z} \cup \{\infty\}$

$x \in \mathbb{Z}$ then $v_p(x)$ is the unique positive integer satisfying,

$$x = p^{v_p(x)} x' \text{ where } p \nmid x'$$

EX: $x = 90$ and $p = 3$ the 3-adic valuation of x to be 2 since,

$$90 = 3^2 \cdot 10$$

p -adic valuation

$x \in \mathbb{Q} - \mathbb{Z}$ then $v_p(x)$, we have $x = \frac{a}{b}$ and $a, b \in \mathbb{Z}$. So we define,

$$v_p(x) = v_p(a) - v_p(b)$$

p -adic valuation

$x \in \mathbb{Q} - \mathbb{Z}$ then $v_p(x)$, we have $x = \frac{a}{b}$ and $a, b \in \mathbb{Z}$. So we define,

$$v_p(x) = v_p(a) - v_p(b)$$

EX $x = \frac{13}{24}$ and $p = 2$ we have then that $a = 13$ and $b = 24$

P -adic valuation

$x \in \mathbb{Q} - \mathbb{Z}$ then $v_p(x)$, we have $x = \frac{a}{b}$ and $a, b \in \mathbb{Z}$. So we define,

$$v_p(x) = v_p(a) - v_p(b)$$

EX $x = \frac{13}{24}$ and $p = 2$ we have then that $a = 13$ and $b = 24$

$$v_2(13) = 0 \quad v_2(24) = 3$$

p -adic valuation

$x \in \mathbb{Q} - \mathbb{Z}$ then $v_p(x)$, we have $x = \frac{a}{b}$ and $a, b \in \mathbb{Z}$. So we define,

$$v_p(x) = v_p(a) - v_p(b)$$

EX $x = \frac{13}{24}$ and $p = 2$ we have then that $a = 13$ and $b = 24$

$$v_2(13) = 0 \quad v_2(24) = 3$$

therefore $v_2(x) = 0 - 3 = -3$

p -adic valuation

$x \in \mathbb{Q} - \mathbb{Z}$ then $v_p(x)$, we have $x = \frac{a}{b}$ and $a, b \in \mathbb{Z}$. So we define,

$$v_p(x) = v_p(a) - v_p(b)$$

EX $x = \frac{13}{24}$ and $p = 2$ we have then that $a = 13$ and $b = 24$

$$v_2(13) = 0 \quad v_2(24) = 3$$

therefore $v_2(x) = 0 - 3 = -3$

Finally $v_p(0) = +\infty$

p -adic absolute value

A function $|\cdot|_p : \mathbb{Q} \rightarrow \mathbb{R}_{\geq 0}$ defined as,

$$|x|_p = \begin{cases} \frac{1}{p^{v_p(x)}} & x \neq 0 \\ 0 & x = 0 \end{cases}$$

p -adic absolute value

A function $|\cdot|_p : \mathbb{Q} \rightarrow \mathbb{R}_{\geq 0}$ defined as,

$$|x|_p = \begin{cases} \frac{1}{p^{v_p(x)}} & x \neq 0 \\ 0 & x = 0 \end{cases}$$

inducing the p -adic metric d_p



Field of p -adic numbers, \mathbb{Q}_p .



Field of p -adic numbers, \mathbb{Q}_p .

Completion of \mathbb{Q} with respect to the p -adic metric.



Field of p -adic numbers, \mathbb{Q}_p .

Completion of \mathbb{Q} with respect to the p -adic metric.

Examples (\mathbb{Z} and \mathbb{Z}_p)

What is a p -adic number?

An example of a p -adic number that is not already a rational/integer number is ...

What is a p -adic number?

An example of a p -adic number that is not already a rational/integer number is ... difficult to show.

What is a p -adic number?

An example of a p -adic number that is not already a rational/integer number is ... difficult to show.

p -adic expansion of an integer is simply that number written in base p

What is a p -adic number?

An example of a p -adic number that is not already a rational/integer number is ... difficult to show.

p -adic expansion of an integer is simply that number written in base p

p -adic expansion of a rational number is the formal power series,
 $r \in \mathbb{Q}$,

$$r = \sum_{i=k}^{\infty} a_i p^i$$

Wrangling with ∞

Closeness is measured by the right expansion

Wrangling with ∞

Closeness is measured by the right expansion

2.71828182...

2.71827071...

Wrangling with ∞

Closeness is measured by the right expansion

2.71828182...

2.71827071...

Difference is 0.0001111...

Wrangling with ∞

Closeness is measured by the right expansion

2.71828182...

2.71827071...

Difference is 0.0001111...

Divided by 10^{-4} is 1.111...

Wrangling with ∞

Closeness is measured by the right expansion

2.71828182...

2.71827071...

Difference is 0.0001111...

Divided by 10^{-4} is 1.111...

Note that $10^{-4} = \frac{1}{10^4}$

Wrangling with ∞

Consider working in the 3 – *adic* system we have,

$$\dots 21120$$
$$\dots 10120$$

where this number continues expanding to the left (infinite).

Wrangling with ∞

Consider working in the 3 – *adic* system we have,

$$\dots 21120$$
$$\dots 10120$$

where this number continues expanding to the left (infinite).

The difference will be $\dots 11000$

Wrangling with ∞

Consider working in the 3 – *adic* system we have,

...21120

...10120

where this number continues expanding to the left (infinite).

The difference will be ...11000 ,

$$\dots 1 \cdot 3^4 + 1 \cdot 3^3 + 0 \cdot 3^2 + 0 \cdot 3^1 + 0 \cdot 3^0$$

Wrangling with ∞

Consider working in the 3 – *adic* system we have,

$$\dots 21120$$

$$\dots 10120$$

where this number continues expanding to the left (infinite).






The difference will be $\dots 11000$,

$$\dots \color{red}{1} \cdot 3^4 + \color{red}{1} \cdot 3^3 + \color{red}{0} \cdot 3^2 + \color{red}{0} \cdot 3^1 + \color{red}{0} \cdot 3^0$$

$$(\dots \color{red}{1} \cdot 3^1 + 1)3^3$$

Number Line?

References

-  Fernando Gouvea (2003) *p-adic Numbers: An Introduction*, Springer Science & Business Media, 2003.
-  Alain M. Robert (2000) *A Course in p-adic Analysis*, Springer; 2000th edition
-  Svetlana Katok (2007) *P-adic Analysis Compared With Real (Student Mathematical Library)*, American Mathematical Society
-  Terence Tao (2016) *Analysis I: Third Edition*, Hindustan Book Agency, 1st ed. 2016 edition
-  James Munkres (2000) *Topology*, Pearson College Div; 2nd edition (January 7, 2000)