

Homework 2

Kevin Guillen

MATH 201 — Algebra II — Winter 2022

Note: I used Symbolab to help do the matrix multiplication for P3 so, I apologize if I didn't write all the work required.

Problem P1 Let F be any field, $n \geq 0$ an integer, V and n -dimensional F -vector space. For any integer k such that $0 \leq k \leq n$, let G_k denote the set of k -dimensional F -subspaces W of V . Prove that the action of $GL_F(V)$ on G_k given by

$$g.W = \{g(w) : w \in W\} \in G_k$$

for all $g \in GL_F(V)$, is transitive.

Proof. We defined in class $GL_F(V)$ as,

$$GL_F(V) = \{f : V \rightarrow V \mid f \text{ is } F\text{-linear isomorphism}\}$$

To show that the provided action is transitive we want to show that for any subspace $W, W' \in G_k$ there exists some $f \in GL_F(V)$ that satisfies:

$$f.W = W'$$

Since both W and W' are in G_k we know they are of dimension k . Meaning they their bases can be expressed as $\{a_1, \dots, a_k\}$ and $\{b_1, \dots, b_k\}$ for W and W' respectively. We know that linear transformations map subspaces to subspaces and there exists $f \in GL_F(V)$ such that,

$$f(\alpha_1 a_1 + \dots + \alpha_k a_k) = \alpha_1 b_1 + \dots + \alpha_k b_k$$

we know this is indeed in $GL_F(V)$ because we can see it is a linear transformation through the following, for any $w_1, w_2 \in W$ we have,

$$\begin{aligned} f((\alpha_1 a_1 + \dots + \alpha_k a_k) + (\gamma_1 a_1 + \dots + \gamma_k a_k)) &= f((\alpha_1 + \gamma_1)a_1 + \dots + (\alpha_k + \gamma_k)a_k) \\ &= (\alpha_1 + \gamma_1)b_1 + \dots + (\alpha_k + \gamma_k)b_k \\ &= (\alpha_1 b_1 + \dots + \alpha_k b_k) + (\gamma_1 b_1 + \dots + \gamma_k b_k) \\ &= f(\alpha_1 a_1 + \dots + \alpha_k a_k) + f(\gamma_1 a_1 + \dots + \gamma_k a_k) \end{aligned}$$

and for $c \in F$,

$$\begin{aligned} cf(\alpha_1 a_1 + \dots + \alpha_k a_k) &= c(\alpha_1 a_1 + \dots + \alpha_k a_k) = c\alpha_1 b_1 + \dots + c\alpha_k b_k \\ &= f(c(\alpha_1 a_1 + \dots + \alpha_k a_k)) \end{aligned}$$

meaning we have $f(W) = W' = f.W$. then from our corollary we proved in class (Jan 18), we have that f is an isomorphism and thereby must be in $GL_F(V)$. Showing that the action is transitive. \square

Problem P2 Let $F = \mathbb{F}_{17}$ be the field with 17 elements. For any integer n , we will denote still by n its image in F .

Apply Gaussian elimination to find all the solutions to the linear system

$$2x + 3y + 5z = 10$$

$$4x + 5y + 8z = 11$$

$$2x + 4y + 7z = 2$$

Proof. We begin by writing our system of equations in matrix form,

$$\begin{bmatrix} 2 & 3 & 5 & 10 \\ 4 & 5 & 8 & 11 \\ 2 & 4 & 7 & 2 \end{bmatrix}$$

Next we will label under each matrix the operation we will be performing,

$$\begin{array}{ccccccc} \begin{bmatrix} 2 & 3 & 5 & 10 \\ 4 & 5 & 8 & 11 \\ 2 & 4 & 7 & 2 \end{bmatrix} & \rightarrow & \begin{bmatrix} 2 & 3 & 5 & 10 \\ 0 & 14 & 11 & 7 \\ 2 & 4 & 7 & 2 \end{bmatrix} & \rightarrow & \begin{bmatrix} 2 & 3 & 5 & 10 \\ 0 & 14 & 11 & 7 \\ 0 & 1 & 2 & 9 \end{bmatrix} & \rightarrow & \begin{bmatrix} 2 & 3 & 5 & 10 \\ 0 & 14 & 11 & 7 \\ 0 & 0 & 0 & 0 \end{bmatrix} \\ R2 = R2 - 2R3 & & R3 = R3 - R1 & & R3 = 14R3 - R2 & & R2 = 4R2 \\ \rightarrow \begin{bmatrix} 2 & 3 & 5 & 10 \\ 0 & 5 & 10 & 11 \\ 0 & 0 & 0 & 0 \end{bmatrix} & \rightarrow & \begin{bmatrix} 10 & 0 & 12 & 10 \\ 0 & 5 & 10 & 11 \\ 0 & 0 & 0 & 0 \end{bmatrix} & \rightarrow & \begin{bmatrix} 1 & 0 & 8 & 0 \\ 0 & 5 & 10 & 11 \\ 0 & 0 & 0 & 0 \end{bmatrix} & \rightarrow & \begin{bmatrix} 1 & 0 & 8 & 0 \\ 0 & 1 & 2 & 9 \\ 0 & 0 & 0 & 0 \end{bmatrix} \\ R1 = 5R1 - 3R2 & & R1 = 12R1 & & R2 = 7R2 & & R2 = 7R2 \end{array}$$

This then gives us,

$$x + 8z = 0$$

$$y + 2z = 9$$

solving for our leading variables, we get all solutions in terms of z for any $z \in \mathbb{F}_{17}$

$$x = 9z$$

$$y = 9 + 15z$$

□

Problem P3 Consider the positively oriented orthonormal vectors in $V = \mathbb{R}^3$:

$$v_1 = \frac{1}{\sqrt{2}}(1, -1, 0), v_2 = \frac{1}{\sqrt{3}}(1, 1, 1), \text{ and } v_3 = v_1 \times v_2$$

(the vector, or cross, product)

Let T be the rotation of $V = \mathbb{R}^3$ about the axis v_3 by 90°

(1) Compute the matrix $[T]_{\mathcal{B}'} = [T]_{\mathcal{B}'}^{\mathcal{B}'}$, with respect to the basis

$$\mathcal{B}' = (v_1, v_2, v_3)$$

(2) Compute the matrix of T with respect to the standard basis $\mathcal{B} = (e_1, e_2, e_3)$

Proof. First we must compute v_3 which evaluates to be,

$$v_3 = \left(-\frac{1}{\sqrt{6}}, -\frac{1}{\sqrt{6}}, \sqrt{\frac{2}{3}}\right) = \frac{1}{\sqrt{6}}(-1, -1, 2)$$

Now because we are rotating only 90 degrees about v_3 , we know v_3 should remain unchanged after our rotation. While $T(v_1) = v_2$ and $T(v_2) = -v_1$, putting this together we have,

$$T(v_1) = v_2$$

$$T(v_2) = -v_1$$

$$T(v_3) = v_3$$

which means we have $[T]_{\mathcal{B}'}^{\mathcal{B}'} = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Now let $P : V \rightarrow V$ be $P(v) = v$. Now we need to calculate $[P]_{\mathcal{B}'}^{\mathcal{B}}$,

$$P(v_1) = \frac{1}{\sqrt{2}}(1, -1, 0) = \frac{1}{\sqrt{2}}e_1 - \frac{1}{\sqrt{2}}e_2$$

$$P(v_2) = \frac{1}{\sqrt{3}}(1, 1, 1) = \frac{1}{\sqrt{3}}e_1 + \frac{1}{\sqrt{3}}e_2 + \frac{1}{\sqrt{3}}e_3$$

$$P(v_3) = \frac{1}{\sqrt{6}}(-1, -1, 2) = -\frac{1}{\sqrt{6}}e_1 - \frac{1}{\sqrt{6}}e_2 + \frac{2}{\sqrt{6}}e_3$$

all together gives us,

$$[P]_{\mathcal{B}'}^{\mathcal{B}} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{3}} & \frac{2}{\sqrt{6}} \end{bmatrix}$$

which then means,

$$\frac{1}{[P]_{\mathcal{B}'}^{\mathcal{B}}} = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{2} & 0 \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} \end{bmatrix}$$

We know then to compute the matrix of T with respect to the standard basis \mathcal{B} , recall though

$$[T]_{\mathcal{B}'}^{\mathcal{B}'} = \frac{1}{[P]_{\mathcal{B}'}^{\mathcal{B}}} [T]_{\mathcal{B}}^{\mathcal{B}} [P]_{\mathcal{B}'}^{\mathcal{B}},$$

so solving for $[T]_{\mathcal{B}}^{\mathcal{B}}$ we get,

$$[T]_{\mathcal{B}}^{\mathcal{B}} = [P]_{\mathcal{B}'}^{\mathcal{B}} [T]_{\mathcal{B}'}^{\mathcal{B}'} \frac{1}{[P]_{\mathcal{B}'}^{\mathcal{B}}},$$

Plugging in what we know we get,

$$\begin{aligned} [T]_{\mathcal{B}}^{\mathcal{B}} &= \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{3}} & \frac{2}{\sqrt{6}} \end{bmatrix} \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{2} & 0 \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{3}} & \frac{2}{\sqrt{6}} \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{3}} \\ \frac{1}{2} & -\frac{1}{2} & 0 \\ -\frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{6} & \frac{-2\sqrt{6}+1}{6} & \frac{-\sqrt{6}-3}{3\sqrt{6}} \\ \frac{2\sqrt{6}+1}{6} & \frac{1}{6} & \frac{3-\sqrt{6}}{3\sqrt{6}} \\ \frac{3-\sqrt{6}}{3\sqrt{6}} & \frac{-\sqrt{6}-3}{3\sqrt{6}} & \frac{2}{3} \end{bmatrix}. \end{aligned}$$

□