

Quantitative morphological characterization of pancreatic islets in HE-stained slides

Daniela F. Rodrigues¹✉, Tiago Bordeira Gaspar^{2,3,4,5}, Paula Sampaio¹, Mafalda Sousa¹

¹ Advanced Light Microscopy Scientific Platform, Institute for Research and Innovation in Health (i3s), University of Porto, Portugal
² Cancer Signalling & Metabolism group, Institute for Research and Innovation in Health (i3s), University of Porto, Portugal
³ Ipatimup – Institute of Molecular Pathology and Immunology of the University of Porto, Portugal
⁴ ICBAS – Institute of Biomedical Sciences Abel Salazar of the University of Porto, Portugal
⁵ FMUP – Medical Faculty of the University of Porto, Portugal

✉ drodrigues@ibmc.up.pt




1. Introduction

Pancreatic islets represent a network of endocrine cells that play an essential role in the maintenance of homeostatic processes. Quantitative assessment of islet morphology is of utmost importance to establish a relationship between morphological and functional changes. Here we propose an image analysis workflow for the quantitative morphological characterization of pancreatic islets in hematoxylin and eosin (HE) stained whole-slide images. As a proof-of-concept, this workflow was applied for characterization and discrimination of age and genotype-associated changes in islet morphology.


2. Methods

Longitudinal study







Islet morphological characterization

HE stained pancreas slides were scanned using NanoZoomer2.0HT (Hamamatsu) at 40x magnification with a resolution of 226 nm/pixel. A deep learning algorithm was trained in HALO® (Indica Labs) for the segmentation of pancreatic islets. The segmentation output was post-processed in Fiji and a set of measurements was extracted for the morphological characterization of pancreatic islets.



3. Results



The quantitative morphological characterization of pancreatic islets can help to identify age- and genotype-associated changes, allowing to reduce the subjectivity associated with this analysis

Acknowledgments

"This work was financed by FEDER - Fundo Europeu de Desenvolvimento Regional funds through the COMPETE 2020 - Operacional Programme for Competitiveness and Internationalisation (POCI), Portugal 2020, and by Portuguese funds through FCT - Fundação para a Ciência e a Tecnologia/Ministério da Ciência, Tecnologia e Ensino Superior in the framework of the project "Institute for Research and Innovation in Health Sciences" (POCI-01-0145-FEDER-007274) and the project "PPBI - Portuguese Platform of Bioimaging" (PPBI-POCI-01-0145-FEDER-022122)."

