# Følner sequence

#### Kai Prince SFHEA

2025-09-17

### 1 Introduction

Recall the definition of a group:

**Definition 1.1** (Group). A *group* is a non-empty set  $\Gamma$  together with a binary operation on  $\Gamma$ , denoted " $\cdot$ ", that combines any two elements  $\gamma$  and  $\gamma'$  of  $\Gamma$  to form an element of  $\Gamma$ , denoted  $\gamma \cdot \gamma'$ , such that the following three requirements, known as *group axioms*, are satisfied:

- Associativity: For all g, h, j in  $\Gamma$ , one has  $(g \cdot h) \cdot j = g \cdot (h \cdot j)$ .
- *Identity:* There exists an element e in  $\Gamma$  such that, for every g in  $\Gamma$ , one has  $e \cdot g = g$  and  $g \cdot e = g$ . Such an element is unique and is called the *identity element*.
- Unique Inverse: For each g in  $\Gamma$ , there exists an element h in  $\Gamma$  such that  $g \cdot h = e$  and  $h \cdot g = e$ , where e is the identity element. For each g, the element h is unique and is called the *inverse* of h and is denoted  $g^{-1}$ .

**Definition 1.2.** We define a *right-Følner sequence* in a group,  $\Gamma$ , as a sequence  $\Phi = (\Phi_N)_{N \in \mathbb{N}}$  of finite subsets of  $\Gamma$  satisfying

$$\lim_{N\to\infty}\frac{|(\Phi_N\cdot\gamma^{-1})\cdot\Phi_N|}{|\Phi_N|}=1$$

for all  $\gamma \in \Gamma$ .

**Definition 1.3.** Similarly, we define a *left-Følner sequence* in a group,  $\Gamma$ , as a sequence  $\Phi = (\Phi_N)_{N \in \mathbb{N}}$  of finite subsets of  $\Gamma$  satisfying

$$\lim_{N\to\infty}\frac{|(\gamma^{-1}\cdot\Phi_N)\cap\Phi_N|}{|\Phi_N|}=1$$

for all  $\gamma \in \Gamma$ .

**Definition 1.4.** We call a sequence a  $F\emptyset$ *lner sequence* if it is both a left and right F $\emptyset$ *lner sequence*.

A related definition is the following:

**Definition 1.5.** We call define density of a subset  $A \subseteq \Gamma$  with respect to a Følner sequence,  $\Phi$ , as

$$d_{\Phi}(A) = \lim_{N \to \infty} \frac{|\Phi_N \cap A|}{|\Phi_N|},$$

if it exists.

For  $\mathbb{N}$ , the *natural density*, d, is defined when the Følner sequence is constructed with  $\Phi_N = [1,...,N]$ .

## 2 Results

**Theorem 2.1** (The Test Theorem). This is a Theorem.



This is a test tip.

## 3 More Information

You can learn more about controlling the appearance of HTML output here: https://quarto.org/docs/output-formats/html-basics.html