

Addressing asymmetrical ties in ranking systems of contestants and judges using primes and irrational numbers

Nico Alberti, Kailey Hua,
Sophie Magaro, Narayan
Rueppel, Brendan Song

Proof School Math Burst 2020
Advisor: Dr. Susan Durst

Ranking system premise & initial setup

- Given n judges and k contestants, assign each contestant a rank from 1 through k .
- We identified a couple immediate issues with commonplace ranking systems:
 - If there is an even number of judges, after averaging their votes, it is possible for contestants to tie for a position in the ranking.
 - One of the judges is crazy and does not vote with reason or logic.
 - Do judges rank each contestant, provide a rating, or pick their favorite p contestants?

Even number of judges?

If there were an even number of judges ranking their contestants, we can arbitrarily ignore a judge's ranking, making an odd number of opinions.

Crazy judge?

For logic and simplicity, we chose to assume that all judges voted under reason.

If a judge really is crazy, we disregard their ranking.

How do the judges cast their votes?

- We started with judges assigning each contestant a ranking from 1-10.
- Ranking each contestant from 1 - k produced better results.

Proposed ranking system and point assignment

- Each judge produces a ranking of contestants from integers 1 through k , with 1 being their most favored contestant and k being their least favorite. Each contestant must receive a unique ranking, and each ranking 1, ..., k must be assigned to a contestant.
 - An example of computing the total scores for a contestant would be 1 point for k 'th place, 2 points for $k-1$ 'th, and so on.
 - The optimal way to do this is to assign each position a prime or an irrational number, making it impossible for them to sum to each other.
- **Symmetrical ties** occur when contestants get the same set of ranks from different judges.
 - We found that this is not preventable.
- **Asymmetrical ties** occur when the total scores of two contestants are equal, but not symmetrical.
 - This occurs if the point values may sum to each other, but using primes and irrationals to assign ranking points, this is not possible.

Examples of symmetrical and asymmetrical ties

- Given judges A, B, and C, give X the rankings 1, 2, 3 and Y the rankings 2, 3, 1, they would be in a symmetrical tie.
- Given judges A, B, and C, give X the rankings 5, 3, 2 and Y the rankings 1, 5, 4, they would be in an asymmetrical tie.

Non prime or irrational ranking system example

- Each ranking has a point value of its distance from $(k+1)/2$.
- For each additional time that someone is ranked at a certain rank, they receive their rank points, plus additional rank points