Nutrition Diagnosis Etiology Matrix

Below are the etiology categories and their definitions. Etiologies are grouped by the type of cause or contributing risk factor. In two specific instances, Access and Behavior etiologies, these alone may be the cause or contributing risk factor of the nutrition diagnosis or the practitioner may determine a more specific root cause, eg, Belief-Attitudes of the problem.

Etiology Category	Definition
Beliefs–Attitudes Etiologies	Cause or contributing risk factors related to the conviction of the truth of some nutrition-related statement or phenomenon; feelings or emotions toward that truth or phenomenon and activities.
Cultural Etiologies	Cause or contributing risk factors related to values, social norms, customs, religious beliefs and/or political systems.
Knowledge Etiologies	Cause or contributing risk factors impacting the level of understanding about food, nutrition and health, or nutrition-related information and guidelines.
Physical Function Etiologies	Cause or contributing risk factors related to physical ability to engage in specific tasks, may be cognitive in nature.
Physiologic–Metabolic Etiologies	Cause or contributing risk factors related to medical/health status that may have a nutritional impact (excludes psychological etiologies—see separate category).
Psychological Etiologies	Cause or contributing risk factors related to a diagnosed or suspected mental health/psychological problem (*Diagnostic and Statistical Manual of Mental Disorders*, DSM)
Social–Personal Etiologies	Cause or contributing risk factors associated with personal and/or social history.
Treatment Etiologies	Cause or contributing risk factors related to medical or surgical treatment or other therapies and management or care.
Access Etiologies	Cause or contributing risk factors that affect intake and the availability of safe, healthful food, water, and food/nutrition-related supplies. A more specific root cause of Access Etiologies may not be known but may eventually reveal Beliefs-Attitudes, Cultural, Knowledge, Physical Function, Psychological, Social-Personal, or Treatment Etiologies.
Behavior Etiologies	Cause or contributing risk factors related to actions which influence achievement of nutrition-related goals. A more specific root cause of Behavior Etiologies may not be known but may eventually reveal Beliefs-Attitudes, Cultural, Knowledge, Physical Function, Psychological, Social-Personal, or Treatment Etiologies.

Category	Etiology	Diagnosis
Beliefs-Attitudes	Altered body image	Poor nutrition quality of life (NB-2.5)
Beliefs-Attitudes	Obsessive desire to be healthy or have a specific body shape	Disordered eating pattern (NB-1.5)
Beliefs-Attitudes	Food preference	Limited food acceptance (NI-2.11), Excessive fat intake (NI-5.5.2), Intake of types of fats inconsistent with needs (*specify*) (NI-5.5.3), Undesirable food choices (NB-1.7)
Beliefs-Attitudes	Limited acceptance need to change	Not ready for diet/lifestyle change (NB-1.3)
Beliefs-Attitudes	Desire for a cure for a chronic disease through the use of alternative therapy	Unsupported beliefs/attitudes about food or nutrition related topics (NB-1.2)
Beliefs-Attitudes	Disbelief in science-based food and nutrition information	Unsupported beliefs/attitudes about food or nutrition related topics (NB-1.2)
Beliefs-Attitudes	End-of-life care if client or supportive individuals do not desire nutrition support	Enteral nutrition composition inconsistent with needs (NI-2.5), Enteral nutrition administration inconsistent with needs (NI-2.6), Parenteral nutrition composition inconsistent with needs (NI-2.9), Parenteral nutrition administration inconsistent with needs (NI-2.10)
Beliefs-Attitudes	Cultural, societal, biological/genetic, and/or environmental related to fear of weight gain	Disordered eating pattern (NB-1.5)
Beliefs-Attitudes	Food faddism	Imbalance of nutrients (NI-5.4), Excessive protein intake (NI-5.6.2), Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Excessive mineral intake (*specify*) (NI-5.10.2)
Beliefs-Attitudes	Limited food acceptance due to food aversion	Inadequate oral intake (NI-2.1), Limited food acceptance (NI-2.11),
Beliefs-Attitudes	Unsupported beliefs/attitudes about food, nutrition, and nutrition related information	Excessive energy intake (NI-1.3), Inadequate oral intake (NI-2.1), Excessive oral intake (NI-2.2), Limited food acceptance (NI-2.11), Excessive alcohol intake (NI-4.3), Imbalance of nutrients (NI-5.4), Excessive fat intake (NI-5.5.2), Intake of types of fats inconsistent with needs (*specify*) (NI-5.5.3), Excessive protein intake (NI-5.6.2), Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1), Excessive fiber intake (NI-5.8.6), Excessive mineral intake (*specify*) (NI-5.10.2), Predicted breastfeeding difficulty (NC-1.5), Underweight (NC-3.1), Growth rate below expected (NC-3.5), Food and nutrition related knowledge deficit (NB-1.1), Not ready for diet/lifestyle change (NB-1.3), Limited adherence to nutrition related recommendations (NB-1.6), Physical inactivity (NB-2.1), Excessive physical activity (NB-2.2), Limited access to food (NB-3.2), Limited access to potable water (NB-3.4)
Beliefs-Attitudes	Irritability	Breastfeeding difficulty (NC-1.3)
Beliefs-Attitudes	Lack of or limited self efficacy for making change or demoralization from previous challenges associated with change	Not ready for diet/lifestyle change (NB-1.3), Limited adherence to nutrition related recommendations (NB-1.6), Poor nutrition quality of life (NB-2.5)
Beliefs-Attitudes	Limited confidence in ability to change	Limited adherence to nutrition related recommendations (NB-1.6)
Beliefs-Attitudes	Limited value for behavior change or competing values	Excessive energy intake (NI-1.3), Excessive oral intake (NI-2.2), Excessive alcohol intake (NI-4.3), Excessive fat intake (NI-5.5.2), Intake of types of fats inconsistent with needs (*specify*) (NI-5.5.3), Self monitoring deficit (NB-1.4), Limited adherence to nutrition related recommendations (NB-1.6), Physical inactivity (NB-2.1), Inability to manage self care (NB-2.3)
Beliefs-Attitudes	Lacks motivation and/or readiness to apply or support systems change	Undesirable food choices (NB-1.7)

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
Beliefs-Attitudes	Negative impact of current or previous medical nutrition therapy (MNT)	Poor nutrition quality of life (NB-2.5)
Beliefs-Attitudes	Not ready for diet/lifestyle change	Overweight, adult or pediatric (NC-3.3.1), Obese, pediatric (NC-3.3.2), Obese, Class I (NC-3.3.3), Obese, Class II (NC-3.3.4), Obese, Class III (NC-3.3.5), Unintended weight gain (NC-3.4), Excessive growth rate (NC-3.6), Self monitoring deficit (NB-1.4), Inability to manage self care (NB-2.3), Poor nutrition quality of life (NB-2.5)
Beliefs-Attitudes	Perception of inadequate milk supply	Breastfeeding difficulty (NC-1.3), Predicted breastfeeding difficulty (NC-1.5)
Beliefs-Attitudes	Perception that lack of or limited resources (eg, time, financial, or interpersonal) prevent:	
	• Selection/food choices consistent with recommendations	Undesirable food choices (NB-1.7)
	• Changes	Not ready for diet/lifestyle change (NB-1.3), Limited adherence to nutrition related recommendations (NB-1.6)
	• Sufficient level of activity	Physical inactivity (NB-2.1)
	• Self monitoring	Self monitoring deficit (NB-1.4), Inability to manage self care (NB-2.3)
Beliefs-Attitudes	Unwilling or disinterested in:	
	• Learning/applying information	Food and nutrition related knowledge deficit (NB-1.1), Not ready for diet/lifestyle change (NB-1.3), Limited adherence to nutrition related recommendations (NB-1.6), Undesirable food choices (NB-1.7), Inability to manage self care (NB-2.3)
	• Reducing energy intake	Excessive energy intake (NI-1.3)
	• Reducing intake	Excessive oral intake (NI-2.2)
	• Modify protein or amino acid intake	Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1)
	• Tracking progress	Self monitoring deficit (NB-1.4)
	• Unwillingness to purchase or consume fiber-containing foods	Inadequate fiber intake (NI-5.8.5)
	• Weight regulation/preoccupation	Disordered eating pattern (NB-1.5)
Cultural	Culture of overeating	Predicted excessive energy intake (NI-1.5)
Cultural	Practices that affect nutrient intake	Predicted inadequate nutrient intake (*specify*) (NI-5.11.1)
Cultural	Cultural practices that affect ability to:	
	• Access to food, fluid, nutrients	Inadequate energy intake (NI-1.2), Inadequate oral intake (NI-2.1), Inadequate fluid intake (NI-3.1), Moderate starvation related malnutrition (undernutrition) (NC-4.1.1.1), Severe starvation related malnutrition (undernutrition) (NC-4.1.1.2), Inadequate protein–energy intake (NI-5.2), Inadequate protein intake (NI-5.6.1), Inadequate carbohydrate intake (NI-5.8.1), Inadequate vitamin intake (*specify*) (NI-5.9.1), Inadequate mineral intake (*specify*) (NI-5.10.1), Unintended weight loss (NC-3.2), Limited access to food (NB-3.2), Limited access to potable water (NB-3.4)
	• Make appropriate food choices	Inadequate fat intake (NI-5.5.1)
	• Breastfeed	Breastfeeding difficulty (NC-1.3), Predicted breastfeeding difficulty (NC-1.5)
	• Learn/apply information	Food and nutrition related knowledge deficit (NB-1.1), Undesirable food choices (NB-1.7)

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
	• Manage self care	Inability to manage self care (NB-2.3)
	• Reduce carbohydrate intake	Excessive carbohydrate intake (NI-5.8.2)
	• Regulate types of protein or amino acids consumed	Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1)
	• Regulate timing of carbohydrate consumption	Inconsistent carbohydrate intake (NI-5.8.4)
	• Regulate types of carbohydrate consumed	Intake of types of carbohydrate inconsistent with needs (*specify*) (NI-5.8.3)
	• Track personal progress	Self monitoring deficit (NB-1.4)
Knowledge	Food and nutrition knowledge deficit	Inadequate energy intake (NI-1.2), Excessive energy intake (NI-1.3), Excessive oral intake (NI-2.2), Inadequate fluid intake (NI-3.1), Excessive fluid intake (NI-3.2), Inadequate bioactive substance intake (*specify*) (NI-4.1), Excessive bioactive substance intake (*specify*) (NI-4.2), Excessive alcohol intake (NI-4.3), Moderate starvation related malnutrition (undernutrition) (NC-4.1.1.1), Severe starvation related malnutrition (undernutrition) (NC-4.1.1.2), Moderate acute disease or injury related malnutrition (undernutrition) (NC-4.1.3.1), Severe acute disease or injury related malnutrition (undernutrition) (NC-4.1.3.2), Inadequate protein–energy intake (NI-5.2), Imbalance of nutrients (NI-5.4), Excessive fat intake (NI-5.5.2), Inadequate protein intake (NI-5.6.1), Excessive protein intake (NI-5.6.2), Inadequate carbohydrate intake (NI-5.8.1), Inadequate fiber intake (NI-5.8.5), Excessive mineral intake (*specify*) (NI-5.10.2), Predicted breastfeeding difficulty (NC-1.5), Overweight, adult or pediatric (NC-3.3.1), Obese, pediatric (NC-3.3.2), Obese, Class I (NC-3.3.3), Obese, Class II (NC-3.3.4), Obese, Class III (NC-3.3.5), Excessive growth rate (NC-3.6), Self monitoring deficit (NB-1.4), Inability to manage self care (NB-2.3), Poor nutrition quality of life (NB-2.5), Limited access to food (NB-3.2), Limited access to nutrition related supplies (NB-3.3), Limited access to potable water (NB-3.4)
Knowledge	Food and nutrition knowledge deficit concerning:	
	• Sufficient oral food/beverage intake	Inadequate oral intake (NI-2.1)
	• Consumption of an appropriate variety of foods	Excessive mineral intake (*specify*) (NI-5.10.2)
	• Potentially unsafe food	Intake of unsafe food (NB-3.1)
	• Proper infant feeding, food/feeding preparation and storage	Intake of unsafe food (NB-3.1)
	• Adequate energy intake	Underweight (NC-3.1)
	• Appropriate amount or types of dietary protein or amino acids	Mild non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.1), Moderate non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.2), Severe non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.3), Inadequate protein–energy intake (NI-5.2), Inadequate protein intake (NI-5.6.1), Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1)
	• Appropriate amount or type of dietary fat	Inadequate protein–energy intake (NI-5.2), Inadequate fat intake (NI-5.5.1), Excessive fat intake (NI-5.5.2), Intake of types of fats inconsistent with needs (*specify*) (NI-5.5.3)

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
	• Appropriate amount and types of dietary carbohydrate	Excessive carbohydrate intake (NI-5.8.2), Intake of types of carbohydrate inconsistent with needs (*specify*) (NI-5.8.3)
	• Appropriate timing of carbohydrate intake	Inconsistent carbohydrate intake (NI-5.8.4)
	• Physiological causes requiring careful timing and consistency in the amount of carbohydrate	Inconsistent carbohydrate intake (NI-5.8.4)
	• Physiological causes altering carbohydrate digestion or metabolism	Intake of types of carbohydrate inconsistent with needs (*specify*) (NI-5.8.3)
	• Physiological causes requiring use of modified carbohydrate intake	Excessive carbohydrate intake (NI-5.8.2)
	• Desirable quantities of fiber	Inadequate fiber intake (NI-5.8.5), Excessive fiber intake (NI-5.8.6)
	• Correct enteral formula needed	Inadequate enteral nutrition infusion (NI-2.3)
	• Food and supplemental sources of vitamins	Inadequate vitamin intake (*specify*) (NI-5.9.1), Excessive vitamin intake (*specify*) (NI-5.9.2)
	• Food and supplemental sources of minerals	Inadequate mineral intake (*specify*) (NI-5.10.1)
	• Recommended dose of vitamin and mineral supplements	Imbalance of nutrients (NI-5.4)
	• Management of diagnosis requiring mineral restriction	Excessive mineral intake (*specify*) (NI-5.10.2)
	• Management of diagnosed genetic disorder altering mineral homeostasis	Excessive mineral intake (*specify*) (NI-5.10.2)
	• Correct amount of enteral/parenteral formula	Excessive energy intake (NI-1.3) Excessive parenteral nutrition infusion (NI-2.8)
	• Correct parenteral nutrition components or administration	Inadequate parenteral nutrition infusion (NI-2.7), Parenteral nutrition composition inconsistent with needs (NI-2.9), Parenteral nutrition administration inconsistent with needs (NI-2.10)
	• Appropriate/correct access for delivering EN/PN	Inadequate enteral nutrition infusion (NI-2.3), Inadequate parenteral nutrition infusion (NI-2.7), Parenteral nutrition administration inconsistent with needs (NI-2.10)
	• Health benefits of physical activity	Physical inactivity (NB-2.1)
	• How to make nutrition related changes	Limited adherence to nutrition related recommendations (NB-1.6)
	• Food drug interactions	Food medication interaction (NC-2.3), Predicted food medication interaction (NC-2.4)
	• On the part of the caregiver	Excessive enteral nutrition infusion (NI-2.4), Excessive parenteral nutrition infusion (NI-2.8), Enteral nutrition composition inconsistent with needs (NI-2.5), Enteral nutrition administration inconsistent with needs (NI-2.6), Parenteral nutrition composition inconsistent with needs (NI-2.9), Parenteral nutrition administration inconsistent with needs (NI-2.10)
	• Consumption of high-dose nutrient supplements	Imbalance of nutrients (NI-5.4)
	• Infant/child hunger cues	Food and nutrition related knowledge deficit (NB-1.1)
Knowledge	Lack of or limited prior exposure or exposure to inaccurate nutrition related information	Food and nutrition related knowledge deficit (NB-1.1), Unsupported beliefs/attitudes about food or nutrition related topics (NB-1.2), Self monitoring deficit (NB-1.4),

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
		Undesirable food choices (NB-1.7), Inability to manage self care (NB-2.3)
Knowledge	Lack of or limited prior exposure to accurate information regarding physical activity	Physical inactivity (NB-2.1)
Knowledge	Limited adjustment for lifestyle changes or restricted mobility and decreased metabolism	Excessive energy intake (NI-1.3)
Physical function	Irritability	Breastfeeding difficulty (NC-1.3)
Physical function	Inability to physically: • Bend elbow at wrist • Grasp cups and utensils • Sit with hips square and back straight • Support neck and/or control head and neck • Coordinate hand movement to mouth	Self feeding difficulty (NB-2.6)
Physical function	Lack of or limited self feeding ability	Unintended weight loss (NC-3.2)
Physical function	Diminished ability to shop	Limited access to food (NB-3.2), Limited access to nutrition related supplies (NB-3.3), Limited access to potable water (NB-3.4)
Physical function	Limited physical strength or range of motion	Self feeding difficulty (NB-2.6)
Physical function	Physical inactivity	Overweight, adult or pediatric (NC-3.3.1), Obese, pediatric (NC-3.3.2), Obese, Class I (NC-3.3.3), Obese, Class II (NC-3.3.4), Obese, Class III (NC-3.3.5), Excessive growth rate (NC-3.6)
Physical function	Change in physical activity anticipated	Predicted inadequate energy intake (NI-1.4), Predicted excessive energy intake (NI-1.5)
Physical function	Voluntary or involuntary physical activity/movement	Increased energy expenditure (NI-1.1)
Physiologic-Metabolic	Age-related demands	Inadequate protein intake (NI-5.6.1), Growth rate below expected (NC-3.5)
Physiologic-Metabolic	Alteration in gastrointestinal tract:	
	• Decreased functional length of GI tract	Increased nutrient needs (*specify*) (NI-5.1), Moderate starvation related malnutrition (undernutrition) (NC-4.1.1.1), Severe starvation related malnutrition (undernutrition) (NC-4.1.1.2), Moderate chronic disease or condition related malnutrition (undernutrition) (NC-4.1.2.1), Severe chronic disease or condition related malnutrition (NC-4.1.2.2), Moderate acute disease or injury related malnutrition (undernutrition) (NC-4.1.3.1), Severe acute disease or injury related malnutrition (undernutrition) (NC-4.1.3.2), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.1), Moderate illness related pediatric malnutrition (NC-4.1.5.2), Severe illness related pediatric malnutrition (NC-4.1.5.3), Altered GI function (NC-1.4), Growth rate below expected (NC-3.5)
	• Alteration in GI anatomical structure	Increased nutrient needs (*specify*) (NI-5.1), Moderate starvation related malnutrition (undernutrition) (NC-4.1.1.1), Severe starvation related malnutrition (undernutrition) (NC-4.1.1.2), Moderate chronic disease or condition related malnutrition (undernutrition) (NC-4.1.2.1), Severe chronic disease or condition related malnutrition (NC-4.1.2.2),Moderate acute disease or injury related malnutrition (undernutrition) (NC-4.1.3.1), Severe acute disease or injury related malnutrition

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
		(undernutrition) (NC-4.1.3.2), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3), Altered GI function (NC-1.4), Inadequate fat intake (NI-5.5.1), Growth rate below expected (NC-3.5)
	• Alteration in GI function	Limited food acceptance (NI-2.11), Inadequate bioactive substance intake (*specify*) (NI-4.1), Excessive bioactive substance intake (*specify*) (NI-4.2), Increased nutrient needs (*specify*) (NI-5.1), Moderate starvation related malnutrition (undernutrition) (NC-4.1.1.1), Severe starvation related malnutrition (undernutrition) (NC-4.1.1.2), Moderate chronic disease or condition related malnutrition (undernutrition) (NC-4.1.2.1), Severe chronic disease or condition related malnutrition (NC-4.1.2.2),Moderate acute disease or injury related malnutrition (undernutrition) (NC-4.1.3.1), Severe acute disease or injury related malnutrition (undernutrition) (NC-4.1.3.2), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3),, Inadequate fat intake (NI-5.5.1), Altered GI function (NC-1.4), Growth rate below expected (NC-3.5)
	• Change in GI tract motility	Altered GI function (NC-1.4)
	• Change in GI related organ function	Increased nutrient needs (*specify*) (NI-5.1), Altered GI function (NC-1.4), Growth rate below expected (NC-3.5)
	• Compromised endocrine function	Impaired nutrient utilization (NC-2.1), Growth rate below expected (NC-3.5)
Physiologic-Metabolic	Altered cholesterol metabolism/regulation	Decreased nutrient needs (*specify*) (NI-5.3)
Physiologic-Metabolic	Breast or nipple abnormality	Breastfeeding difficulty (NC-1.3), Predicted breastfeeding difficulty (NC-1.5)
Physiologic-Metabolic	Changes in taste, appetite	Excessive fat intake (NI-5.5.2), Intake of types of fats inconsistent with needs (*specify*) (NI-5.5.3)
Physiologic-Metabolic	Conditions leading to excess fluid loss	Inadequate fluid intake (NI-3.1)
Physiologic-Metabolic	Craniofacial malformations	Biting/Chewing (masticatory) difficulty (NC-1.2)
Physiologic-Metabolic	Decreased energy needs	Overweight, adult or pediatric (NC-3.3.1), Obese, pediatric (NC-3.3.2), Obese, Class I (NC-3.3.3), Obese, Class II (NC-3.3.4), Obese, Class III (NC-3.3.5), Excessive growth rate (NC-3.6)
Physiologic-Metabolic	Decreased ability to consume sufficient energy, nutrients	Inadequate energy intake (NI-1.2), Inadequate oral intake (NI-2.1), Inadequate protein–energy intake (NI-5.2), Inadequate protein intake (NI-5.6.1), Inadequate vitamin intake (*specify*) (NI-5.9.1), Inadequate mineral intake (*specify*) (NI-5.10.1), Unintended weight loss (NC-3.2), Growth rate below expected (NC-3.5)
Physiologic-Metabolic	Decreased nutrient needs related to low activity levels due to chronic disease or organ failure	Excessive enteral nutrition infusion (NI-2.4), Excessive parenteral nutrition infusion (NI-2.8), Excessive vitamin intake (*specify*) (NI-5.9.2)
Physiologic-Metabolic	Decreased total fat need or recommendation	Excessive fat intake (NI-5.5.2)
Physiologic-Metabolic	Altered fatty acid need or recommendation	Intake of types of fats inconsistent with needs (*specify*) (NI-5.5.3)
Physiologic-Metabolic	Developmental delay	Limited food acceptance (NI-2.11)

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
Physiologic-Metabolic	Difficulty chewing or swallowing high-fiber foods	Inadequate fiber intake (NI-5.8.5)
Physiologic-Metabolic	Difficulty latching on	Breastfeeding difficulty (NC-1.3)
Physiologic-Metabolic	Excessive energy intake	Overweight, adult or pediatric (NC-3.3.1), Obese, pediatric (NC-3.3.2), Obese, Class I (NC-3.3.3), Obese, Class II (NC-3.3.4), Obese, Class III (NC-3.3.5), Excessive growth rate (NC-3.6)
Physiologic-Metabolic	Food allergies and aversions impeding food choices consistent with guidelines	Undesirable food choices (NB-1.7)
Physiologic-Metabolic	Food intolerances	Decreased nutrient needs (*specify*) (NI-5.3)
Physiologic-Metabolic	Genetic predisposition to overweight/obesity	Predicted excessive energy intake (NI-1.5)
Physiologic-Metabolic	Heart failure	Decreased nutrient needs (*specify*) (NI-5.3)
Physiologic-Metabolic	Illness causing unexpected weight gain because of head trauma, immobility, paralysis or related condition	Unintended weight gain (NC-3.4)
Physiologic-Metabolic	Impaired cognitive ability, including learning disabilities, neurological or sensory impairment, and dementia	Inadequate fluid intake (NI-3.1), Moderate starvation related malnutrition (undernutrition) (NC-4.1.1.1), Severe starvation related malnutrition (undernutrition) (NC-4.1.1.2), Food and nutrition related knowledge deficit (NB-1.1), Not ready for diet/lifestyle change (NB-1.3), Self monitoring deficit (NB-1.4), Undesirable food choices (NB-1.7), Inability to manage self care (NB-2.3), Impaired ability to prepare foods/meals (NB-2.4), Self feeding difficulty (NB-2.6), Limited access to food (NB-3.2)
Physiologic-Metabolic	Inadequate energy intake	Underweight (NC-3.1), Moderate chronic disease or condition related malnutrition (undernutrition) (NC-4.1.2.1), Severe chronic disease or condition related malnutrition (NC-4.1.2.2), Moderate acute disease or injury related malnutrition (undernutrition) (NC-4.1.3.1), Severe acute disease or injury related malnutrition (undernutrition) (NC-4.1.3.2),
Physiologic-Metabolic	Inadequate milk supply	Breastfeeding difficulty (NC-1.3)
Physiologic-Metabolic	Increased energy needs	Underweight (NC-3.1)
Physiologic-Metabolic	Injury, condition, physical disability or limitation that reduces physical activity or activities of daily living	Physical inactivity (NB-2.1), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3),
Physiologic-Metabolic	Intolerance of EN/PN	Inadequate enteral nutrition infusion (NI-2.3), Inadequate parenteral nutrition infusion (NI-2.7), Mild non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.1), Moderate non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.2), Severe non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.3), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3),

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
Physiologic-Metabolic	Kidney, liver, cardiac, endocrine, neurologic, and/or pulmonary dysfunction	Limited food acceptance (NI-2.11), Excessive fluid intake (NI-3.2), Biting/ Chewing (masticatory) difficulty (NC-1.2), Altered nutrition related laboratory values (*specify*) (NC-2.2), Growth rate below expected (NC-3.5)
Physiologic-Metabolic	Lack of or limited developmental readiness to perform self management tasks	Inability to manage self care (NB-2.3)
Physiologic-Metabolic	Lethargy, sleepiness	Breastfeeding difficulty (NC-1.3)
Physiologic-Metabolic	Limited vision	Self feeding difficulty (NB-2.6)
Physiologic-Metabolic	Limited food acceptance	Inadequate oral intake (NI-2.1), Growth rate below expected (NC-3.5)
Physiologic-Metabolic	Liver dysfunction	Decreased nutrient needs (*specify*) (NI-5.3), Excessive protein intake (NI-5.6.2), Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1)
Physiologic-Metabolic	Loss of appetite awareness	Excessive oral intake (NI-2.2)
Physiologic-Metabolic	Malnutrition/malabsorption	Increased nutrient needs (*specify*) (NI-5.1), Breastfeeding difficulty (NC-1.3), Predicted breastfeeding difficulty (NC-1.5), Growth rate below expected (NC-3.5)
Physiologic-Metabolic	Mastitis and/or painful breasts, nipples	Breastfeeding difficulty (NC-1.3), Predicted breastfeeding difficulty (NC-1.5)
Physiologic-Metabolic	Mechanical issues such as inflammation, surgery, stricture, or oral, pharyngeal and esophageal tumors, mechanical ventilation	Swallowing difficulty (NC-1.1), Biting/ Chewing (masticatory) difficulty (NC-1.2)
Physiologic-Metabolic	Inborn errors of metabolism	Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1), Altered nutrition related laboratory values (*specify)* (NC-2.2), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3),
Physiologic-Metabolic	Metabolic abnormality	Excessive protein intake (NI-5.6.2), Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3),
Physiologic-Metabolic	Metabolic disorders	Impaired nutrient utilization (NC-2.1), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3),
Physiologic-Metabolic	Motor causes related to neurological or muscular disorders	Swallowing difficulty (NC-1.1)
Physiologic-Metabolic	Oral pain	Breastfeeding difficulty (NC-1.3)
Physiologic-Metabolic	Other organ dysfunction that leads to biochemical changes	Altered nutrition related laboratory values (*specify*) (NC-2.2)
Physiologic-Metabolic	Partial or complete edentulism	Biting/Chewing (masticatory) difficulty (NC-1.2)

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
Physiologic-Metabolic	Physical disability	Impaired ability to prepare foods/meals (NB-2.4), Limited access to food (NB-3.2), Limited access to potable water (NB-3.4)
Physiologic-Metabolic	Physiological causes that put emphasis on food, weight, or shape	Disordered eating pattern (NB-1.5)
Physiologic-Metabolic	Physiologic causes requiring modified amount or timing of carbohydrate intake	Excessive carbohydrate intake (NI-5.8.2), Intake of types of carbohydrate inconsistent with needs (*specify*) (NI-5.8.3), Inconsistent carbohydrate intake (NI-5.8.4)
Physiologic-Metabolic	Physiological causes increasing nutrient needs due to:	
Physiologic-Metabolic	• Accelerated growth or anabolism	Increased energy expenditure (NI-1.1), Inadequate enteral nutrition infusion (NI-2.3), Inadequate parenteral nutrition infusion (NI-2.7), Increased nutrient needs (*specify*) (NI-5.1), Inadequate mineral intake (*specify*)(NI-5.10.1), Growth rate below expected (NC-3.5)
	• Altered absorption or metabolism	Inadequate fluid intake (NI-3.1), Increased nutrient needs (*specify*) (NI-5.1), Inadequate protein intake (NI-5.6.1), Inadequate carbohydrate intake (NI-5.8.1), Inadequate vitamin intake (*specify*) (NI-5.9.1), Inadequate mineral intake (*specify*) (NI-5.10.1), Predicted inadequate nutrient intake (*specify*) (NI-5.11.1), Moderate chronic disease or condition related malnutrition (undernutrition) (NC-4.1.2.1), Severe chronic disease or condition related malnutrition (NC-4.1.2.2),Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3),Inadequate protein–energy intake (NI-5.2), Unintended weight loss (NC-3.2), Growth rate below expected (NC-3.5)
	• Disease/condition	Inadequate enteral nutrition infusion (NI-2.3), Inadequate parenteral nutrition infusion (NI-2.7), Inadequate fluid intake (NI-3.1), Increased nutrient needs (*specify*) (NI-5.1), Moderate chronic disease or condition related malnutrition (undernutrition) (NC-4.1.2.1), Severe chronic disease or condition related malnutrition (NC-4.1.2.2),Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3), Inadequate protein–energy intake (NI-5.2), Inadequate protein intake (NI-5.6.1), Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1), Inadequate vitamin intake (*specify*) (NI-5.9.1), Unintended weight loss (NC-3.2), Growth rate below expected (NC-3.5)
	• Maintenance of body temperature	Increased energy expenditure (NI-1.1), Inadequate fluid intake (NI-3.1), Growth rate below expected (NC-3.5)
	• Prolonged catabolic illness	Inadequate energy intake (NI-1.2), Inadequate oral intake (NI-2.1), Inadequate fluid intake (NI-3.1), Moderate chronic disease or condition related malnutrition (undernutrition) (NC-4.1.2.1), Severe chronic disease or condition related malnutrition (NC-4.1.2.2),Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3),Inadequate protein–energy intake (NI-5.2), Inadequate protein intake (NI-5.6.1), Inadequate vitamin

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
		intake (*specify*) (NI-5.9.1), Inadequate mineral intake (*specify*) (NI-5.10.1), Unintended weight loss (NC-3.2), Growth rate below expected (NC-3.5)
Physiologic-Metabolic	Altered metabolism	Predicted excessive energy intake (NI-1.5), Predicted excessive nutrient intake (NI-5.11.2)
Physiologic-Metabolic	Poor sucking ability	Breastfeeding difficulty (NC-1.3), Predicted breastfeeding difficulty (NC-1.5), Growth rate below expected (NC-3.5)
Physiologic-Metabolic	Prematurity	Altered nutrition related laboratory values (NC-2.2), Growth rate below expected (NC-3.5), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3),
Physiologic-Metabolic	Renal dysfunction	Decreased nutrient needs (*specify*) (NI-5.3), Excessive protein intake (NI-5.6.2), Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1)
Physiologic-Metabolic	Small for gestational age, intrauterine growth retardation/restriction and/or Lack of or limited progress/appropriate weight gain per day	Underweight (NC-3.1), Growth rate below expected (NC-3.5)
Physiologic-Metabolic	Soft tissue disease (primary or oral manifestations of a systemic disease)	Biting/Chewing (masticatory) difficulty (NC-1.2), Moderate starvation related malnutrition (undernutrition) (NC-4.1.1.1), Severe starvation related malnutrition (undernutrition) (NC-4.1.1.2),
Physiologic-Metabolic	Swallowing difficulty, and altered suck and breathing patterns in infants	Swallowing difficulty (NC-1.1), Breastfeeding difficulty (NC-1.3), Predicted breastfeeding difficulty (NC-1.5)
Physiologic-Metabolic	Xerostomia	Biting/Chewing (masticatory) difficulty (NC-1.2)
Psychological	Alcohol or drug addiction	Excessive alcohol intake (NI-4.3), Impaired nutrient utilization (NC-2.1)
Psychological	Addictive personality	Excessive physical activity (NB-2.2)
Psychological	Mental illness, confusion, or altered awareness	Intake of unsafe food (NB-3.1), Excessive oral intake (NI-2.2)
Psychological	Personality characteristics or temperament associated with eating disorders	Disordered eating pattern (NB-1.5)
Psychological	Psychological causes that put emphasis on food, weight, or shape	Disordered eating pattern (NB-1.5)
Psychological	Psychological causes such as depression and disordered eating	Inadequate energy intake (NI-1.2), Inadequate oral intake (NI-2.1), Inadequate fluid intake (NI-3.1), Excessive fluid intake (NI-3.2), Moderate starvation related malnutrition (undernutrition) (NC-4.1.1.1), Severe starvation related malnutrition (undernutrition) (NC-4.1.1.2), Non illness related pediatric malnutrition (undernutrition) (NC-4.1.4), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3), Inadequate protein–energy intake (NI-5.2), Inadequate fat intake (NI-5.5.1), Inadequate protein intake (NI-5.6.1), Inadequate carbohydrate intake (NI-5.8.1), Excessive carbohydrate intake (NI-5.8.2), Intake of types of carbohydrate inconsistent with needs (*specify*) (NI-5.8.3), Inconsistent carbohydrate intake (NI-5.8.4), Inadequate fiber intake (NI-5.8.5), Inadequate vitamin intake (*specify*) (NI-5.9.1), Excessive vitamin intake (*specify*) (NI-5.9.2), Inadequate mineral intake (*specify*)

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
		(NI-5.10.1), Unintended weight loss (NC-3.2), Growth rate below expected (NC-3.5), Undesirable food choices (NB-1.7), Excessive physical activity (NB-2.2), Limited access to food (NB-3.2), Limited access to nutrition related supplies (NB-3.3)
Social-Personal	Lack of or limited role models	Breastfeeding difficulty (NC-1.3), Physical inactivity (NB-2.1)
Social-Personal	Lack of or limited social support for implementing changes	Breastfeeding difficulty (NC-1.3), Predicted breastfeeding difficulty (NC-1.5), Mild non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.1), Moderate non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.2), Severe non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.3), Not ready for diet/lifestyle change (NB-1.3), Self monitoring deficit (NB-1.4), Limited adherence to nutrition related recommendations (NB-1.6), Physical inactivity (NB-2.1), Inability to manage self care (NB-2.3), Poor nutrition quality of life (NB-2.5)
Social-Personal	Family or social history of overeating	Predicted excessive energy intake (NI-1.5)
Social-Personal	Increased psychological/life stress	Predicted breastfeeding difficulty (NC-1.5), Overweight, adult or pediatric (NC-3.3.1), Obese, pediatric (NC-3.3.2), Obese, Class I (NC-3.3.3), Obese, Class II (NC-3.3.4), Obese, Class III (NC-3.3.5), Predicted inadequate energy intake (NI-1.4), Predicted excessive energy intake (NI-1.5)
Social-Personal	Traumatic event(s) that causes a physical or psychological stress reaction	Disordered eating pattern (NB-1.5)
Social-Personal	Change in living situation	Predicted inadequate energy intake (NI-1.4), Predicted excessive energy intake (NI-1.5), Predicted inadequate nutrient intake (*specify*) (NI-5.11.1), Predicted breastfeeding difficulty (NC-1.5)
Social-Personal	Living in a geographic location with danger for environmental emergency	Predicted inadequate nutrient intake (*specify*) (NI-5.11.1)
Treatment	Accidental vitamin and/or mineral overdose from oral, enteral or parenteral sources	Excessive vitamin intake (*specify*) (NI-5.9.2), Excessive mineral intake (*specify*) (NI-5.10.2)
Treatment	Calories/kcal/kJ unaccounted for from IV infusion and/or medications	Excessive energy intake (NI-1.3)
Treatment	Changes in taste, appetite	Excessive fat intake (NI-5.5.2), Intake of types of fats inconsistent with needs (*specify*) (NI-5.5.3)
Treatment	Changes in GI tract motility	Altered GI function (NC-1.4)
Treatment	Chronic use of medications known to cause weight gain, such as use of certain antidepressants, antipsychotics, corticosteroids, certain HIV medications	Unintended weight gain (NC-3.4), Excessive growth rate (NC-3.6)
Treatment	Difficulty chewing or swallowing high-fiber foods	Inadequate fiber intake (NI-5.8.5)
Treatment	Excessive energy intake	Overweight, adult or pediatric (NC-3.3.1), Obese, pediatric (NC-3.3.2), Obese, Class I (NC-3.3.3), Obese, Class II (NC-3.3.4), Obese, Class III (NC-3.3.5),
Treatment	Food intolerances	Decreased nutrient needs (*specify*) (NI-5.3)
Treatment	High level of fatigue or other side effect of therapy	Undesirable food choices (NB-1.7), Impaired ability to prepare foods/meals (NB-2.4)
Treatment	Improvement in client status, allowing return to total or partial oral diet; changes in the course of disease resulting in changes in nutrient requirements	Enteral nutrition composition inconsistent with needs (NI-2.5), Enteral nutrition administration inconsistent with needs (NI-2.6), Parenteral nutrition composition inconsistent with needs (NI-2.9), Parenteral nutrition administration inconsistent with needs (NI-2.10)

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
Treatment	Inadequate energy intake	Underweight (NC-3.1)
Treatment	Infusion volume not reached or schedule for infusion interrupted	Inadequate enteral nutrition infusion (NI-2.3), Inadequate parenteral nutrition infusion (NI-2.7)
Treatment	Insufficient electrolyte replacement when initiating feeding (PN/EN, including oral)	Imbalance of nutrients (NI-5.4)
Treatment	Lack of or limited, compromised, or incorrect access for delivering EN/PN	Inadequate enteral nutrition infusion (NI-2.3), Inadequate parenteral nutrition infusion (NI-2.7)
Treatment	Mechanical issues such as inflammation, surgery, stricture, or oral, pharyngeal and esophageal tumors, mechanical ventilation	Swallowing difficulty (NC-1.1), Biting/ Chewing (masticatory) difficulty (NC-1.2)
Treatment	Medications that increase appetite	Excessive energy intake (NI-1.3), Excessive oral intake (NI-2.2),
Treatment	Medications that increase nutrient needs	Increased nutrient needs (*specify*) (NI-5.1)
Treatment	Medications that affect nutrient metabolism	Impaired nutrient utilization (NC-2.1)
Treatment	Medications that increase fluid needs or decrease thirst	Inadequate fluid intake (NI-3.1)
Treatment	Misused specialized protein products	Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3)
Treatment	Misused specialized amino acid products	Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1)
Treatment	Misdiagnosis of lactose intolerance/lactase deficiency	Inadequate mineral intake (*specify*) (NI-5.10.1)
Treatment	Nutrient/nutrient interaction and/or drug/nutrient interaction	Inadequate enteral nutrition infusion (NI-2.3), Inadequate parenteral nutrition infusion (NI-2.7), Inadequate vitamin intake (*specify*) (NI-5.9.1), Inadequate mineral intake (*specify*) (NI-5.10.1), Food Medication Interaction (NC-2.3), Predicted food medication interaction (NC-2.4)
Treatment	Overfeeding of parenteral/enteral nutrition (PN/EN)	Excessive energy intake (NI-1.3)
Treatment	Planned procedure, therapy or medication predicted to increase energy expenditure or nutrient need	Predicted inadequate energy intake (NI-1.4), Predicted inadequate nutrient intake (*specify*) (NI-5.11.1)
Treatment	Planned therapy or medication predicted to reduce energy/nutrient need or metabolic rate/metabolism	Predicted excessive energy intake (NI-1.5), Predicted excessive nutrient intake (NI-5.11.2)
Treatment	Planned procedure, therapy or medication predicted to hinder breastfeeding	Predicted breastfeeding difficulty (NC-1.5)
Treatment	Planned procedure, therapy or medication predicted to decrease ability to consume sufficient energy or nutrients	Predicted inadequate energy intake (NI-1.4), Predicted inadequate nutrient intake (*specify*) (NI-5.11.1)
Treatment	Prolonged adherence to a low-fiber or low-residue diet	Inadequate fiber intake (NI-5.8.5)
Treatment	Prolonged hospitalization	Unintended weight loss (NC-3.2)
Treatment	Renal dysfunction	Decreased nutrient needs (*specify*) (NI-5.3), Excessive protein intake (NI-5.6.2), Intake of types of proteins inconsistent with needs (*specify*) (NI-5.6.3), Intake types of amino acids inconsistent with needs (*specify*) (NI-5.7.1)
Treatment	Xerostomia	Biting/Chewing (masticatory) difficulty (NC-1.2)
Access	Access to foods and supplements in excess of needs	Excessive vitamin intake (*specify*) (NI-5.9.2)
Access	Caregiver intentionally or unintentionally not providing access to food or nutrition related supplies	Limited access to food (NB-3.2), Limited access to nutrition related supplies (NB-3.3), Limited access to potable water (NB-3.4)

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
Access	Community and geographical constraints	Inadequate vitamin intake (*specify*) (NI-5.9.1), Limited access to food (NB-3.2), Limited access to nutrition related supplies (NB-3.3), Limited access to potable water (NB-3.4)
Access	Environmental causes (eg, inadequately tested nutrient bioavailability of fortified foods, beverages, and supplements; marketing of fortified foods, beverages, supplements as a substitute for natural food source of nutrient(s))	Inadequate vitamin intake (NI-5.9.1), Inadequate mineral intake (*specify*) (NI-5.10.1)
Access	Exposure to contaminated water or food (eg, community outbreak of illness documented by surveillance and/or response agency)	Intake of unsafe food (NB-3.1), Limited access to food (NB-3.2), Limited access to potable water (NB-3.4)
Access	Reluctance to participate in federal food programs such as WIC, National School Breakfast/Lunch Program, food stamps	Limited access to food (NB-3.2), Limited access to potable water (NB-3.4)
Access	Financial constraints that may prevent sufficient level of activity (eg, to address cost of equipment or shoes or club membership to gain access)	Physical inactivity (NB-2.1)
Access	Lack of, or limited access to:	
	• Adaptive foods or eating devices conducive for self feeding	Self feeding difficulty (NB-2.6)
	• Available and safe exercise environment and/or equipment	Physical inactivity (NB-2.1)
	• Fluid	Inadequate fluid intake (NI-3.1), Inadequate fiber intake (NI-5.8.5)
	• Fortified foods and beverages	Inadequate mineral intake (*specify*) (NI-5.10.1)
	• Specialized protein products	Excessive protein intake (NI-5.6.2)
	• Food or artificial nutrition	Inadequate energy intake (NI-1.2), Inadequate oral intake (NI-2.1), Moderate starvation related malnutrition (undernutrition) (NC-4.1.1.1), Severe starvation related malnutrition (undernutrition) (NC-4.1.1.2), Mild non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.1), Moderate non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.2), Severe non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.3), Inadequate protein energy intake (NI-5.2), Inadequate fat intake (NI-5.5.1), Inadequate protein intake (NI-5.6.1), Intake of types of proteins inconsistent with needs (NI-5.6.3), Intake of types of amino acids inconsistent with needs (NI-5.7.2), Inadequate carbohydrate intake (NI-5.8.1), Inadequate vitamin intake (*specify*) (NI-5.9.1), Inadequate mineral intake (*specify*) (NI-5.10.1), Underweight (NC-3.1), Unintended weight loss (NC-3.2),), Growth rate below expected (NC-3.5), Poor nutrition quality of life (NB-2.5)
	• Fiber-containing foods	Inadequate fiber intake (NI-5.8.5)
	• Food that contains a bioactive substance	Inadequate bioactive substance intake (NI-4.1)
	• Healthy food choices	Excessive energy intake (NI-1.3), Excessive oral intake (NI-2.2), Excessive fat intake (NI-5.5.2), Intake of types of fats inconsistent with needs (*specify*) (NI-5.5.3)
	• Recommended foods	Undesirable food choices (NB-1.7)

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
	• Sufficient quantity or variety of culturally appropriate healthful food or water	Limited access to food (NB-3.2); Limited access to potable water (NB-3.4)
	• Safe and/or clear and accurately labeled food supply	Excessive bioactive substance intake (NI-4.2) Intake of unsafe food (NB-3.1)
	• Food storage equipment/facilities	Intake of unsafe food (NB-3.1)
	• Self management tools or decision guides or other nutrition related supplies	Inability to manage self care (NB-2.3), Limited access to nutrition related supplies (NB-3.3)
Access	Limited or absent, participation in community supplemental food programs such as food pantries, emergency kitchens, or shelters, with a sufficient variety of culturally appropriate healthful foods or nutrition related supplies	Limited access to food (NB-3.2), Limited access to nutrition related supplies (NB-3.3), Limited access to potable water (NB-3.4)
Access	Schools lacking nutrition/wellness policies or application of policies ensuring convenient, appetizing, competitively priced culturally appropriate healthful foods at meals, snacks, and school sponsored activities	Limited access to food (NB-3.2), Limited access to potable water (NB-3.4)
Behavior	Addictive behavior	Excessive physical activity (NB-2.2)
Behavior	Consumption of high-dose nutrient supplements	Imbalance of nutrients (NI-5.4)
Behavior	Eating behavior serves a purpose other than nourishment (eg, pica)	Unsupported beliefs/attitudes about food or nutrition related topics (NB-1.2), Limited food acceptance (NI-2.11)
Behavior	Excessive energy intake	Overweight, adult or pediatric (NC-3.3.1), Obese, pediatric (NC-3.3.2), Obese, Class I (NC-3.3.3), Obese, Class II (NC-3.3.4), Obese, Class III (NC-3.3.5),
Behavior	Excessive physical activity	Underweight (NC-3.1)
Behavior	Food and nutrition adherence limitations	Excessive carbohydrate intake (NI-5.8.2), Intake of types of carbohydrate inconsistent with needs (*specify*) (NI-5.8.3), Inconsistent carbohydrate intake (NI-5.8.4)
Behavior	Food or activity behavior-related difficulty	Poor nutrition quality of life (NB-2.5)
Behavior	Food preparation or eating patterns that involve only high-fiber foods to the exclusion of other nutrient-dense foods	Excessive fiber intake (NI-5.8.6)
Behavior	Disordered eating pattern	Excessive physical activity (NB-2.2), Underweight (NC-3.1), Overweight, adult or pediatric (NC-3.3.1), Obese, pediatric (NC-3.3.2), Obese, Class I (NC-3.3.3), Obese, Class II (NC-3.3.4), Obese, Class III (NC-3.3.5), Mild non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.1), Moderate non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.2), Severe non illness related pediatric malnutrition (undernutrition) (NC-4.1.4.3), Mild illness related pediatric malnutrition (undernutrition) (NC-4.1.5.), Moderate illness related

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.

Category	Etiology	Diagnosis
		pediatric malnutrition (undernutrition) (NC-4.1.5.2), Severe illness related pediatric malnutrition (undernutrition) (NC-4.1.5.3)
Behavior	Feeding via bottle or other route that may affect breastfeeding	Breastfeeding difficulty (NC-1.3), Predicted breastfeeding difficulty (NC-1.5)
Behavior	Frequent intake of foods containing bioactive substances	Excessive bioactive substance intake (NI-4.2)
Behavior	Limited food acceptance due to behavioral issues	Inadequate oral intake (NI-2.1), Growth rate below expected (NC-3.5)
Behavior	Inability to limit or refuse offered foods	Excessive oral intake (NI-2.2)
Behavior	Inadequate energy intake	Underweight (NC-3.1)
Behavior	Lack of or limited focus and attention to detail, difficulty with time management and/or organization	Self monitoring deficit (NB-1.4)
Behavior	Lifestyle change that reduces physical activity or activities of daily living	Physical inactivity (NB-2.1)
Behavior	Over consumption of a limited variety of foods	Excessive mineral intake (*specify*) (NI-5.10.2)
Behavior	Poor food planning, purchasing and preparation practices	Excessive oral intake (NI-2.2), Inadequate fiber intake (NI-5.8.5), Limited access to food (NB-3.2)
Behavior	Reluctance or avoidance of self feeding	Self feeding difficulty (NB-2.6)

Supplemental material for the electronic *Nutrition Care Process Terminology (eNCPT)*. Copyright 2020. Academy of Nutrition and Dietetics.