## Physicochemical, Microbiological and Sensory Properties of Drinking Yoghurt Incorporated with *Kithul (Caryota urens)* Flour

## Rasana S., Gunasekara D.C.S.1, Jayathilaka. S.N.T.1, Vidanarachchi J.K. and Prasanna P.H.P.\*

Department of Animal Science, Faculty of Agriculture, University of Peradeniya, Peradeniya, Sri Lanka

The current study was carried out to investigate physicochemical, microbiological, and sensory attributes of drinking-yoghurt prepared with varying concentrations of *Kithul* (*Caryota urens*) flour (0.1%, 0.2%, and 0.5%). The final products were stored at 4 °C for 21 days and physicochemical and microbiological properties were evaluated weekly during the storage. The results of sensory evaluation revealed that treatment containing 0.2% (v/v) *Kithul* flour had the highest preference. pH values of all yoghurt samples reduced significantly (p<0.05) during the storage period. Total coliform count, yeast and mould count were negative in all sample during 21 days of storage at 4 °C. Therefore, the results of the current study suggested that *Kithul* flour could be successfully utilized to manufacture drinking-yoghurt with enhanced physicochemical and sensory properties.

**Keywords:** Drinking yoghurt, *Kithul* flour, Microbiological properties, Physicochemical properties

Funding from the National Plant Quarantine Service of Sri Lanka is gratefully acknowledged

204

<sup>&</sup>lt;sup>1</sup>CIC Dairies (Pvt) Ltd., Ibbankatuwa, Dambulla, Sri Lanka

<sup>\*</sup>phpprasanna@agri.pdn.ac.lk