

Problem Set 1

Applied Stats/Quant Methods 1

Due: October 9, 2025

Instructions

- Please show your work! You may lose points by simply writing in the answer. If the problem requires you to execute commands in **R**, please include the code you used to get your answers. Please also include the **.R** file that contains your code. If you are not sure if work needs to be shown for a particular problem, please ask.
- Your homework should be submitted electronically on GitHub.
- This problem set is due before 23:59 on Thursday October 9, 2025. No late assignments will be accepted.

Question 1: Education

A school counselor was curious about the average of IQ of the students in her school and took a random sample of 25 students' IQ scores. The following is the data set:

```
1 y <- c(105, 69, 86, 100, 82, 111, 104, 110, 87, 108, 87, 90, 94, 113, 112, 98,  
      80, 97, 95, 111, 114, 89, 95, 126, 98)
```

1. Find a 90% confidence interval for the average student IQ in the school.

Finding a 90% confidence interval

Step 1: Calculate the sample mean. The sample mean equals the sum of the values divided by the number of values.

```
1 n <- length(y)
2 mean_y <- sum(y)/n
3
4 #or you can use the mean function
5 mean(y)
```

```
[1] 98.44
```

Step 2: Calculate the standard deviation

Find how much each data point differs from the mean and square each value.

```
1 distance_from_mean <- y - mean(y)
2
3 distance_from_mean_squared <- distance_from_mean^2
```

Add the squared values to find the sum of squares

```
1 sum_of_squares <- sum(distance_from_mean_squared)
```

Find the variance by dividing the sum of squared by the degrees of freedom (the number of values minus 1)

```
1 variance <- sum_of_squares/(n-1)
```

Find the standard deviation (the square root of variance)

```
1 standard_deviation_y <- sqrt(variance)
2 standard_deviation_y
```

```
[1] 13.09287
```

Step 3: Calculate the standard error, which equals the standard deviation divided by the square root of n(the number of observations in the sample)

```
1 standard_error_y <- standard_deviation_y/(sqrt(n))
2 standard_error_y
```

Step 4: Calculate the t-score by inputting (1 minus the confidence coefficient divided by 2) and the degrees of freedom (n-1)

```
1 t_score <- qt(p=(.1/2), n-1)
2 t_score
```

```
[1] -1.710882
```

Step 6: Calculate the lower limit of the confidence interval by doing the sample mean + (t-score * standard error)

```
1 lower_limit <- mean(y) + (t_score * standard_error_y)
2 lower_limit
```

```
[1] 93.95993
```

Step 7: Calculate the upper limit of the confidence interval by doing the sample mean - (t-score * standard error)

```
1 upper_limit <- mean(y) - (t_score * standard_error_y)
2 upper_limit
```

```
[1] 102.9201
```

Conclusion: Based on the confidence interval, we are 90 percent confident that the true mean of IQ at the school is between 93.95993 and 102.9201.

2. Next, the school counselor was curious whether the average student IQ in her school is higher than the average IQ score (100) among all the schools in the country.

Using the same sample, conduct the appropriate hypothesis test with $\alpha = 0.05$.

Conducting a hypothesis test

Step 1: Assumptions

- Due to the fact that n is less than 30, we cannot assume a normal distribution, so I will perform a t-test instead of a z-test
- Data is continuous
- The sample was random

Step 2: State Hypotheses

- Alternative Hypothesis: The average student IQ in the school counselor's school is higher than 100
- Null Hypothesis: The average student IQ in the school counselor's school is less than or equal to 100

Step 3: Calculate the test statistic (sample mean - population mean)/(sample SE)

```
1 test_statistic <- (mean(y) - 100)/(standard_error_y)
2 test_statistic
```

```
[1] -0.5957439
```

Step 4: Calculate the p-value by inputting the test statistic and the degrees of freedom (sample size - 1). lower.tail = FALSE was used because it is a one-sided hypothesis in which we are examining the upper tail of the curve.

```
1 pt(test_statistic, df=24, lower.tail=FALSE)
```

```
[1] -0.5957439
```

Conclusion: The calculated p value is greater than the critical value of 0.05. Therefore, we fail to reject the null hypothesis that the average student IQ in the school counselor's school is less than or equal to 100.

Question 2: Political Economy

Researchers are curious about what affects the amount of money communities spend on addressing homelessness. The following variables constitute our data set about social welfare expenditures in the USA.

State	<i>50 states in US</i>
Y	<i>per capita expenditure on shelters/housing assistance in state</i>
X1	<i>per capita personal income in state</i>
X2	<i>Number of residents per 100,000 that are "financially insecure" in state</i>
X3	<i>Number of people per thousand residing in urban areas in state</i>
Region	<i>1=Northeast, 2= North Central, 3= South, 4=West</i>


Explore the `expenditure` data set and import data into R.

- Please plot the relationships among Y , $X1$, $X2$, and $X3$? What are the correlations among them (you just need to describe the graph and the relationships among them)?
- Please plot the relationship between Y and $Region$? On average, which region has the highest per capita expenditure on housing assistance?
- Please plot the relationship between Y and $X1$? Describe this graph and the relationship. Reproduce the above graph including one more variable $Region$ and display different regions with different types of symbols and colors.

Plotting the relationships among Y, X1, X2, and X3

```
1 ggpairs(data=expenditure, columns = 2:5, title = "U.S. Social Welfare  
Expenditures")
```

Figure 1: Plot of Y, X1, X2, and X3




- Per capita personal income in state (X1) and per capita expenditure on shelters/housing assistance in state (Y) are positively correlated. The form is moderately spread out. A correlation coefficient of 0.532 indicates that it is a moderate positive correlation.
- Per capita expenditure on shelters/housing assistance in state (Y) is positively correlated with number of residents per 100,000 that are "financially insecure" in state (X2). The form is relatively tight. The form appears to curve indicating that a linear measurement may be the best way of analyzing this data. A correlation coefficient of 0.448 indicates a moderate positive linear correlation.
- Per capita personal income in state (X1) is positively correlated with the number of residents per 100,000 that are "financially insecure" in state (X2). The form is relatively spread out. A correlation coefficient of 0.221 suggests there is a weak correlation.

- Per capita expenditure on shelters/housing assistance in state (Y) is positively correlated with the number of people per thousand residing in urban areas in state (X3). The form is moderately tight. A correlation coefficient of 0.464 suggests a moderate positive correlation.
- Per capita personal income in state (X1) is positively correlated with the number of people per thousand residing in urban areas in state (X3). The distribution appears relatively tight. A correlation coefficient of 0.595 indicates a moderate positive association. This is the largest correlation coefficient and the tightest grouping that we see among variables in this data set.
- The number of residents per 100,000 that are "financially insecure" in state (X2) and the number of people per thousand residing in urban areas in state (X3) are positively correlated. The distribution appears very spread out. A correlation coefficient of 0.221 indicates a weak/negligible positive correlation.

Plotting the relationship between Y and Region

```
1 ggplot(expenditure) +  
2   geom_boxplot(aes(x = as.factor(expenditure$Region), y=expenditure$Y)) +  
3   labs(x="Region", y="Per capita expenditure on shelters/housing assistance in  
4     state (Y)",  
5   title = "Per Capita Expenditure on Shelters/Housing by Region") +  
6   theme(plot.title = element_text(hjust = 0.5)) +
```

Figure 2: Plot of Expenditure on Housing Assistance by Region
Per Capita Expenditure on Shelters/Housing by Region




On average, the West has the highest per capita expenditure on housing assistance. The North Central region has the next highest average expenditure, followed by the Northeast. Lastly, the South has the lowest average per capita expenditure on housing assistance.

Plotting the relationship between Y and X1

```
1 #Please plot the relationship between Y and X1
2 ggplot(expenditure, aes(x=X1, y=Y))+
3   geom_point() +
4   geom_text(label= show.legend = TRUE) +
5   labs(x="Per capita personal income in state (X1)",
```



Figure 4: Plot of Expenditure on Housing Assistance by Personal Income



This graph suggests that the positive linear correlation between per capita personal income in state and per capita expenditure on shelters/housing in state may have region as a confounding variable. Some regions tend to have data points that are clustered together. In particular, the South has a grouping of points that have both low per capita personal income in state and low expenditure on shelter/housing assistance. This example highlights the danger of assuming a causal relationship based on correlation.