

ROS_Normal Documentation

Date: 2025_10_23

Class:

Semester: Fall 2025

General

- Vital signs: 37.2 C (route unknown), P 71/min, RR 13/min, BP 111/72 mmHg (sitting, upper Rt extremity, manual), SpO2 99% RA, Ht 177.8 cm, Wt 90.7 kg
- General: NAD, A&O x3. well appeared. well groomed. well nourished. normal mood and affect. appears stated age. No acute distress

Skin

- Skin: warm, dry, intact, good turgor. no noted rashes or lesions. no abnormal bruising or pigmentation. appropriate color. no edema noted. no significant scarring. no bromhidrosis.
- Hair: normal texture and distribution. normal scalp integrity.
- Nails: normal color. no deformities. no pitting, clubbing, or cyanosis.

HEENT

Head

Head: normal head contour, atraumatic, normocephalic, facial symmetry. no edema, erythema, or masses. no tenderness of scalp, facial structures, joints, and temporal artery on palpation. skull integrity intact. hair texture normal, full even distribution of hair, no masses, facial drooping, fasciculations, thickening of skin, scaling/dryness, bony protuberance. B/L TMJ has no tenderness on palpation, crepitus, or edema. B/L parotid gland is not tender on palpation, firm, or enlarged. B/L sub-mandibular gland is not tender on palpation, firm, or enlarged.

focus PMHx always chronic, meds, allergies -> other we decide

no need to ask things we dont know like heart/pulmonary

top 3 and cant miss -rational pertinent +/- (cant miss can be in top 3 but rewrite on why dont miss focus)

topic resources have physical exam findings

Eyes

- External: BL eyebrow normal size, distribution, shape, extends to outer canthus. BL orbital area has no edema, lesions, masses, redundant skin. BL eyelids are symmetric, have no ptosis, edema, erythema, masses, lesions, scaling, infestations, or fasciculations. BL lacrimal gland unremarkable. Conjunctiva and Sclera pink, no injections, discharge, edema, pterygium, scleral icterus, or FB. Cornea clear. Iris intact. PERRLA, BL 3mm in size. No photo-phobia. EOM intact, no nystagmus. Visual Acuity: Gross visual acuity intact. Using the tumbling E chart, OD, OS, OU: 20/20, no corrective lens used. BL Visual fields are full via confrontation. BL corneal reflex intact. Palpation: no tenderness, masses, lesions, crepitus, edema noted on eyelids, globes, and orbital bones. Fundoscopic exam: BL red reflex. Unremarkable disc-cup ratio. Disc well-defined and cream colored. No haemorrhages, exudates, AV nicking. Macula intact and yellow-free.

Ears

- External: BL auricle aligned with outer cathus. Auricle, tragus, mastoid, lobule have no tenderness, erythema, edema, masses, or lesions. Internal: BL IAC has mild cerumen. BL IAC has no erythema, edema, discharge, FB, lesions, or masses. BL TM is translucent, pearly, intact, and convex. No effusion, retraction, scarring. BL TM is mobile with bony landmarks and cone of light visualized. Tests: Conversational hearing intact. BL unremarkable finger rub test. Weber lateralize BL. Rinne: AC>BC BL.

Nose

- External: External inspection normal and symmetric. Skin appropriate color, no masses, lesions, edema, ecchymoses, or rashes. No tenderness of bone, cartilage on palpation. Frontal sinus', maxillary sinus', sphenoid sinus' non-tender when palpated and percussed. Transillumination normal. BL nare patency intact. Internal: Muscosa pink, moist and septum mid-line. No discharge, polyps, congestion, or FB.

Throat

- External: Lips pink, intact, symmetric, well-defines. No edema, lesions, ecchymoses, or crust. Mouth: Normal bite. Good dentition, no missing teeth, no tenderness on palpation. Buccal mucosa pink and moist, no lesions. Gingiva pink and firm, no tenderness, lesions, or sores. Tongue is pink and mid-line, no edema or lesions. Parotid glad, submandibular gland, and sublingual glands are not tender, firm, or enlarged. Pharynx: Pharynx pink, moist, and clear. Soft palate pink and symmetric. Uvula midline with symmetric rise. BL Tonsils +1 without exudates. Gag reflex intact.

Neck

Neck: no skin dryness, masses, discoloration, lesions, scars, or edema. no webbing, fullness in the neck base, carotid prominence, jugular distension, overlying skin change or deformities. SCM and trapezius is symmetric. B/L Anterior and posterior cervical triangles present and unobstructed. No spinal deviation or tracheal deviation. No tenderness on palpation of cervical spine and para vertebral muscles. Swallow symmetry. Range of motion intact. Thyroid: is symmetric with normal consistency of the thyroid tissue and configuration. no tenderness on palpation, enlargement, nodules, or edema. Symmetric coordinated swallow. negative thyroid auscultation. Lymph Nodes: no overlying erythema, lymphangitic streaking, visual edema, visual enlargement. Nodes are soft, freely mobile, and discrete with no enlargement, warmth, and tenderness in the occipital, pre/post auricular, retro-pharyngeal, parotid, submandibular, submental, anterior/posterior cervical, supraclavicular, infraclavicular, epitrochlear, inguinal, femoral, popliteal lymph nodes.

Lymph Nodes

Lymph Nodes: no overlying erythema, lymphangitic streaking, visual edema, visual enlargement. Nodes are soft, freely mobile, and discrete with no enlargement, warmth, and tenderness in the occipital, pre/post auricular, retro-pharyngeal, parotid, submandibular, submental, anterior/posterior cervical, supraclavicular, infraclavicular, epitrochlear, inguinal, femoral, popliteal lymph nodes.

Endocrine

Cardiac

General: pt appears in NAD. (-) cyanosis, anxiety, diaphoresis, Levine Sign, labored breathing, SOB.

Skin: (-) nail clubbing, splinter hemorrhages, Osler's Nodes, Janeway Lesions. Hair Distributed normally.

Cardio: (-) Levine's Sign

no scars or chest wall deformities. PMI not visible, palpated at midclavicular line in 5th ICS. (-) lifts, heaves, thrills on inspection or palpation. (-) tenderness on palpation. Auscultation: RRR, S1 and S2 heard in all auscultation points, no rubs, clicks/gallops/snaps, murmur grade: 0. No audible S3 or S4.

Vascular: Inspection: (-) carotid pulsations, varicose veins. Auscultation: (-) carotid bruit.

Palpation: peripheral edema grade: 0. JVP 3cm above sternal angle @ 30' elevation. (-) JVD and hepatjugular reflex. capillary refill less than 3 sec. Carotid, radial, femoral, popliteal, PT, DP pulses present (+2) and equal B/L. (-) distal changes in color or temp.

Lung
