

Problem 1:

① Orthogonal Basis

We want $P(x)$ in P_2 such that $P(0) = P(1) = 0$.

$$P(x) = x(x-1),$$

$$P(0) = 0, P(1) = 0.$$

Basis Functions

$$P_1(x) = x(1-x),$$

We want P_2 orthogonal to P_1 .

number of basis functions.

+

Orthogonal Basis Functions

$$P(x) = \sum_{i=1}^n w_i x_i(x)$$

$$= w_1 x_1(x) + w_2 x_2(x) \\ = w_1 x(1-x) + w_2 x(1-x)^2$$

$$\frac{dP}{dx} = x^2 + x + 0$$

$$P(x) = x^2 + x^2 + 0 \times x + 0$$

$$P(0) = 0, P(1) = 0$$

$$P(1) = 1^2 + 1^2 + 0 = 2$$

$$P(1) = 2$$

percentage

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

$$P(x) = 1 - x^2$$

Handwritten notes in blue ink, possibly describing a process or algorithm.

1.

Ans. Part 1:
 $100 \times 100 = 10000$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 10000 & 0 & 0 \\ 0 & 10000 & 0 \\ 0 & 0 & 10000 \end{pmatrix} = \begin{pmatrix} 10000 & 0 & 0 \\ 0 & 10000 & 0 \\ 0 & 0 & 10000 \end{pmatrix}$$

$$= 10000 \times (1-x) + 10000 \times (1-x)^2$$

Ans. Part 2: $P(x) = 10000 \times (1-x) + 10000 \times (1-x)^2$

Calculation:

$$P(x) = 10000 \times (1-x) + 10000 \times (1-x)^2$$

$$= 10000 \times (1-x) + 10000 \times (1-x)^2$$

Ans. Part 3:
 $10000 \times (1-x) + 10000 \times (1-x)^2$

Ans. Part 4:

Problem 2:

1.

$$\frac{d}{dt}T + U^2 T - (U_1 + U_2)T = 0.$$

6.

Using the standard solver, we have
 Part A: $\frac{d}{dt}T + U^2 T - (U_1 + U_2)T = 0$ + the condition that T is a function of x only.
 Part B: $2T + U^2 T = U_1 T + U_2 T$ implies that:

$$T + U^2 T = U_1 T + U_2 T \Rightarrow T + (U_1 + U_2)T = 0 \Rightarrow T = 0$$

$$T + (U_1 + U_2)T = 0 \Rightarrow T = 0$$

$$T + U^2 T = U_1 T + U_2 T \Rightarrow T = 0$$

Part C: $\frac{d}{dt}T + U^2 T - (U_1 + U_2)T = 0$ + the condition that T is a function of x only.

$$\frac{d}{dt}T + U^2 T - (U_1 + U_2)T = 0 \Rightarrow T = 0$$

For B: $\frac{\partial \psi}{\partial t} = -i^2 \psi - 2i\alpha\psi \frac{\partial \psi}{\partial x} + \alpha^2 \psi = 0$

$\Rightarrow \frac{\partial \psi}{\partial t} = k^2 \psi_k + 2k\alpha\psi_k \psi_k + \alpha^2 \psi_k \psi_k = 0$

$\Rightarrow \frac{\partial \psi}{\partial t} = -i(k^2 + 2k\alpha\psi_k + \alpha^2 \psi_k) \psi_k = 0$

$\Rightarrow \psi_k(t) = e^{-i \int_0^t (k^2 + 2k\alpha\psi_k + \alpha^2 \psi_k) dt} \psi_k$

For i Riccati
Eq.
(Approximation)

For A: $\frac{\partial \psi}{\partial t} = 0 \cdot \psi = 0$

$\Rightarrow \partial_x \psi = -i \cdot 0 \cdot \psi \Rightarrow \psi(x) = e^{-i \cdot 0 \cdot x} \psi(0)$

Final solution (approximation):

$\psi_A(t, 0) = e^{-i \cdot 0 \cdot t} \psi^{-1} \left| \frac{-i(k^2 + 2k\alpha\psi_k + \alpha^2 \psi_k)t}{\psi} \right| e^{-i \cdot 0 \cdot t}$

Send that: ~~coll~~

$\psi(t) = e^{-i \cdot 0 \cdot t} \psi^{-1} \left| \frac{-i(k^2 + 2k\alpha\psi_k + \alpha^2 \psi_k)t}{\psi} \right| e^{-i \cdot 0 \cdot t} \psi(0) + O(t^3)$

Now, the time step:

$\Delta t < \frac{1}{\sqrt{x^2 + y^2 + z^2 + w^2}} \Rightarrow \Delta t < \frac{1}{\sqrt{x^2 + y^2 + z^2 + w^2}}$

Also, the force on ψ

$t = \frac{1}{\sqrt{x^2 + y^2 + z^2 + w^2}} + |a| \cdot x$

Not a strong
restriction or
before

At least:

$\Delta t < \frac{1}{\sqrt{x^2 + y^2 + z^2 + w^2}} + |a| \cdot x$

Step:
Since Δt is $\frac{1}{\sqrt{x^2 + y^2 + z^2 + w^2}}$,
the time step Δt is
small.

Step: Δt is small

good (approximation)

①. Now, we have a different problem: the workfunction will spread along distance.

due to the "prominent deformation of the electron".
 Instead, we can order our workfunction in a way such that we don't need as big of a simulation box. This will be done by the application of the translation operator $\hat{Q}(t)$.

(equivalent to changing the reference frame)

$$\hat{Q}(t) = \prod_{n=0}^{t-1} \left(\hat{Q}(t, n) \right) = \frac{1}{\hbar} \int \hat{Q}(t, n) \hat{Q}(t, n-1) \dots \hat{Q}(t, 0) \hat{Q}(t, -1) \dots$$

To verify that this corresponds to a translation, we can do:

$$\hat{Q}(\vec{r}, t) = \int d\vec{k} e^{-i\vec{k} \cdot \vec{r}} \phi_{\vec{k}}(\vec{r}, t)$$

Applying the operator: this explicit

$$\hat{Q}(t) \hat{Q}(\vec{r}, t) = \int d\vec{k} \hat{Q}(t) e^{-i\vec{k} \cdot \vec{r}} \phi_{\vec{k}}(\vec{r}, t)$$

$$\int d\vec{k} \left[\frac{e^{-i\vec{k} \cdot \vec{r}}}{e^{-i\vec{k} \cdot \vec{r}}} \cdot e^{-i\vec{k} \cdot \vec{r}} \right] \phi_{\vec{k}}(\vec{r}, t)$$

$$\int d\vec{k} e^{-i\vec{k} \cdot (\vec{r} + \alpha(t) \hat{x})} \phi_{\vec{k}}(\vec{r}, t)$$

$$\hat{Q}(\vec{r} + \alpha(t) \hat{x}, t) = \hat{Q}(\vec{r} + \alpha(t) \hat{x}, t)$$

translation is evident here.

In a variational approximation:

$$\hat{H}_0 = -\frac{\partial^2}{\partial x^2} = -\frac{\partial^2 (U_0 + U_1)}{\partial x^2}$$

$$= -\frac{\partial^2 U_0}{\partial x^2} = -\alpha \psi_1$$

we can neglect U_1 for any constant

$$\hat{H}_0 \psi_1 = -\alpha \psi_1 \quad \text{this time}$$

Introducing this in the total

$$\hat{H} \psi = \left(-\frac{\partial^2}{\partial x^2} + U_1 \right) \psi = 0$$

we can neglect U_1 for any constant

$$\hat{H}_0 \psi = 0 \Rightarrow$$

$$= -\frac{\partial^2 \psi}{\partial x^2} + U_1 \psi = 0 \Rightarrow$$

$$\Rightarrow x \partial_x \phi = -\partial^2 \phi - i \partial_x \phi + (i \partial_x + \partial_x) \phi \Rightarrow$$

$$\Rightarrow \partial_x \phi = -i(-\partial^2 - i \partial_x \phi - i \partial_x \phi) \quad \text{4.}$$

Ans 10:

$$\partial_x \phi = -i(-\partial^2 - i \partial_x \phi) \Rightarrow$$

$$\Rightarrow \partial_x \phi = -i(k^2 + k_x \partial_x) \phi \Rightarrow$$

$$\Rightarrow \phi_{k+x} = e^{-i(k^2 + k_x \partial_x) t}$$

Ans 11:

$$\partial_x \phi = -i(i \partial_x + \partial_x) \phi \Rightarrow \partial_x \phi = i \partial_x \phi + \partial_x \phi$$

$$\Rightarrow \partial_x \phi = i \partial_x \phi + \partial_x \phi \Rightarrow \partial_x \phi = i \partial_x \phi + \partial_x \phi$$

Ans 12:

$$\phi(t) = e^{-i(k^2 + k_x \partial_x) t} \Rightarrow \phi(t) = e^{-i(k^2 + k_x \partial_x) t}$$

Use Poisson's equation to solve with this
 for. Use in order to solve in order to
 solve. Solve in order to solve.

1. If don't want to solve the
 for the first part of the
 for the first part of the
 for the first part of the