

Fundamental limits for thermodynamic control with quantum feedback

Kaiyuan Ji (Cornell)

Joint work with Gilad Gour (Technion) and Mark M. Wilde (Cornell)



[arXiv:2503.09012](https://arxiv.org/abs/2503.09012)

Overview

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

Q: Does a system still obey the laws of thermodynamics when it is subject to feedback control?

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

Q: Does a system still obey the laws of thermodynamics when it is subject to feedback control?

A: **No.**

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)



Maxwell's demon

a classical feedback controller

- violates the second law

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)



Overview

Thermodynamics


- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)



Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)



Maxwell's demon

a classical feedback controller

- violates the second law
- limits on the violation [Parrondo et al. 2015]

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)



Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

- incoherent (classical) feedback



Maxwell's demon

a classical feedback controller

- violates the second law
- limits on the violation [Parrondo et al. 2015]

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)



Maxwell's demon

a classical feedback controller

- violates the second law
- limits on the violation [Parrondo et al. 2015]

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

- incoherent (classical) feedback



PHYSICAL REVIEW A, VOLUME 62, 022108

Coherent quantum feedback

Seth Lloyd*

d'Arbeloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)



Maxwell's demon

a classical feedback controller

- violates the second law
- limits on the violation [Parrondo et al. 2015]

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

- incoherent (classical) feedback
- **coherent (quantum) feedback**



PHYSICAL REVIEW A, VOLUME 62, 022108

Coherent quantum feedback

Seth Lloyd*

d'Arbeloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

- incoherent (classical) feedback
- **coherent (quantum) feedback**



Fully quantum Maxwell's demon
a controller with coherent feedback

PHYSICAL REVIEW A, VOLUME 62, 022108

Coherent quantum feedback

Seth Lloyd*

d'Arbeloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

- incoherent (classical) feedback
- **coherent (quantum) feedback**



Fully quantum Maxwell's demon
a controller with coherent feedback

- could violate the second law even more

PHYSICAL REVIEW A, VOLUME 62, 022108

Coherent quantum feedback

Seth Lloyd*

d'Arbeloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.

Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

- incoherent (classical) feedback
- **coherent (quantum) feedback**



Fully quantum Maxwell's demon a controller with coherent feedback

- could violate the second law even more
- limits on the violation **this work**

PHYSICAL REVIEW A, VOLUME 62, 022108

Coherent quantum feedback

Seth Lloyd*


d'Arbeloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.


Thermodynamics & feedback control

Second law: the average work consumed by a system is at least its change in (nonequilibrium) free energy




Thermodynamics & feedback control

Maxwell's demon, who holds side information, can violate the second law **locally**



Thermodynamics & feedback control


Maxwell's demon, who holds side information, can violate the second law **locally**



Thermodynamics & feedback control

Maxwell's demon, who holds side information, can violate the second law **locally**


- because of correlations between the system and the demon's memory



Thermodynamics & feedback control

Maxwell's demon, who holds side information, can violate the second law **locally**


- because of correlations between the system and the demon's memory
- the violation is limited: “second law with (classical) feedback” [Parrondo et al. 2015]



Thermodynamics & feedback control


Maxwell's demon, who holds side information, can violate the second law **locally**

- because of correlations between the system and the demon's memory
- the violation is limited: “second law with (classical) feedback” [Parrondo et al. 2015]



Thermodynamics & feedback control


Quantum generalizations have been studied [Sagawa & Ueda 2008, 2009]
[Narasimhachar & Gour 2017] [Minagawa et al. 2025]



Thermodynamics & feedback control

Quantum generalizations have been studied [Sagawa & Ueda 2008, 2009]
[Narasimhachar & Gour 2017] [Minagawa et al. 2025]


- ⚠ But side information is always collected by a measurement
- side information & feedback are incoherent (a classical demon)



Thermodynamics & feedback control

Quantum generalizations have been studied [Sagawa & Ueda 2008, 2009]
[Narasimhachar & Gour 2017] [Minagawa et al. 2025]

- ⚠ But side information is always collected by a measurement
- side information & feedback are incoherent (a classical demon)



PHYSICAL REVIEW A, VOLUME 62, 022108

Coherent quantum feedback

Seth Lloyd*

d'Arbelloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139


(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.

Thermodynamics & feedback control

Quantum generalizations have been studied [Sagawa & Ueda 2008, 2009]
[Narasimhachar & Gour 2017] [Minagawa et al. 2025]

- ⚠ But side information is always collected by a measurement
- side information & feedback are incoherent (a classical demon)
 - the existing “second law with (classical) feedback” is not universal: it could be violated by coherent (quantum) feedback



PHYSICAL REVIEW A, VOLUME 62, 022108

Coherent quantum feedback

Seth Lloyd*

d'Arbelloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.

Incoherent (classical) vs. coherent (quantum) feedback

PHYSICAL REVIEW A, VOLUME 62, 022108


Coherent quantum feedback

Seth Lloyd*

d'Arbelloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.



Incoherent (classical) vs. coherent (quantum) feedback

A **fully quantum demon** holds a quantum memory, and acquires, processes, and feeds back **quantum** side information coherently

PHYSICAL REVIEW A, VOLUME 62, 022108


Coherent quantum feedback

Seth Lloyd*

d'Arbelloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.



Incoherent (classical) vs. coherent (quantum) feedback

A **fully quantum demon** holds a quantum memory, and acquires, processes, and feeds back **quantum** side information coherently

- Some control tasks, e.g., state exchange and entanglement transfer, are achievable only with quantum feedback

PHYSICAL REVIEW A, VOLUME 62, 022108


Coherent quantum feedback

Seth Lloyd*

d'Arbelloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.



Incoherent (classical) vs. coherent (quantum) feedback

A **fully quantum demon** holds a quantum memory, and acquires, processes, and feeds back **quantum** side information coherently

- Some control tasks, e.g., state exchange and entanglement transfer, are achievable only with quantum feedback
- Second-law-like limits are universal only if they apply to quantum feedback

PHYSICAL REVIEW A, VOLUME 62, 022108


Coherent quantum feedback

Seth Lloyd*

d'Arbelloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.



Incoherent (classical) vs. coherent (quantum) feedback

A **fully quantum demon** holds a quantum memory, and acquires, processes, and feeds back **quantum** side information coherently

- Some control tasks, e.g., state exchange and entanglement transfer, are achievable only with quantum feedback
- Second-law-like limits are universal only if they apply to quantum feedback (**missing so far**)

PHYSICAL REVIEW A, VOLUME 62, 022108


Coherent quantum feedback

Seth Lloyd*

d'Arbelloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.



Overview

Thermodynamics

- Second law
 - convertibility between states
 - work cost / extractable work
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

- incoherent (classical) feedback
- **coherent (quantum) feedback**



Fully quantum Maxwell's demon a controller with coherent feedback

- could violate the second law even more
- limits on the violation **this work**

PHYSICAL REVIEW A, VOLUME 62, 022108

Coherent quantum feedback

Seth Lloyd*

d'Arbeloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.

Overview

Thermodynamics

- Second law ✓ with quantum feedback
 - convertibility between states ✓
 - work cost / extractable work ✓
- Resource theory (operational machinery)
 - macroscopic (asymptotic / average)
 - microscopic (single-shot)

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

- incoherent (classical) feedback
- **coherent (quantum) feedback**



Fully quantum Maxwell's demon a controller with coherent feedback

- could violate the second law even more
- limits on the violation **this work**

PHYSICAL REVIEW A, VOLUME 62, 022108

Coherent quantum feedback

Seth Lloyd*

d'Arbeloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.

Overview

Thermodynamics

- Second law ✓ with quantum feedback
 - convertibility between states ✓
 - work cost / extractable work ✓
- Resource theory (operational machinery) ✓
 - macroscopic (asymptotic / average) ✓
 - microscopic (single-shot) ✓

Feedback control

Interact with the system (e.g., measure it) to acquire side information

Control the system better based on the side information (feedback)

- incoherent (classical) feedback
- **coherent (quantum) feedback**



Fully quantum Maxwell's demon a controller with coherent feedback

- could violate the second law even more
- limits on the violation **this work**

PHYSICAL REVIEW A, VOLUME 62, 022108

Coherent quantum feedback

Seth Lloyd*

d'Arbeloff Laboratory, Mechanical Engineering, MIT 3-160, Cambridge, Massachusetts 02139

(Received 21 December 1998; published 14 July 2000)

In the conventional picture of quantum feedback control, sensors perform measurements on the system, a classical controller processes the results of the measurements, and actuators supply semiclassical potentials to alter the behavior of the quantum system. In this picture, the sensors tend to destroy coherence in the process of making measurements, and although the controller can use the actuators to act coherently on the quantum system, it is processing and feeding back classical information. This paper proposes an alternative method for quantum feedback control, in which the sensors, controller, and actuators are quantum systems that interact coherently with the system to be controlled. In this picture, the controller gets, processes, and feeds back *quantum* information. Controllers that operate using such quantum feedback loops can perform tasks such as entanglement transfer that are not possible using classical feedback. Necessary and sufficient conditions are presented for Hamiltonian quantum systems to be controllable and observable using both classical and quantum feedback.

Operational machinery

Resource theory of thermal nonequilibrium in the presence of quantum side information

Free operations

In thermodynamics, the resource is work **expendable on the system**

Free operations: feedback-control schemes that can be performed **without injecting work** into the system

Free operations

In thermodynamics, the resource is work **expendable on the system**

Free operations: feedback-control schemes that can be performed **without injecting work** into the system

But they are hard to delineate precisely (we still don't know how)

Axiomatic approach: identify properties that every such scheme must satisfy; regard every quantum channel with these properties as free

Free operations

In thermodynamics, the resource is work **expendable on the system**

Free operations: feedback-control schemes that can be performed **without injecting work** into the system

But they are hard to delineate precisely (we still don't know how)

Axiomatic approach: identify properties that every such scheme must satisfy; regard every quantum channel with these properties as free

- a larger set than the truly free operations;
but that is fine for no-go limits: if they hold for these permissive free operations, then they hold for the truly free operations as well

Free operations

In thermodynamics, the resource is work **expendable on the system**


Free operations: feedback-control schemes that can be performed **without injecting work** into the system

Free operations

In thermodynamics, the resource is work **expendable on the system**

Free operations: feedback-control schemes that can be performed **without injecting work** into the system

- ① One-way communication from memory to system



Free operations


In thermodynamics, the resource is work **expendable on the system**

Free operations: feedback-control schemes that can be performed **without injecting work** into the system

- ① One-way communication from memory to system

$$\mathcal{N}_{SM \rightarrow S'M'} = \mathcal{F}_{SL \rightarrow S'} \circ \mathcal{E}_{M \rightarrow LM'}$$

L is a quantum system



Free operations

In thermodynamics, the resource is work **expendable on the system**

Free operations: feedback-control schemes that can be performed **without injecting work** into the system

- ① One-way communication from memory to system

$$\mathcal{N}_{SM \rightarrow S'M'} = \mathcal{F}_{SL \rightarrow S'} \circ \mathcal{E}_{M \rightarrow LM'}$$


L is a quantum system

- ② Gibbs-preserving on the system if no side information is supplied (Kelvin's principle)

$$\mathcal{F}_{SL \rightarrow S'}[\gamma_S \otimes \rho_L] = \gamma'_{S'} \quad \forall \rho_L$$

Gibbs states $\gamma \propto e^{-\beta \hat{H}}$

feedback (L) only supplies information, not work



Free operations: QFGOs

In thermodynamics, the resource is work **expendable on the system**

Free operations: feedback-control schemes that can be performed **without injecting work** into the system

- ① One-way communication from memory to system

$$\mathcal{N}_{SM \rightarrow S'M'} = \mathcal{F}_{SL \rightarrow S'} \circ \mathcal{E}_{M \rightarrow LM'}$$


L is a quantum system

- ② Gibbs-preserving on the system if no side information is supplied (Kelvin's principle)

$$\mathcal{F}_{SL \rightarrow S'}[\gamma_S \otimes \rho_L] = \gamma'_{S'} \quad \forall \rho_L$$

Gibbs states $\gamma \propto e^{-\beta \hat{H}}$

feedback (L) only supplies information, not work



$\Rightarrow$ Quantum-feedback-assisted Gibbs-preserving operations (**QFGOs**)

Quantifying work

Pure qubits as the currency of work [Szilárd 1929]

- one pure qubit ($|0\rangle\langle 0|$, $\hat{H} = 0$) $\rightleftharpoons \beta^{-1} \ln 2$ of work
- one pure qudit of dimension d $\rightleftharpoons \beta^{-1} \ln d$ of work


Quantifying work

Pure qubits as the currency of work [Szilárd 1929]

- one pure qubit ($|0\rangle\langle 0|$, $\hat{H} = 0$) $\Leftrightarrow \beta^{-1} \ln 2$ of work
- one pure qudit of dimension d $\Leftrightarrow \beta^{-1} \ln d$ of work

Battery model [Faist, Dupuis, Oppenheim & Renner 2015]

- put the system alongside a battery S_0 , a pure qudit
- system & battery jointly undergo a QFGO (no external work supply)
- the battery ends up as a pure qudit S_1 of another dimension



Quantifying work


Pure qubits as the currency of work [Szilárd 1929]

- one pure qubit ($|0\rangle\langle 0|$, $\hat{H} = 0$) $\Leftrightarrow \beta^{-1} \ln 2$ of work
- one pure qudit of dimension d $\Leftrightarrow \beta^{-1} \ln d$ of work

Battery model [Faist, Dupuis, Oppenheim & Renner 2015]

- put the system alongside a battery S_0 , a pure qudit
- system & battery jointly undergo a QFGO (no external work supply)
- the battery ends up as a pure qudit S_1 of another dimension

$\Rightarrow$ work consumed by the system = decrease in battery charge = $\beta^{-1} (\ln |S_0| - \ln |S_1|)$ ($|S|$: dimension of S)



Definition of work cost

Task: transform an initial into a target (state of system & memory, Gibbs state of system)

$$(\rho_{SM}, \gamma_S) \longrightarrow (\rho'_{S'M'}, \gamma'_{S'})$$

↙ Gibbs state: encodes the Hamiltonian of S

Definition of work cost

Task: transform an initial into a target (state of system & memory, Gibbs state of system)


$$(\rho_{SM}, \gamma_S) \longrightarrow (\rho'_{S'M'}, \gamma'_{S'})$$

↖ Gibbs state: encodes the Hamiltonian of S

Work cost, allowing error ε and treating QFGOs as free:

$$W^\varepsilon((\rho_{SM}, \gamma_S) \rightarrow (\rho'_{S'M'}, \gamma'_{S'})) := \inf_{\substack{S_0, S_1, \\ \mathcal{N} \in \text{QFGO}}} \left\{ \beta^{-1} (\ln |S_0| - \ln |S_1|) : \mathcal{N}[|0\rangle\langle 0|_{S_0} \otimes \rho_{SM}] \approx_\varepsilon |0\rangle\langle 0|_{S_1} \otimes \rho'_{S'M'} \right\}$$

↖ in trace distance




Example: quantum feedback beats classical feedback

A system S correlated with a memory M ; all Hamiltonians trivial

- S : classical, 4 states $s \in \{0, 1, 2, 3\}$, uniformly mixed (thermal)
- M : one qubit holding the record $\hat{\sigma}_s |\psi\rangle$ ($\hat{\sigma}_0 = \mathbb{1}$; $\hat{\sigma}_{1,2,3}$: Pauli)
with $|\psi\rangle\langle\psi| = \frac{1}{2}\left(\hat{\sigma}_0 + \frac{1}{\sqrt{3}}(\hat{\sigma}_1 + \hat{\sigma}_2 + \hat{\sigma}_3)\right)$

S alone is thermal: no work is extractable without feedback



Bloch vectors of the 4 records:
vertices of a regular tetrahedron

Example: quantum feedback beats classical feedback

A system S correlated with a memory M ; all Hamiltonians trivial


- S : classical, 4 states $s \in \{0, 1, 2, 3\}$, uniformly mixed (thermal)
- M : one qubit holding the record $\hat{\sigma}_s |\psi\rangle$ ($\hat{\sigma}_0 = \mathbb{1}$; $\hat{\sigma}_{1,2,3}$: Pauli)
with $|\psi\rangle\langle\psi| = \frac{1}{2}(\hat{\sigma}_0 + \frac{1}{\sqrt{3}}(\hat{\sigma}_1 + \hat{\sigma}_2 + \hat{\sigma}_3))$

S alone is thermal: no work is extractable without feedback


Work extractable from S with feedback from M :

quantum feedback  $\beta^{-1} \ln 2$ (single shot)

scheme at right; a QFGO: with S thermal & independent of L , the controlled Pauli twirls L into $\mathbb{1}/2$, so it creates no work (Kelvin)



Bloch vectors of the 4 records:
vertices of a regular tetrahedron



Example: quantum feedback beats classical feedback

A system S correlated with a memory M ; all Hamiltonians trivial

- S : classical, 4 states $s \in \{0, 1, 2, 3\}$, uniformly mixed (thermal)
- M : one qubit holding the record $\hat{\sigma}_s |\psi\rangle$ ($\hat{\sigma}_0 = \mathbb{1}$; $\hat{\sigma}_{1,2,3}$: Pauli)
with $|\psi\rangle\langle\psi| = \frac{1}{2}(\hat{\sigma}_0 + \frac{1}{\sqrt{3}}(\hat{\sigma}_1 + \hat{\sigma}_2 + \hat{\sigma}_3))$

S alone is thermal: no work is extractable without feedback

Work extractable from S with feedback from M :

quantum feedback  $\beta^{-1} \ln 2$ (single shot)


scheme at right; a QFGO: with S thermal & independent of L , the controlled Pauli twirls L into $\mathbb{1}/2$, so it creates no work (Kelvin)

classical feedback  $\leq \beta^{-1} \ln \frac{4}{3}$ (even asymptotically)


measure $M \rightarrow X$; second law with classical feedback:

$$W_{\text{extr}} \leq \beta^{-1} I(S; X) \leq \beta^{-1} I_{\text{acc}} = \beta^{-1} \ln \frac{4}{3}$$

(I_{acc} : accessible information of the 4 records) [Holevo 1973]



Bloch vectors of the 4 records:
vertices of a regular tetrahedron



Main results

Single-shot & asymptotic work costs with quantum feedback

Task 1: state formation with quantum feedback

Starting from thermal equilibrium, create a target state (ρ_{SM}, γ_S) at minimum work cost

- Application: writing quantum data from an external memory into a device (e.g., a hard drive)
 - the device can end up entangled with the memory: impossible with classical feedback

Task 1: state formation with quantum feedback

Starting from thermal equilibrium, create a target state (ρ_{SM}, γ_S) at minimum work cost

- Application: writing quantum data from an external memory into a device (e.g., a hard drive)
 - the device can end up entangled with the memory: impossible with classical feedback



Task 1: state formation with quantum feedback

Starting from thermal equilibrium, create a target state (ρ_{SM}, γ_S) at minimum work cost

- Application: writing quantum data from an external memory into a device (e.g., a hard drive)
 - the device can end up entangled with the memory: impossible with classical feedback
- Prior work: classical feedback, classical data, asymptotic [Narasimhachar et al. 2019]



Task 1: state formation with quantum feedback

Starting from thermal equilibrium, create a target state (ρ_{SM}, γ_S) at minimum work cost

- Application: writing quantum data from an external memory into a device (e.g., a hard drive)
 - the device can end up entangled with the memory: impossible with classical feedback
- Prior work: classical feedback, classical data, asymptotic [Narasimhachar et al. 2019]



Definition: single-shot work of formation under QFGOs

$$W_{\text{form}}^{\varepsilon}(\rho_{SM}, \gamma_S) := W^{\varepsilon}(\emptyset \rightarrow (\rho_{SM}, \gamma_S))$$

no system & memory to start with (equivalently, S thermal)

Task 1: state formation with quantum feedback

Theorem 1. Single-shot & asymptotic work of formation under QFGOs:

$$W_{\text{form}}^{\varepsilon}(\rho_{SM}, \gamma_S) = \beta^{-1} I_{\max}^{\uparrow, \varepsilon}(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in [0, 1]$$
$$\lim_{n \rightarrow \infty} \frac{1}{n} W_{\text{form}}^{\varepsilon}(\rho_{SM}^{\otimes n}, \gamma_S^{\otimes n}) = \beta^{-1} I(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in (0, 1)$$

Task 1: state formation with quantum feedback

Theorem 1. Single-shot & asymptotic work of formation under QFGOs:

$$W_{\text{form}}^{\varepsilon}(\rho_{SM}, \gamma_S) = \beta^{-1} I_{\max}^{\uparrow, \varepsilon}(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in [0, 1]$$
$$\lim_{n \rightarrow \infty} \frac{1}{n} W_{\text{form}}^{\varepsilon}(\rho_{SM}^{\otimes n}, \gamma_S^{\otimes n}) = \beta^{-1} I(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in (0, 1)$$

Smoothed max-generalized mutual information (GMI): **SDP computable**


$$I_{\max}^{\uparrow, \varepsilon}(\rho_{SM} \| \gamma_S) := \inf_{t, \tau_{SM}, \sigma_M} \left\{ \ln t : \tau_{SM} \leq t \gamma_S \otimes \sigma_M, \tau_M \leq \sigma_M, \tau_{SM} \approx_{\varepsilon} \rho_{SM} \right\}$$

 τ_{SM} : subnormalized state; σ_M : state

Umegaki GMI:

$$I(\rho_{SM} \| \gamma_S) := D(\rho_{SM} \| \gamma_S \otimes \rho_M) = \underbrace{D(\rho_S \| \gamma_S)}_{\text{nonequilibrium of } S} + \underbrace{I(S; M)_{\rho}}_{\text{correlation (quantum mutual information)}}$$


Task 1: optimal protocol



Protocol

- $\mathcal{E}$: prepare $|M| \omega_M^{1/2} \Phi_{LM} \omega_M^{1/2}$ (Φ : maximally entangled) and feed $L \cong M$ back
- $\mathcal{F}$: measure the battery S_0 ; outcome 0 $\Rightarrow$ apply $\mathcal{F}_{L \rightarrow S_1 S}^{(0)}$ (output $\omega_{S_1 S M}$); otherwise $\Rightarrow$ apply $\mathcal{F}_{L \rightarrow S_1 S}^\perp$

Task 1: optimal protocol




Protocol

- $\mathcal{E}$: prepare $|M| \omega_M^{1/2} \Phi_{LM} \omega_M^{1/2}$ (Φ : maximally entangled) and feed $L \cong M$ back
- $\mathcal{F}$: measure the battery S_0 ; outcome 0 $\Rightarrow$ apply $\mathcal{F}_{L \rightarrow S_1 S}^{(0)}$ (output $\omega_{S_1 S M}$); otherwise $\Rightarrow$ apply $\mathcal{F}_{L \rightarrow S_1 S}^\perp$

Why it works

- Kelvin: a thermal battery gives outcome 0 only w.p. $1/|S_0|$; $\mathcal{F}^\perp$ makes the average output $\frac{\mathbb{1}_{S_1}}{|S_1|} \otimes \gamma_S$ for any L , so the scheme is a QFGO

Task 1: optimal protocol



Protocol

- $\mathcal{E}$: prepare $|M| \omega_M^{1/2} \Phi_{LM} \omega_M^{1/2}$ (Φ : maximally entangled) and feed $L \cong M$ back
- $\mathcal{F}$: measure the battery S_0 ; outcome 0 $\Rightarrow$ apply $\mathcal{F}_{L \rightarrow S_1 S}^{(0)}$ (output $\omega_{S_1 S M}$); otherwise $\Rightarrow$ apply $\mathcal{F}_{L \rightarrow S_1 S}^\perp$

Why it works

- Kelvin: a thermal battery gives outcome 0 only w.p. $1/|S_0|$; $\mathcal{F}^\perp$ makes the average output $\frac{\mathbb{1}_{S_1}}{|S_1|} \otimes \gamma_S$ for any L , so the scheme is a QFGO
- $\mathcal{F}^\perp$ is CPTP as long as $|S_0| \geq e^{D_{\max}(\omega_{S_1 S M} \| \frac{\mathbb{1}_{S_1}}{|S_1|} \otimes \gamma_S \otimes \omega_M)}$. So, for a suitable S_1 and $\omega_{S_1 S M}$, the minimum cost is

$$\beta^{-1}(\ln |S_0| - \ln |S_1|) \approx \beta^{-1}(D_{\max}(\omega_{S_1 S M} \| \frac{\mathbb{1}_{S_1}}{|S_1|} \otimes \gamma_S \otimes \omega_M) - \ln |S_1|) = \beta^{-1} I_{\max}^{\uparrow, \epsilon}(\rho_{SM} \| \gamma_S)$$

Optimal protocol for state formation

Alternative expression of the smoothed max-GMI:

$$I_{\max}^{\uparrow, \varepsilon}(\rho_{SM} \| \gamma_S) = \inf_{S_1, \omega_{S_1 SM}} \left\{ D_{\max}(\omega_{S_1 SM} \| \frac{\mathbb{1}_{S_1}}{|S_1|} \otimes \gamma_S \otimes \omega_M) - \ln |S_1| : \omega_{S_1 SM} \approx_{\varepsilon} |0\rangle\langle 0|_{S_1} \otimes \rho_{SM} \right\}$$

- (i) prepare $|M| \omega_M^{1/2} \Phi_{LM} \omega_M^{1/2}$ with $L \cong M$ and $\Phi_{LM} \equiv \frac{1}{|M|} \sum_{i,j} |i\rangle\langle j|_L \otimes |i\rangle\langle j|_M$; (ii) battery $|S_0| = \lceil e^{D_{\max}(\omega_{S_1 SM} \| \frac{\mathbb{1}_{S_1}}{|S_1|} \otimes \gamma_S \otimes \omega_M)} \rceil$
 (iii) measure S_0 : outcome 0 $\Rightarrow$ apply $\mathcal{F}_{L \rightarrow S_1 S}^{(0)}$; otherwise $\Rightarrow$ apply $\mathcal{F}_{L \rightarrow S_1 S}^{\perp}$, where $(\omega_M$ full rank without loss of generality)

$$\mathcal{F}_{L \rightarrow S_1 S}^{(0)}[\cdot] := \text{Tr}_L[\omega_L^{-\frac{1}{2}} \omega_{S_1 S L} \omega_L^{-\frac{1}{2}} (\cdot)_L^{\top}], \quad \mathcal{F}_{L \rightarrow S_1 S}^{\perp}[\cdot] := \frac{1}{|S_0|-1} \left(|S_0| \frac{\mathbb{1}_{S_1}}{|S_1|} \otimes \gamma_S \otimes \text{Tr}_L[\cdot] - \mathcal{F}_{L \rightarrow S_1 S}^{(0)}[\cdot] \right)$$

Channel: $\mathcal{N}_{S_0 \rightarrow S_1 SM} = \mathcal{F}_{S_0 L \rightarrow S_1 S} \circ \mathcal{E}_{\emptyset \rightarrow LM}$, $\mathcal{E}_{\emptyset \rightarrow LM} := |M| \omega_M^{1/2} \Phi_{LM} \omega_M^{1/2}$,

$$\mathcal{F}_{S_0 L \rightarrow S_1 S}[\cdot] = (\text{Tr}_{S_0} \otimes \mathcal{F}_{L \rightarrow S_1 S}^{(0)})[|0\rangle\langle 0|_{S_0} (\cdot)_{S_0 L}] + (\text{Tr}_{S_0} \otimes \mathcal{F}_{L \rightarrow S_1 S}^{\perp})[(\mathbb{1}_{S_0} - |0\rangle\langle 0|_{S_0}) (\cdot)_{S_0 L}]$$

$\mathcal{F}^{(0)}$ is a channel (Choi operator $\omega_L^{-\frac{1}{2}} \omega_{S_1 S L} \omega_L^{-\frac{1}{2}}$); $\mathcal{F}^{\perp}$ is CPTP iff $|S_0| \frac{\mathbb{1}_{S_1}}{|S_1|} \otimes \gamma_S \otimes \omega_M \geq \omega_{S_1 SM}$, i.e., iff $|S_0| \geq e^{D_{\max}(\dots)}$

Kelvin: $\mathcal{F}_{S_0 L \rightarrow S_1 S}[\frac{\mathbb{1}_{S_0}}{|S_0|} \otimes (\cdot)_L] = \frac{\mathbb{1}_{S_1}}{|S_1|} \otimes \gamma_S \otimes \text{Tr}_L[\cdot]$ Output: $\mathcal{N}[|0\rangle\langle 0|_{S_0}] = \omega_{S_1 SM} \approx_{\varepsilon} |0\rangle\langle 0|_{S_1} \otimes \rho_{SM}$

Cost: $\beta^{-1}(\ln |S_0| - \ln |S_1|) \approx \beta^{-1}(D_{\max}(\dots) - \ln |S_1|)$, whose infimum over S_1 and $\omega_{S_1 SM}$ is $\beta^{-1} I_{\max}^{\uparrow, \varepsilon}(\rho_{SM} \| \gamma_S)$

Task 2: work extraction with quantum feedback

Starting from (ρ_{SM}, γ_S) , extract as much work as possible

- Prior work: classical feedback, asymptotic [Morris, Lami & Adesso 2019]

Task 2: work extraction with quantum feedback

Starting from (ρ_{SM}, γ_S) , extract as much work as possible

- Prior work: classical feedback, asymptotic [Morris, Lami & Adesso 2019]
- Quantum feedback can extract strictly more
 - asymptotically, the gap is the quantum discord [Ollivier & Zurek 2001] [Manzano et al. 2018]

Task 2: work extraction with quantum feedback

Starting from (ρ_{SM}, γ_S) , extract as much work as possible

- Prior work: classical feedback, asymptotic [Morris, Lami & Adesso 2019]
- Quantum feedback can extract strictly more
 - asymptotically, the gap is the quantum discord [Ollivier & Zurek 2001] [Manzano et al. 2018]
 - e.g., the example: $\beta^{-1} \ln 2$ with quantum feedback vs. $\leq \beta^{-1} \ln \frac{4}{3}$ with classical feedback

Task 2: work extraction with quantum feedback

Starting from (ρ_{SM}, γ_S) , extract as much work as possible

- Prior work: classical feedback, asymptotic [Morris, Lami & Adesso 2019]
- Quantum feedback can extract strictly more
 - asymptotically, the gap is the quantum discord [Ollivier & Zurek 2001] [Manzano et al. 2018]
 - e.g., the example: $\beta^{-1} \ln 2$ with quantum feedback vs. $\leq \beta^{-1} \ln \frac{4}{3}$ with classical feedback

Definition: single-shot extractable work under QFGOs

$$W_{\text{extr}}^{\varepsilon}(\rho_{SM}, \gamma_S) := -W^{\varepsilon}((\rho_{SM}, \gamma_S) \rightarrow \emptyset)$$

 the initial state is used up

Task 2: work extraction with quantum feedback

Theorem 2. Single-shot & asymptotic extractable work under QFGOs:

$$W_{\text{extr}}^{\varepsilon}(\rho_{SM}, \gamma_S) = \beta^{-1} I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in [0, 1]$$
$$\lim_{n \rightarrow \infty} \frac{1}{n} W_{\text{extr}}^{\varepsilon}(\rho_{SM}^{\otimes n}, \gamma_S^{\otimes n}) = \beta^{-1} I(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in (0, 1)$$

Task 2: work extraction with quantum feedback

Theorem 2. Single-shot & asymptotic extractable work under QFGOs:

$$W_{\text{extr}}^{\varepsilon}(\rho_{SM}, \gamma_S) = \beta^{-1} I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \parallel \gamma_S) \quad \forall \varepsilon \in [0, 1]$$
$$\lim_{n \rightarrow \infty} \frac{1}{n} W_{\text{extr}}^{\varepsilon}(\rho_{SM}^{\otimes n}, \gamma_S^{\otimes n}) = \beta^{-1} I(\rho_{SM} \parallel \gamma_S) \quad \forall \varepsilon \in (0, 1)$$

the example: $\beta^{-1} I(\rho_{SM} \parallel \frac{1_S}{4}) = \beta^{-1} \ln 2$
 $\Rightarrow$ its quantum scheme is optimal

Task 2: work extraction with quantum feedback

Theorem 2. Single-shot & asymptotic extractable work under QFGOs:

$$W_{\text{extr}}^\varepsilon(\rho_{SM}, \gamma_S) = \beta^{-1} I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in [0, 1]$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} W_{\text{extr}}^\varepsilon(\rho_{SM}^{\otimes n}, \gamma_S^{\otimes n}) = \beta^{-1} I(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in (0, 1)$$

the example: $\beta^{-1} I(\rho_{SM} \| \frac{1_S}{4}) = \beta^{-1} \ln 2$
 $\Rightarrow$ its quantum scheme is optimal

Smoothed min-GMI: **SDP computable**

$$I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \| \gamma_S) := \sup_{\Lambda_{SM}} \left\{ -\ln \|\text{Tr}_S[\Lambda_{SM} \gamma_S]\|_\infty : \text{Tr}[\Lambda_{SM} \rho_{SM}] \geq 1 - \varepsilon, \ 0 \leq \Lambda_{SM} \leq \mathbb{1}_{SM} \right\}$$

$\uparrow$
a test: hypothesis testing between ρ_{SM} and $\{\gamma_S \otimes \sigma_M : \sigma_M \text{ a state}\}$

Task 2: work extraction with quantum feedback

Theorem 2. Single-shot & asymptotic extractable work under QFGOs:

$$W_{\text{extr}}^\varepsilon(\rho_{SM}, \gamma_S) = \beta^{-1} I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in [0, 1]$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} W_{\text{extr}}^\varepsilon(\rho_{SM}^{\otimes n}, \gamma_S^{\otimes n}) = \beta^{-1} I(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in (0, 1)$$

the example: $\beta^{-1} I(\rho_{SM} \| \frac{\mathbb{1}_S}{4}) = \beta^{-1} \ln 2$
 $\Rightarrow$ its quantum scheme is optimal

Smoothed min-GMI: **SDP computable**

$$I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \| \gamma_S) := \sup_{\Lambda_{SM}} \left\{ -\ln \|\text{Tr}_S[\Lambda_{SM} \gamma_S]\|_\infty : \text{Tr}[\Lambda_{SM} \rho_{SM}] \geq 1 - \varepsilon, 0 \leq \Lambda_{SM} \leq \mathbb{1}_{SM} \right\}$$

$\uparrow$ a test: hypothesis testing between ρ_{SM} and $\{\gamma_S \otimes \sigma_M : \sigma_M \text{ a state}\}$

Erasing S ($\hat{H}_S = 0$), i.e., resetting it to $|0\rangle\langle 0|_S$, with quantum feedback: a single-shot Landauer principle

$$W_{\text{erase}}^\varepsilon = \beta^{-1} \ln |S| - W_{\text{extr}}^\varepsilon(\rho_{SM}, \frac{\mathbb{1}_S}{|S|}) = \beta^{-1} H_{\max}^{\uparrow, \varepsilon}(S|M)_\rho$$

Task 2: work extraction with quantum feedback

Theorem 2. Single-shot & asymptotic extractable work under QFGOs:

$$W_{\text{extr}}^\varepsilon(\rho_{SM}, \gamma_S) = \beta^{-1} I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in [0, 1]$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} W_{\text{extr}}^\varepsilon(\rho_{SM}^{\otimes n}, \gamma_S^{\otimes n}) = \beta^{-1} I(\rho_{SM} \| \gamma_S) \quad \forall \varepsilon \in (0, 1)$$

the example: $\beta^{-1} I(\rho_{SM} \| \frac{1_S}{4}) = \beta^{-1} \ln 2$
 $\Rightarrow$ its quantum scheme is optimal

Smoothed min-GMI: **SDP computable**

$$I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \| \gamma_S) := \sup_{\Lambda_{SM}} \left\{ -\ln \|\text{Tr}_S[\Lambda_{SM} \gamma_S]\|_\infty : \text{Tr}[\Lambda_{SM} \rho_{SM}] \geq 1 - \varepsilon, 0 \leq \Lambda_{SM} \leq \mathbb{1}_{SM} \right\}$$

$\uparrow$ a test: hypothesis testing between ρ_{SM} and $\{\gamma_S \otimes \sigma_M : \sigma_M \text{ a state}\}$


Erasing S ($\hat{H}_S = 0$), i.e., resetting it to $|0\rangle\langle 0|_S$, with quantum feedback: a single-shot Landauer principle

$$W_{\text{erase}}^\varepsilon = \beta^{-1} \ln |S| - W_{\text{extr}}^\varepsilon(\rho_{SM}, \frac{1_S}{|S|}) = \beta^{-1} H_{\max}^{\uparrow, \varepsilon}(S|M)_\rho$$

can be negative: cooling rather than heating

cf. [del Rio et al. 2011]


Task 2: optimal protocol



Protocol

- $\mathcal{E}$: feed the memory back unchanged ($M \rightarrow L$)
- $\mathcal{F}$: test S_0SL with Λ_{SL} (accepts ρ_{SM} w.p. $1 - \varepsilon$); accept $\Rightarrow |0\rangle\langle 0|_{S_1}$, reject $\Rightarrow \frac{\mathbb{1}_{S_1} - |0\rangle\langle 0|_{S_1}}{|S_1| - 1}$

Task 2: optimal protocol



Protocol

- $\mathcal{E}$: feed the memory back unchanged ($M \rightarrow L$)
- $\mathcal{F}$: test S_0SL with Λ_{SL} (accepts ρ_{SM} w.p. $1 - \epsilon$); accept $\Rightarrow |0\rangle\langle 0|_{S_1}$, reject $\Rightarrow \frac{\mathbb{1}_{S_1} - |0\rangle\langle 0|_{S_1}}{|S_1| - 1}$

Why it works

- Kelvin: if S is thermal, it accepts w.p. $\leq \|\text{Tr}_S[\Lambda_{SL}\gamma_S]\|_{\infty}$ for any L ; a QFGO if $\frac{1}{|S_1|} \approx \frac{1}{|S_0|} \|\text{Tr}_S[\Lambda_{SL}\gamma_S]\|_{\infty}$
- work extracted: $\beta^{-1} (\ln |S_1| - \ln |S_0|) \approx -\beta^{-1} \ln \|\text{Tr}_S[\Lambda_{SM}\gamma_S]\|_{\infty} = \beta^{-1} I_{\min}^{\downarrow, \epsilon}(\rho_{SM} \parallel \gamma_S)$

Extracting work = telling ρ_{SM} apart from “thermal S , any feedback”

Optimal protocol for work extraction

(i) optimal test Λ_{SM} for $I_{\min}^{\downarrow, \varepsilon}$; (ii) $\mathcal{E}_{M \rightarrow L}[\cdot] = (\cdot)_{M \rightarrow L}$; (iii) measure S_0SL with $(\Omega_{S_0SL}, \mathbb{1} - \Omega_{S_0SL})$,

$$\Omega_{S_0SL} := |0\rangle\langle 0|_{S_0} \otimes \Lambda_{SL} + |1\rangle\langle 1|_{S_0} \otimes \mathbb{1}_S \otimes \left(\frac{|S_0|}{|S_1|} \mathbb{1}_L - \text{Tr}_S[\Lambda_{SL} \gamma_S] \right), \quad |S_1| = \left\lfloor \frac{|S_0|}{\|\text{Tr}_S[\Lambda_{SM} \gamma_S]\|_{\infty}} \right\rfloor$$

so that $0 \leq \Omega_{S_0SL} \leq \mathbb{1}$; accept $\Rightarrow S_1$ in $|0\rangle\langle 0|_{S_1}$, reject $\Rightarrow S_1$ in $(\mathbb{1}_{S_1} - |0\rangle\langle 0|_{S_1})/(|S_1| - 1)$:

$$\mathcal{F}_{S_0SL \rightarrow S_1}[\cdot] = \text{Tr}[\Omega_{S_0SL}(\cdot)] |0\rangle\langle 0|_{S_1} + \text{Tr}[(\mathbb{1} - \Omega_{S_0SL})(\cdot)] \frac{\mathbb{1}_{S_1} - |0\rangle\langle 0|_{S_1}}{|S_1| - 1}$$

Kelvin: $\mathcal{F}_{S_0SL \rightarrow S_1} \left[\frac{\mathbb{1}_{S_0}}{|S_0|} \otimes \gamma_S \otimes (\cdot)_L \right] = \frac{\mathbb{1}_{S_1}}{|S_1|} \otimes \text{Tr}_L[\cdot]$

Output: $\mathcal{N}[|0\rangle\langle 0|_{S_0} \otimes \rho_{SM}] = \text{Tr}[\Lambda \rho] |0\rangle\langle 0|_{S_1} + (1 - \text{Tr}[\Lambda \rho]) \frac{\mathbb{1}_{S_1} - |0\rangle\langle 0|_{S_1}}{|S_1| - 1} \approx_{\varepsilon} |0\rangle\langle 0|_{S_1}$

Work: $\beta^{-1} (\ln |S_1| - \ln |S_0|) \approx -\beta^{-1} \ln \|\text{Tr}_S[\Lambda_{SM} \gamma_S]\|_{\infty} = \beta^{-1} I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \| \gamma_S)$

Generalized second law with quantum feedback

Theorem 3. Asymptotic work cost under QFGOs, $\forall \varepsilon \in (0, 1)$:

$$\lim_{n \rightarrow \infty} \frac{1}{n} W^\varepsilon((\rho_{SM}^{\otimes n}, \gamma_S^{\otimes n}) \rightarrow (\rho_{S'M'}^{\otimes n}, \gamma_{S'}^{\otimes n})) = (F(S'|M')_{\rho'} - F(S')_{\gamma'}) - (F(S|M)_\rho - F(S)_\gamma)$$

Conditional nonequilibrium free energy: [Bera et al. 2017]

$$F(S|M)_\rho := \text{Tr}[\hat{H}_S \rho_S] - \beta^{-1} H(S|M)_\rho = F(S)_\rho + \beta^{-1} I(S; M)_\rho$$

$H(S|M)$: conditional entropy, which can be negative (entanglement); $F(S)_\gamma$: equilibrium free energy

Generalized second law with quantum feedback

Theorem 3. Asymptotic work cost under QFGOs, $\forall \varepsilon \in (0, 1)$:

$$\lim_{n \rightarrow \infty} \frac{1}{n} W^\varepsilon((\rho_{SM}^{\otimes n}, \gamma_S^{\otimes n}) \rightarrow (\rho_{S'M'}^{\otimes n}, \gamma_{S'}^{\otimes n})) = (F(S'|M')_{\rho'} - F(S')_{\gamma'}) - (F(S|M)_\rho - F(S)_\gamma)$$

Conditional nonequilibrium free energy: [Bera et al. 2017]

$$F(S|M)_\rho := \text{Tr}[\hat{H}_S \rho_S] - \beta^{-1} H(S|M)_\rho = F(S)_\rho + \beta^{-1} I(S; M)_\rho$$

$H(S|M)$: conditional entropy, which can be negative (entanglement); $F(S)_\gamma$: equilibrium free energy

QFGOs include every physical feedback-control scheme, so the average work consumed by the system obeys

$$\langle W \rangle \geq \Delta F(S|M) = \Delta F_{\text{sys}} + \beta^{-1} \Delta I_q \quad (\text{Theorem 3 + change in equilibrium free energy})$$

Generalized second law with quantum feedback

Theorem 3. Asymptotic work cost under QFGOs, $\forall \varepsilon \in (0, 1)$:

$$\lim_{n \rightarrow \infty} \frac{1}{n} W^\varepsilon((\rho_{SM}^{\otimes n}, \gamma_S^{\otimes n}) \rightarrow (\rho_{S'M'}^{\otimes n}, \gamma_{S'}^{\otimes n})) = (F(S'|M')_{\rho'} - F(S')_{\gamma'}) - (F(S|M)_\rho - F(S)_\gamma)$$

Conditional nonequilibrium free energy: [Bera et al. 2017]

$$F(S|M)_\rho := \text{Tr}[\hat{H}_S \rho_S] - \beta^{-1} H(S|M)_\rho = F(S)_\rho + \beta^{-1} I(S; M)_\rho$$

$H(S|M)$: conditional entropy, which can be negative (entanglement); $F(S)_\gamma$: equilibrium free energy

QFGOs include every physical feedback-control scheme, so the average work consumed by the system obeys


$$\langle W \rangle \geq \Delta F(S|M) = \Delta F_{\text{sys}} + \beta^{-1} \Delta I_q \quad (\text{Theorem 3 + change in equilibrium free energy})$$

↑
change in **quantum** mutual information (classical before)

Consistency with the traditional second law

Maxwell's demon violates the second law only **locally**

- no global violation over a closed cycle: the work gained through feedback is paid back elsewhere (measurement, reset) [Bennett 1982] [Sagawa & Ueda 2009] [Minagawa et al. 2025]




Consistency with the traditional second law

Maxwell's demon violates the second law only **locally**

- no global violation over a closed cycle: the work gained through feedback is paid back elsewhere (measurement, reset) [Bennett 1982] [Sagawa & Ueda 2009] [Minagawa et al. 2025]

? Does this survive for a fully quantum demon?




Consistency with the traditional second law

Maxwell's demon violates the second law only **locally**


- no global violation over a closed cycle: the work gained through feedback is paid back elsewhere (measurement, reset) [Bennett 1982] [Sagawa & Ueda 2009] [Minagawa et al. 2025]

❓ Does this survive for a fully quantum demon?

- most general cycle: quantum memory throughout; acquisition by any interaction (not a measurement); quantum processing & feedback



Consistency with the traditional second law




Consistency with the traditional second law

From the generalized second law alone:

$$\langle W_{\text{sys}} \rangle \geq \Delta F_{\text{sys}} + \beta^{-1} \Delta I_q$$

generalized second law with quantum feedback



Consistency with the traditional second law


From the generalized second law alone:

$$\langle W_{\text{sys}} \rangle \geq \Delta F_{\text{sys}} + \beta^{-1} \Delta I_q$$

generalized second law with quantum feedback

$$\langle W_{\text{mem}} \rangle \geq \Delta F_{\text{mem}}$$

the memory itself receives no feedback



Consistency with the traditional second law

From the generalized second law alone:

$$\langle W_{\text{sys}} \rangle \geq \Delta F_{\text{sys}} + \beta^{-1} \Delta I_{\text{q}}$$


generalized second law with quantum feedback

$$\langle W_{\text{mem}} \rangle \geq \Delta F_{\text{mem}}$$

the memory itself receives no feedback

$$\langle W_{\text{acqn}} \rangle + \langle W_{\text{rst}} \rangle \geq -\Delta F_{\text{sys\&mem}} = -(\Delta F_{\text{sys}} + \Delta F_{\text{mem}} + \beta^{-1} \Delta I_{\text{q}})$$

no feedback here; the cycle closes



Consistency with the traditional second law

From the generalized second law alone:

$$\langle W_{\text{sys}} \rangle \geq \Delta F_{\text{sys}} + \beta^{-1} \Delta I_{\text{q}}$$

generalized second law with quantum feedback

$$\langle W_{\text{mem}} \rangle \geq \Delta F_{\text{mem}}$$


the memory itself receives no feedback

$$\langle W_{\text{acqn}} \rangle + \langle W_{\text{rst}} \rangle \geq -\Delta F_{\text{sys\&mem}} = -(\Delta F_{\text{sys}} + \Delta F_{\text{mem}} + \beta^{-1} \Delta I_{\text{q}})$$

no feedback here; the cycle closes

$$\Rightarrow \langle W_{\text{sys}} \rangle + \langle W_{\text{mem}} \rangle + \langle W_{\text{acqn}} \rangle + \langle W_{\text{rst}} \rangle \geq 0$$

as the traditional second law predicts



Conclusion

Fundamental limits on thermodynamic control with coherent (quantum) feedback, on both microscopic & macroscopic scales

- QFGOs: axiomatic and deliberately permissive $\Rightarrow$ the limits hold for every physical feedback-control scheme

Conclusion

Fundamental limits on thermodynamic control with coherent (quantum) feedback, on both microscopic & macroscopic scales

- QFGOs: axiomatic and deliberately permissive $\Rightarrow$ the limits hold for every physical feedback-control scheme
- Single shot: exact work costs, with optimal protocols
 - formation $\beta^{-1} I_{\max}^{\uparrow, \varepsilon}(\rho_{SM} \| \gamma_S)$, extraction $\beta^{-1} I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \| \gamma_S)$
 - erasure costs $\beta^{-1} H_{\max}^{\uparrow, \varepsilon}(S|M)_{\rho}$, which can be negative: cooling rather than heating

Conclusion

Fundamental limits on thermodynamic control with coherent (quantum) feedback, on both microscopic & macroscopic scales

- QFGOs: axiomatic and deliberately permissive $\Rightarrow$ the limits hold for every physical feedback-control scheme
- Single shot: exact work costs, with optimal protocols
 - formation $\beta^{-1} I_{\max}^{\uparrow, \varepsilon}(\rho_{SM} \parallel \gamma_S)$, extraction $\beta^{-1} I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \parallel \gamma_S)$
 - erasure costs $\beta^{-1} H_{\max}^{\uparrow, \varepsilon}(S|M)_{\rho}$, which can be negative: cooling rather than heating
- Asymptotic: generalized second law $\langle W \rangle \geq \Delta F_{\text{sys}} + \beta^{-1} \Delta I_q$
 - consistent with the traditional second law over a closed cycle

Conclusion

Fundamental limits on thermodynamic control with coherent (quantum) feedback, on both microscopic & macroscopic scales

- QFGOs: axiomatic and deliberately permissive $\Rightarrow$ the limits hold for every physical feedback-control scheme
- Single shot: exact work costs, with optimal protocols
 - formation $\beta^{-1} I_{\max}^{\uparrow, \varepsilon}(\rho_{SM} \| \gamma_S)$, extraction $\beta^{-1} I_{\min}^{\downarrow, \varepsilon}(\rho_{SM} \| \gamma_S)$
 - erasure costs $\beta^{-1} H_{\max}^{\uparrow, \varepsilon}(S|M)_{\rho}$, which can be negative: cooling rather than heating
- Asymptotic: generalized second law $\langle W \rangle \geq \Delta F_{\text{sys}} + \beta^{-1} \Delta I_q$
 - consistent with the traditional second law over a closed cycle
- Quantum feedback can strictly beat classical feedback (example: $\beta^{-1} \ln 2$ vs. $\leq \beta^{-1} \ln \frac{4}{3}$, even for a classical system, without entanglement)

Outlook

Outlook

- Our limits are attained within QFGOs, which are maximally permissive: are they attainable in practice, theoretically and experimentally?
 - e.g., a quantum-feedback-assisted generalization of thermal operations [Janzing et al. 2000] [Horodecki & Oppenheim 2013]

Outlook

- Our limits are attained within QFGOs, which are maximally permissive: are they attainable in practice, theoretically and experimentally?
 - e.g., a quantum-feedback-assisted generalization of thermal operations [Janzing et al. 2000] [Horodecki & Oppenheim 2013]
- Work costs of logical processes (a channel acting on a given input) [Faist et al. 2015] [Faist & Renner 2018]

Outlook

- Our limits are attained within QFGOs, which are maximally permissive: are they attainable in practice, theoretically and experimentally?
 - e.g., a quantum-feedback-assisted generalization of thermal operations [Janzing et al. 2000] [Horodecki & Oppenheim 2013]
- Work costs of logical processes (a channel acting on a given input) [Faist et al. 2015] [Faist & Renner 2018]
- Energetic costs of adaptive agents that receive feedback from their environment, with the agent and/or the environment quantum [Elliott et al. 2022] [Thompson et al. 2025]

Outlook

- Our limits are attained within QFGOs, which are maximally permissive: are they attainable in practice, theoretically and experimentally?
 - e.g., a quantum-feedback-assisted generalization of thermal operations [Janzing et al. 2000] [Horodecki & Oppenheim 2013]
- Work costs of logical processes (a channel acting on a given input) [Faist et al. 2015] [Faist & Renner 2018]
- Energetic costs of adaptive agents that receive feedback from their environment, with the agent and/or the environment quantum [Elliott et al. 2022] [Thompson et al. 2025]
- Quantum error correction (QEC) as feedback control
 - a round of QEC is a feedback-control cycle
 - measurement-based / measurement-free QEC = classical / coherent feedback [Heußen, Locher & Müller 2024]
 - ...

