Supplementary Table S4. Common palliative radiotherapy regimens and the respective BED and EQD2

Regimen	α/β ratio 3		α/β ratio 10	
	BED3	EQD2	BED10	EQD2
40 Gy in 10 fractions (current study)	93.3 Gy	56.0 Gy	56.0 Gy	46.7 Gy
8 Gy in 1 fraction	29.3 Gy	17.6 Gy	14.4 Gy	12.0 Gy
10 Gy in 1 fraction	43.3 Gy	26.0 Gy	20.0 Gy	16.7 Gy
16 Gy in 2 fractions	58.7 Gy	35.2 Gy	28.8 Gy	24.0 Gy
17 Gy in 2 fractions	65.2 Gy	39.1 Gy	31.5 Gy	26.2 Gy
20 Gy in 5 fractions	46.7 Gy	28.0 Gy	28.0 Gy	23.3 Gy
30 Gy in 10 fractions	60.0 Gy	36.0 Gy	39.0 Gy	32.5 Gy
36 Gy in 12 fractions	72.0 Gy	43.2 Gy	46.8 Gy	39.0 Gy
39 Gy in 13 fractions	78.0 Gy	46.8 Gy	50.7 Gy	42.3 Gy

BED, biologically effective dose; EQD2, equivalent dose at 2 Gy per fraction.