MTL104: Linear Algebra Spring 2020-21

Lecture 4

Date: 10 February 2021 Scribe: Dhananjay Kajla

Note: LaTeX template courtesy of UC Berkeley EECS dept.

Disclaimer: These notes are **unofficial** and were meant for personal use. Therefore accuracy of these notes are not guaranteed. And therefore the liability for any factual errors does not lie with either the author or the instructor.

4.1 Properties of subspaces

- V(F) is also a subspace, so is $\{0\}$.
- Let V be the vector space \mathbb{R}^3 . Then the set W consisting of those vectors whose third component is zero, i.e. $w = \{a, b, 0 : a, b \in \mathbb{R}\}$ is a subspace of \mathbb{R}^3
- Let V be the vector space of all $n \times n$ matrices. Then the set W consisting of these matrices $A = [a_{ij}]$ for which $a_{ji} = a_{ij}$ (Symmetric matrices) is a subspace of V.
- Let V be the vector space of polynomials. Then the set W consisting of polynomials with degree $\leq n$, for a fixed n, is a subspace of V.
- Let V be the vector space of all functions for a non-empty set X into the real field \mathbb{R} . Then the set consisting of all bounded functions in V is a subspace of V.
 - A function $f \in V$ is bounded iff $\exists M \in R$ such that $|f(x)| \leq M$
- Let S be a non-empty subset of V(F), The set of all linear comination of vectors in S, denoted by L(S), is a subspace of V containing S.
- Furthermore, if W is any other subspace of V containing S, then $L(S) \leq W$.
- The solution space of a system of linear equations $\subseteq R^{n\times 1}$
- In F^n , the set of all n-tuples $(x_1, x_2, ..., x_n)$ with $x_1 = 0$ is a subspace.
- The set of all hermitian matrices is **NOT** a subspace of the space of all $n \times n$ matrices over \mathbb{C} . The set of all $n \times n$ complex hermitian matrices is a vector space over the field of real numbers.
- Suppose W_1 and W_2 are subspaces of a vector space V(F)

•