

## Lecture 3

Date : 8 February 2021

Scribe: Dhananjay Kajla

**Note:** *LaTeX template courtesy of UC Berkeley EECS dept.*

**Disclaimer:** These notes are **unofficial** and were meant for personal use. Therefore accuracy of these notes are not guaranteed. And therefore the liability for any factual errors does not lie with either the author or the instructor.

## 3.1 Linear Combinations and Independence

### 3.1.1 Linear Independence

- A finite set  $\{x_1, x_2, \dots, x_n\}$  of vectors in  $V$  is called linearly independent iff :

$$\alpha_1 x_1 + \alpha_2 x_2 + \alpha_3 x_3 \dots + \alpha_n x_n = \mathbf{0} \implies \alpha_1 = \alpha_2 = \dots = \alpha_n = 0$$

- Let  $X \subseteq V$ ,  $X$  is said to be linearly independent if every finite subset of  $X$  is linearly independent.

### 3.1.2 Properties of Linear Independence

- A set of vectors which contains  $\mathbf{0}$  as one of its elements is linearly dependent.
- The set  $\{x\}$  is linearly independent iff  $x \neq 0$
- Any subset of a linearly independent set is linearly independent.
- Any superset of a linearly dependent set is linearly dependent.
- If  $\{x_1, x_2, \dots, x_n\}$  is linearly independent and if  $\{a_1, a_2, \dots, a_n\} \in F$  and  $\{b_1, b_2, \dots, b_n\} \in F$ , then

$$a_1 x_1 + a_2 x_2 + a_3 x_3 + \dots + a_n x_n = b_1 x_1 + b_2 x_2 + b_3 x_3 + \dots + b_n x_n \implies a_1 = b_1, a_2 = b_2, \dots, a_n = b_n$$

- **Lemma :** The set of non-zero vectors  $\{x_1, x_2, \dots, x_n\} \in V$  is linearly dependent if and only if any one of them, say  $x_j$  can be expressed as a *linear combination* of the other vectors.

### 3.1.3 Linear Combination of vectors

- If  $V(F)$  is a vector space and  $\{x_1, x_2, \dots, x_n\} \in V$  and  $\{\alpha_1, \alpha_2, \dots, \alpha_n\} \in F$ , then the vector  $x = \alpha_1 x_1 + \alpha_2 x_2 + \dots + \alpha_n x_n = \sum_{i=1}^n (\alpha_i x_i)$  is called the linear combination of vectors  $\{x_1, x_2, \dots, x_n\}$

### 3.1.4 Linear Span of a set

- The linear span of a non-empty subset  $S$  of  $V(F)$  is the set of all linear combinations of any finite number of elements of  $S$  and is denoted by  $L(S)$  (Some textbooks denote it as  $\langle S \rangle$ )

### 3.1.5 Basis of a set

- If  $S$  is a subset of a vector space  $V(F)$  such that :
  1.  $S$  is a linearly independent set of vectors of  $V(F)$
  2. Every vector  $v$  in  $V(F)$  is a linear combination of elements of  $S$  or in other words  $L(S) = V$ , then  $S$  is called the basis of  $V(F)$ .
- Zero vector ( $\mathbf{0}$ ) cannot be an element of any basis of  $V(F)$

### 3.1.6 Finitely Generated Vector Spaces

- A vector space  $V(F)$  is said to be finitely generated if there exists a finite subset  $S$  of  $V$  such that  $V = L(S)$  or if  $V(S)$  has a finite spanning set.
- Equivalently a vector space is finitely generated if it's basis is finite.
- In a finitely generate vector space  $V(F)$ , whose basis set is  $B = x_1, x_2, \dots, x_n$ , every vector  $x \in V$  is uniquely expressible as a linear combination of vectors in  $B$ .
- **Existence Theorem :** There exists a basis for each finite dimensional vector space.

### 3.1.7 Dimension of a vector space

- The dimension of a finitely generated vector space  $V(F)$  is the cardinality of any basis of  $V(F)$  and is represented as  $\dim(V)$ .
- **Extension Theorem :** If  $V(F)$  is a finitely generated vector space, and  $A = x_1, x_2, \dots, x_m$  is any linearly independent set of vectors in  $V$ , then unless the  $x_i$ s already form a basis, we can find the vectors  $\{y_1, y_2, \dots, y_{n-m}\}$  so that the extended set of  $n$  vectors  $\{x_1, x_2, \dots, x_m, y_1, y_2, \dots, y_{n-m}\}$  is a basis.
- In a finitely generated vector space, the cardinality of every basis is the same and is equal to the dimension of the vector space.

### 3.1.8 Subspace of a vector space

- A subset  $W$  of a vector space  $V$  over a field  $F$  is called a subspace of  $V$  if  $W$  is a vector space over  $F$  with the same operations of addition and multiplication as defined for  $V$ .
- **Theorem :** Let  $V$  be a vector space and  $W$  be a subset of  $V$ . Then  $W$  is a subspace of  $V$  if and only if the following three conditions hold for operations defined in  $V$ .
  1.  $\mathbf{0} \in W$
  2.  $x + y \in W$  whenever  $x \in W$  and  $y \in W$
  3.  $\alpha x \in W$  whenever  $x \in W$  and  $\alpha \in F$