

Lecture 4

Date : 10 February 2021

Scribe: Dhananjay Kajla

Note: *LaTeX template courtesy of UC Berkeley EECS dept.*

Disclaimer: These notes are **unofficial** and were meant for personal use. Therefore accuracy of these notes are not guaranteed. And therefore the liability for any factual errors does not lie with either the author or the instructor.

4.1 Subspaces

4.1.1 Examples of subspaces

- $V(F)$ is also a subspace, so is $\{0\}$.
- Let V be the vector space \mathbb{R}^3 . Then the set W consisting of those vectors whose third component is zero, i.e. $w = \{a, b, 0 : a, b \in \mathbb{R}\}$ is a subspace of \mathbb{R}^3
- Let V be the vector space of all $n \times n$ matrices. Then the set W consisting of these matrices $A = [a_{ij}]$ for which $a_{ji} = a_{ij}$ (Symmetric matrices) is a subspace of V .
- Let V be the vector space of polynomials. Then the set W consisting of polynomials with degree $\leq n$, for a fixed n , is a subspace of V .
- Let V be the vector space of all functions for a non-empty set X into the real field \mathbb{R} . Then the set consisting of all bounded functions in V is a subspace of V .
 - A function $f \in V$ is bounded iff $\exists M \in \mathbb{R}$ such that $|f(x)| \leq M$
- Let S be a non-empty subset of $V(F)$, The set of all linear combination of vectors in S , denoted by $L(S)$, is a subspace of V containing S .
- Furthermore, if W is any other subspace of V containing S , then $L(S) \subseteq W$.
- The solution space of a system of linear equations $\subseteq \mathbb{R}^{n \times 1}$
- In F^n , the set of all n -tuples (x_1, x_2, \dots, x_n) with $x_1 = 0$ is a subspace.
- The set of all hermitian matrices is **NOT** a subspace of the space of all $n \times n$ matrices over \mathbb{C} . The set of all $n \times n$ complex hermitian matrices is a vector space over the field of real numbers.

4.1.2 Few Properties of subspaces

1. Suppose W_1 and W_2 are subspaces of a vector space $V(F)$, Then
 - $W_1 \cup W_2$ need not be a subspace of V
 - For example. Consider the vector space of \mathbb{R}^2 , with $W_1 = \{(a, 0); a \in \mathbb{R}\}$ and $W_2 = \{(0, b); b \in \mathbb{R}\}$. Then $(1, 0) \in W_1$ and $(0, 1) \in W_2$, but their addition $(1, 0) + (0, 1) = (1, 1) \notin W_1 \cup W_2$.

- Hence addition is not closed, hence $W_1 \cup W_2$ is not a subspace of V
- $W_1 \cap W_2$ is a subspace of V . Proof :
 - $\mathbf{0} \in W_1, \mathbf{0} \in W_2 \implies \mathbf{0} \in W_1 \cap W_2$
 - Let $x, y \in W_1 \cap W_2, \alpha \in F$,
 - Now $x \in W_1 \cap W_2 \implies x \in W_1$ and $x \in W_2$
 - Similarly $y \in W_1 \cap W_2 \implies y \in W_1$ and $y \in W_2$
 - Since W_1 is a subspace, $\alpha(x + y) \in W_1$
 - Similarly W_2 is a subspace, so $\alpha(x + y) \in W_2$
 - Therefore $\alpha(x + y) \in W_1 \cap W_2$
- 2. Let V be a vector space over the field F , then intersection of any collection of subspaces (i.e. Arbitrary intersection of subspaces) is a subspace in V .

4.1.3 Linear Sum of Subspaces

- Let W_1 and W_2 be two subspaces of a vector space $V(F)$, Then the linear sum of subspaces, denoted by $W_1 + W_2$ is defined as :

$$W_1 + W_2 = \{x_1 + x_2 : x_1 \in W_1, x_2 \in W_2\}$$

- **Theorem :** If W_1 and W_2 are subspaces of a vector space $V(F)$, then $W_1 + W_2$ is a subspace of $V(F)$
- Proof :

- $\mathbf{0} = \mathbf{0} + \mathbf{0}$ where $\mathbf{0} \in W_1$ and $\mathbf{0} \in W_2$
- let $x, y \in W_1 + W_2, \alpha \in F$, then :
 - * $x = x_1 + x_2$ where $x_1 \in W_1$ and $x_2 \in W_2$
 - * $y = y_1 + y_2$ where $y_1 \in W_1$ and $y_2 \in W_2$
- Consider $\alpha x + y = \alpha(x_1 + x_2) + (y_1 + y_2) = (\alpha x_1 + y_1) + (\alpha x_2 + y_2)$
 - * Now $(\alpha x_1 + y_1) \in W_1$ as W_1 is a subspace
 - * Similarly $(\alpha x_2 + y_2) \in W_2$ as W_2 is a subspace
 - * Therefore $\alpha x + y \in W_1 + W_2$
- Therefore $W_1 + W_2$ is a subspace of V

- If W_1, W_2, \dots, W_k are subspaces of the vector space $V(F)$. Then their linear sum, denote by $W_1 + W_2 + \dots + W_k$ is defined as :

$$W_1 + W_2 + \dots + W_k = \{x_1 + x_2 + \dots + x_k : x_i \in W_i, 1 \leq i \leq k\}$$

- $W_1 + W_2 + \dots + W_k$ is a subspace of $V(F)$

4.1.4 Direct Sum of Subspaces

- The vector space V over the field F is the direct sum of two vector spaces W_1 and W_2 if
 1. $V = W_1 + W_2$
 2. Every vector z in V can be uniquely expressed as the sum of $x_1 + x_2$ where $x_1 \in W_1$ and $x_2 \in W_2$
- If the vector space $V(F)$ is the direct sum of two subspaces, W_1 and W_2 , we write,

$$V = W_1 \oplus W_2$$

- **Theorem :** $V = W_1 \oplus W_2 \iff V = W_1 + W_2$ and $W_1 \cap W_2 = \{0\}$