

Theoretical papers are [1–3].

X-ray diffraction papers are [4, 5].

Bibliography

- [1] T. C. Lubensky and F. C. MacKintosh, “Theory of ripple phases of lipid bilayers,” *Physical Review Letters*, vol. 71, no. 10, p. 1565, 1993.
- [2] C. M. Chen, T. C. Lubensky, and F. C. MacKintosh, “Phase transitions and modulated phases in lipid bilayers,” *Physical Review E*, vol. 51, no. 1, p. 504, 1995.
- [3] M. A. Kamal, A. Pal, V. A. Raghunathan, and M. Rao, “Theory of the asymmetric ripple phase in achiral lipid membranes,” *Europhysics Letters*, vol. 95, no. 4, p. 48004, 2011.
- [4] W. J. Sun, S. Tristram-Nagle, R. M. Suter, and J. F. Nagle, “Structure of the ripple phase in lecithin bilayers,” *Proceedings of the National Academy of Sciences*, vol. 93, no. 14, pp. 7008–7012, 1996.
- [5] K. Sengupta, V. A. Raghunathan, and J. Katsaras, “Structure of the ripple phase of phospholipid multibilayers,” *Physical Review E*, vol. 68, p. 031710, Sep 2003.
- [6] K. Akabori and J. F. Nagle, “Comparing lipid membranes in different environments,” *ACS Nano*, vol. 8, no. 4, pp. 3123–3127, 2014.
- [7] U. Bernchou, H. Midtby, J. H. Ipsen, and A. C. Simonsen, “Correlation between the ripple phase and stripe domains in membranes,” *Biochimica et Biophysica Acta (BBA) - Biomembranes*, vol. 1808, no. 12, pp. 2849 – 2858, 2011.
- [8] J. P. Bradshaw, M. S. Edenborough, P. J. Sizer, and A. Watts, “Observation of rippled dioleoylphosphatidylcholine bilayers by neutron diffraction,” *Biochimica et Biophysica Acta (BBA)-Biomembranes*, vol. 987, no. 1, pp. 111–114, 1989.

- [9] J. M. Carlson and J. P. Sethna, “Theory of the ripple phase in hydrated phospholipid bilayers,” *Physical Review A*, vol. 36, no. 7, p. 3359, 1987.
- [10] G. Cevc, “Polymorphism of the bilayer membranes in the ordered phase and the molecular origin of the lipid pretransition and rippled lamellae,” *Biochimica et Biophysica Acta (BBA)-Biomembranes*, vol. 1062, no. 1, pp. 59–69, 1991.
- [11] D. Chapman, R. M. Williams, and B. D. Ladbroke, “Physical studies of phospholipids. vi. thermotropic and lyotropic mesomorphism of some 1, 2-diacyl-phosphatidylcholines (lecithins),” *Chemistry and Physics of Lipids*, vol. 1, no. 5, pp. 445–475, 1967.
- [12] L. V. Chernomordik and M. M. Kozlov, “Protein-lipid interplay in fusion and fission of biological membranes,” *Annual review of biochemistry*, vol. 72, no. 1, pp. 175–207, 2003.
- [13] S. D. Connell and D. A. Smith, “The atomic force microscope as a tool for studying phase separation in lipid membranes (review),” *Molecular membrane biology*, vol. 23, no. 1, pp. 17–28, 2006.
- [14] B. R. Copeland and H. M. McConnell, “The rippled structure in bilayer membranes of phosphatidylcholine and binary mixtures of phosphatidylcholine and cholesterol,” *Biochimica et Biophysica Acta (BBA) - Biomembranes*, vol. 599, no. 1, pp. 95 – 109, 1980.
- [15] B. A. Cunningham, A.-D. Brown, D. H. Wolfe, W. P. Williams, and A. Brain, “Ripple phase formation in phosphatidylcholine: Effect of acyl chain relative length, position, and unsaturation,” *Physical Review E*, vol. 58, no. 3, p. 3662, 1998.
- [16] <http://henke.lbl.gov/optical.constants>.
- [17] D. M. Czajkowsky, C. Huang, and Z. Shao, “Ripple phase in asymmetric unilamellar bilayers with saturated and unsaturated phospholipids,” *Biochemistry*, vol. 34, no. 39, pp. 12501–12505, 1995.
- [18] L. Dahbi, M. Arbel-Haddad, P. Lesieur, C. Bourgaux, and M. Ollivon, “A long ripple phase in dlpcdecylglucoside mixture evidenced by synchrotron {SAXS} coupled to {DSC},” *Chemistry and Physics of Lipids*, vol. 139, no. 1, pp. 43 – 53, 2006.

- [19] A. Debnath, K. G. Ayappa, V. Kumaran, and P. K. Maiti, “The influence of bilayer composition on the gel to liquid crystalline transition,” *The Journal of Physical Chemistry B*, vol. 113, no. 31, pp. 10660–10668, 2009. PMID: 19594148.
- [20] A. H. de Vries, S. Yefimov, A. E. Mark, and S. J. Marrink, “Molecular structure of the lecithin ripple phase,” *Proceedings of the National Academy of Sciences*, vol. 102, no. 15, pp. 5392–5396, 2005.
- [21] S. Doniach, “A thermodynamic model for the monoclinic (ripple) phase of hydrated phospholipid bilayers,” *The Journal of Chemical Physics*, vol. 70, no. 10, pp. 4587–4596, 1979.
- [22] W. Dowhan, “Molecular basis for membrane phospholipid diversity: why are there so many lipids?,” *Annual review of biochemistry*, vol. 66, no. 1, pp. 199–232, 1997.
- [23] P. F. Fahey and W. W. Webb, “Lateral diffusion in phospholipid bilayer membranes and multilamellar liquid crystals,” *Biochemistry*, vol. 17, no. 15, pp. 3046–3053, 1978.
- [24] R. E. Goldstein and S. Leibler, “Model for lamellar phases of interacting lipid membranes,” *Physical Review Letters*, vol. 61, no. 19, p. 2213, 1988.
- [25] S. Guler, D. D. Ghosh, J. Pan, J. C. Mathai, M. L. Zeidel, J. F. Nagle, and S. Tristram-Nagle, “Effects of ether vs. ester linkage on lipid bilayer structure and water permeability,” *Chemistry and Physics of Lipids*, vol. 160, no. 1, pp. 33–44, 2009.
- [26] F. Harb, A. Simon, and B. Tinland, “Ripple formation in unilamellar-supported lipid bilayer revealed by frapp,” *The European Physical Journal E*, vol. 36, no. 12, pp. 1–7, 2013.
- [27] M. H. Hawton and W. J. Keeler, “van der Waals energy of lecithins in the ripple phase,” *Physical Review A*, vol. 33, no. 5, p. 3333, 1986.
- [28] T. Heimburg, “A model for the lipid pretransition: coupling of ripple formation with the chain-melting transition,” *Biophysical Journal*, vol. 78, no. 3, pp. 1154–1165, 2000.

- [29] M. P. Hentschel and F. Rustichelli, “Structure of the ripple phase $P_{\beta'}$ in hydrated phosphatidylcholine multimembranes,” *Physical Review Letters*, vol. 66, pp. 903–906, Feb 1991.
- [30] K. Honda and H. Kimura, “Theory on formation of the ripple phase in bilayer membranes,” *Journal of the Physical Society of Japan*, vol. 60, no. 4, pp. 1212–1215, 1991.
- [31] M. J. Janiak, D. M. Small, and G. G. Shipley, “Nature of the thermal pretransition of synthetic phospholipids: dimyristoyl- and dipalmitoyl-lecithin,” *Biochemistry*, vol. 15, no. 21, pp. 4575–4580, 1976.
- [32] M. J. Janiak, D. M. Small, and G. G. Shipley, “Temperature and compositional dependence of the structure of hydrated dimyristoyl lecithin,” *Journal of Biological Chemistry*, vol. 254, no. 13, pp. 6068–6078, 1979.
- [33] M. A. Kamal, A. Pal, V. A. Raghunathan, and M. Rao, “Phase behavior of two-component lipid membranes: Theory and experiments,” *Physical Review E*, vol. 85, no. 5, p. 051701, 2012.
- [34] M. A. Kamal, A. Pal, and V. A. Raghunathan, “Modulated phases of lipid membranes induced by sterol derivatives,” *Soft Matter*, vol. 8, no. 46, pp. 11600–11603, 2012.
- [35] S. Karmakar and V. A. Raghunathan, “Cholesterol-induced modulated phase in phospholipid membranes,” *Physical Review Letters*, vol. 91, no. 9, p. 098102, 2003.
- [36] S. Karmakar and V. A. Raghunathan, “Structure of phospholipid-cholesterol membranes: An X-ray diffraction study,” *Physical Review E*, vol. 71, no. 6, pp. 061924.1–061924.10, 2005.
- [37] T. Kaasgaard, C. Leidy, J. H. Crowe, O. G. Mouritsen, and K. Jørgensen, “Temperature-controlled structure and kinetics of ripple phases in one-and two-component supported lipid bilayers,” *Biophysical Journal*, vol. 85, no. 1, pp. 350–360, 2003.
- [38] J. Katsaras and V. A. Raghunathan, “Molecular chirality and the ripple phase of phosphatidylcholine multibilayers,” *Physical Review Letters*, vol. 74, pp. 2022–2025, Mar 1995.

- [39] J. Katsaras, S. Tristram-Nagle, Y. Liu, R. Headrick, E. Fontes, P. Mason, and J. F. Nagle, “Clarification of the ripple phase of lecithin bilayers using fully hydrated, aligned samples,” *Physical Review E*, vol. 61, no. 5, p. 5668, 2000.
- [40] M. A. Kiselev, E. V. Zemlyanaya, V. K. Aswal, and R. H. Neubert, “What can we learn about the lipid vesicle structure from the small-angle neutron scattering experiment?,” *European Biophysics Journal*, vol. 35, no. 6, pp. 477–493, 2006.
- [41] R. Koynova, A. Koumanov, and B. Tenchov, “Metastable rippled gel phase in saturated phosphatidylcholines: calorimetric and densitometric characterization,” *Biochimica et Biophysica Acta (BBA) - Biomembranes*, vol. 1285, no. 1, pp. 101 – 108, 1996.
- [42] R. Koynova and B. Tenchov, “Transitions between lamellar and non-lamellar phases in membrane lipids and their physiological roles,”
- [43] B. D. Ladbroke and D. Chapman, “Thermal analysis of lipids, proteins and biological membranes a review and summary of some recent studies,” *Chemistry and Physics of Lipids*, vol. 3, no. 4, pp. 304–356, 1969.
- [44] C. Leidy, T. Kaasgaard, J. H. Crowe, O. G. Mouritsen, and K. Jørgensen, “Ripples and the formation of anisotropic lipid domains: imaging two-component supported double bilayers by atomic force microscopy,” *Biophysical Journal*, vol. 83, no. 5, pp. 2625–2633, 2002.
- [45] O. Lenz and F. Schmid, “Structure of symmetric and asymmetric “ripple” phases in lipid bilayers,” *Physical Review Letters*, vol. 98, p. 058104, Jan 2007.
- [46] L. Li and J.-X. Cheng, “Coexisting stripe-and patch-shaped domains in giant unilamellar vesicles,” *Biochemistry*, vol. 45, no. 39, pp. 11819–11826, 2006.
- [47] <http://cars9.uchicago.edu/software/python/lmfit/>.
- [48] E. J. Luna and H. M. McConnell, “The intermediate monoclinic phase of phosphatidylcholines,” *Biochimica et Biophysica Acta (BBA)-Biomembranes*, vol. 466, no. 3, pp. 381–392, 1977.

- [49] M. Marder, H. L. Frisch, J. S. Langer, and H. M. McConnell, "Theory of the intermediate rippled phase of phospholipid bilayers," *Proceedings of the National Academy of Sciences*, vol. 81, no. 20, pp. 6559–6561, 1984.
- [50] D. Marsh, "Molecular motion in phospholipid bilayers in the gel phase: long axis rotation," *Biochemistry*, vol. 19, no. 8, pp. 1632–1637, 1980.
- [51] D. Marsh, "Lateral order in gel, subgel and crystalline phases of lipid membranes: Wide-angle X-ray scattering," *Chemistry and Physics of Lipids*, vol. 165, no. 1, pp. 59–76, 2012.
- [52] P. C. Mason, B. D. Gaulin, R. M. Epand, G. D. Wignall, and J. S. Lin, "Small angle neutron scattering and calorimetric studies of large unilamellar vesicles of the phospholipid dipalmitoylphosphatidylcholine," *Physical Review E*, vol. 59, no. 3, p. 3361, 1999.
- [53] P. C. Mason, B. D. Gaulin, R. M. Epand, G. D. Wignall, and J. S. Lin, "Small-angle scattering studies of the fully hydrated phospholipid DPPC," *Physical Review E*, vol. 59, no. 1, p. 921, 1999.
- [54] P. C. Mason, J. F. Nagle, R. M. Epand, and J. Katsaras, "Anomalous swelling in phospholipid bilayers is not coupled to the formation of a ripple phase," *Physical Review E*, vol. 63, no. 3, p. 030902, 2001.
- [55] T. J. McIntosh, "Differences in hydrocarbon chain tilt between hydrated phosphatidylethanolamine and phosphatidylcholine bilayers. a molecular packing model," *Biophysical Journal*, vol. 29, no. 2, pp. 237–245, 1980.
- [56] W. S. McCullough and H. L. Scott, "Statistical-mechanical theory of the ripple phase of lipid bilayers," *Physical Review Letters*, vol. 65, pp. 931–934, Aug 1990.
- [57] C. E. Miller, J. Majewski, E. B. Watkins, D. J. Mulder, T. Gog, and T. L. Kuhl, "Probing the local order of single phospholipid membranes using grazing incidence X-ray diffraction," *Physical Review Letters*, vol. 100, no. 5, p. 058103, 2008.
- [58] T. T. Mills, G. E. Toombes, S. Tristram-Nagle, D.-M. Smilgies, G. W. Feigenson, and J. F. Nagle, "Order parameters and areas in fluid-phase

- oriented lipid membranes using wide angle X-ray scattering,” *Biophysical Journal*, vol. 95, no. 2, pp. 669–681, 2008.
- [59] C. Misbah, J. Duplat, and B. Houchmandzadeh, “Transition to ripple phases in hydrated amphiphiles,” *Physical Review Letters*, vol. 80, no. 20, p. 4598, 1998.
 - [60] K. Mortensen, W. Pfeiffer, E. Sackmann, and W. Knoll, “Structural properties of a phosphatidylcholine-cholesterol system as studied by small-angle neutron scattering: ripple structure and phase diagram,” *Biochimica et Biophysica Acta (BBA)-Biomembranes*, vol. 945, no. 2, pp. 221–245, 1988.
 - [61] J. F. Nagle and S. Tristram-Nagle, “Structure of lipid bilayers,” *Biochimica et Biophysica Acta (BBA) - Reviews on Biomembranes*, vol. 1469, no. 3, pp. 159 – 195, 2000.
 - [62] J. Nagle, “Theory of lipid monolayer and bilayer phase transitions: Effect of headgroup interactions,” *The Journal of Membrane Biology*, vol. 27, no. 1, pp. 233–250, 1976.
 - [63] J. F. Nagle, “Theory of the main lipid bilayer phase-transition,” *Annual Review of Physical Chemistry*, vol. 31, pp. 157–195, 1980.
 - [64] G. Pabst, H. Amenitsch, D. P. Kharakoz, P. Laggner, and M. Rappolt, “Structure and fluctuations of phosphatidylcholines in the vicinity of the main phase transition,” *Physical Review E*, vol. 70, no. 2, p. 021908, 2004.
 - [65] R. A. Parente and B. R. Lentz, “Phase behavior of large unilamellar vesicles composed of synthetic phospholipids,” *Biochemistry*, vol. 23, no. 11, pp. 2353–2362, 1984.
 - [66] S. Potekhin, A. Senin, N. Abdurakhmanov, and R. Khusainova, “High pressure effect on the main transition from the ripple gel p phase to the liquid crystal (l) phase in dipalmitoylphosphatidylcholine. microcalorimetric study,” *Biochimica et Biophysica Acta (BBA) - Biomembranes*, vol. 1778, no. 11, pp. 2588 – 2593, 2008.

- [67] M. Rappolt and G. Rapp, “Structure of the stable and metastable ripple phase of dipalmitoylphosphatidylcholine,” *European Biophysics Journal*, vol. 24, no. 6, pp. 381–386, 1996.
- [68] M. Rappolt, G. Pabst, G. Rapp, M. Kriechbaum, H. Amenitsch, C. Krenn, S. Bernstorff, and P. Lagner, “New evidence for gel-liquid crystalline phase coexistence in the ripple phase of phosphatidylcholines,” *European Biophysics Journal*, vol. 29, no. 2, pp. 125–133, 2000.
- [69] S. Rifci, C. Crupi, G. D’Angelo, G. Di Marco, G. Sabatino, V. Conti Nibali, A. Trimarchi, and U. Wanderlingh, “Effects of a short length alcohol on the dimyristoylphosphatidylcholine system,” *Philosophical Magazine*, vol. 91, no. 13-15, pp. 2014–2020, 2011.
- [70] K. A. Riske, R. P. Barroso, C. C. Vequi-Suplicy, R. Germano, V. B. Henriques, and M. T. Lamy, “Lipid bilayer pre-transition as the beginning of the melting process,” *Biochimica et Biophysica Acta (BBA) - Biomembranes*, vol. 1788, no. 5, pp. 954 – 963, 2009.
- [71] D. Ruppel and E. Sackmann, “On defects in different phases of two-dimensional lipid bilayers,” *Journal de Physique*, vol. 44, no. 9, pp. 1025–1034, 1983.
- [72] F. Schmid, S. Dolezel, O. Lenz, and S. Meinhardt, “On ripples and rafts: Curvature induced nanoscale structures in lipid membranes,” in *Journal of Physics Conference Series*, vol. 487, p. 2004, 2014.
- [73] H. L. Scott, “Monte carlo studies of a general model for lipid bilayer condensed phases,” *The Journal of Chemical Physics*, vol. 80, no. 5, pp. 2197–2202, 1984.
- [74] K. Sengupta, V. Raghunathan, and J. Katsaras, “Structure of the ripple phase in chiral and racemic dimyristoylphosphatidylcholine multibilayers,” *Physical Review E*, vol. 59, no. 2, p. 2455, 1999.
- [75] K. Sengupta, V. Raghunathan, and J. Katsaras, “Novel structural features of the ripple phase of phospholipids,” *Europhysics Letters*, vol. 49, no. 6, p. 722, 2000.

- [76] K. Sengupta, V. Raghunathan, and Y. Hatwalne, “Role of tilt order in the asymmetric ripple phase of phospholipid bilayers,” *Physical Review Letters*, vol. 87, no. 5, pp. 055705_1–055705_4, 2001.
- [77] A. Schäfer, T. Salditt, and M. C. Rheinstädter, “Atomic force microscopy study of thick lamellar stacks of phospholipid bilayers,” *Physical Review E*, vol. 77, no. 2, p. 021905, 2008.
- [78] M. B. Schneider, W. K. Chan, and W. W. Webb, “Fast diffusion along defects and corrugations in phospholipid $P_{\beta'}$, liquid crystals,” *Biophysical Journal*, vol. 43, no. 2, pp. 157–165, 1983.
- [79] G. S. Smith, E. B. Sirota, C. R. Safinya, and N. A. Clark, “Structure of the $L_{\beta'}$ phases in a hydrated phosphatidylcholine multilamellar membrane,” *Physical Review Letters*, vol. 60, pp. 813–816, Feb 1988.
- [80] W. J. Sun, R. M. Suter, M. A. Knewtson, C. R. Worthington, S. Tristram-Nagle, R. Zhang, and J. F. Nagle, “Order and disorder in fully hydrated unoriented bilayers of gel-phase dipalmitoylphosphatidylcholine,” *Physical Review E*, vol. 49, pp. 4665–4676, May 1994.
- [81] A. Tardieu, V. Luzzati, and F. Reman, “Structure and polymorphism of the hydrocarbon chains of lipids: A study of lecithin-water phases,” *Journal of Molecular Biology*, vol. 75, no. 4, pp. 711 – 733, 1973.
- [82] H. Truble and D. H. Haynes, “The volume change in lipid bilayer lamellae at the crystalline-liquid crystalline phase transition,” *Chemistry and Physics of Lipids*, vol. 7, no. 4, pp. 324 – 335, 1971.
- [83] S. Tristram-Nagle, R. Zhang, R. M. Suter, C. R. Worthington, W. J. Sun, and J. F. Nagle, “Measurement of chain tilt angle in fully hydrated bilayers of gel phase lecithins,” *Biophysical Journal*, vol. 64, no. 4, pp. 1097–1109, 1993.
- [84] S. Tristram-Nagle, Y. Isaacson, Y. Lyatskaya, Y. Liu, K. Brummond, J. Katsaras, and J. F. Nagle, “Polymorphism in myristoylpalmitoylphosphatidylcholine,” *Chemistry and Physics of Lipids*, vol. 100, no. 1, pp. 101–113, 1999.

- [85] S. Tristram-Nagle, Y. Liu, J. Legleiter, and J. F. Nagle, "Structure of gel phase DMPC determined by X-ray diffraction," *Biophysical Journal*, vol. 83, no. 6, pp. 3324 – 3335, 2002.
- [86] S. A. Tristram-Nagle, "Preparation of oriented, fully hydrated lipid samples for structure determination using X-ray scattering," *Methods of Molecular Biology*, vol. 400, pp. 63–75, 2007.
- [87] G. H. Vineyard, "Grazing-incidence diffraction and the distorted-wave approximation for the study of surfaces," *Physical Review B*, vol. 26, no. 8, p. 4146, 1982.
- [88] D. C. Wack and W. W. Webb, "Synchrotron X-ray study of the modulated lamellar phase $P_{\beta'}$ in the lecithin-water system," *Physical Review A*, vol. 40, pp. 2712–2730, Sep 1989.
- [89] B. E. Warren, *X-ray Diffraction*. Courier Dover Publications, 1969.
- [90] E. Watkins, C. Miller, J. Majewski, and T. Kuhl, "Membrane texture induced by specific protein binding and receptor clustering: active roles for lipids in cellular function," *Proceedings of the National Academy of Sciences*, vol. 108, no. 17, pp. 6975–6980, 2011.
- [91] E. B. Watkins, C. E. Miller, W.-P. Liao, and T. L. Kuhl, "Equilibrium or quenched: Fundamental differences between lipid monolayers, supported bilayers, and membranes," *ACS Nano*, vol. 8, no. 4, pp. 3181–3191, 2014.
- [92] M. C. Wiener, R. M. Suter, and J. F. Nagle, "Structure of the fully hydrated gel phase of dipalmitoylphosphatidylcholine," *Biophysical Journal*, vol. 55, no. 2, pp. 315 – 325, 1989.
- [93] R. Wittebort, C. Schmidt, and R. Griffin, "Solid-state carbon-13 nuclear magnetic resonance of the lecithin gel to liquid-crystalline phase transition," *Biochemistry*, vol. 20, no. 14, pp. 4223–4228, 1981.
- [94] J. Woodward IV and J. Zasadzinski, "Amplitude, wave form, and temperature dependence of bilayer ripples in the $P_{\beta'}$ phase," *Physical Review E*, vol. 53, no. 4, p. R3044, 1996.

- [95] J. Woodward IV and J. Zasadzinski, “High-resolution scanning tunneling microscopy of fully hydrated ripple-phase bilayers,” *Biophysical Journal*, vol. 72, no. 2, pp. 964–976, 1997.
- [96] C. Worthington, G. King, and T. McIntosh, “Direct structure determination of multilayered membrane-type systems which contain fluid layers,” *Biophysical Journal*, vol. 13, no. 5, pp. 480 – 494, 1973.
- [97] L. Yang and M. Fukuto, “Modulated phase of phospholipids with a two-dimensional square lattice,” *Physical Review E*, vol. 72, no. 1, p. 010901, 2005.
- [98] H. Yao, S. Matuoka, B. Tenchov, and I. Hatta, “Metastable ripple phase of fully hydrated dipalmitoylphosphatidylcholine as studied by small angle X-ray scattering,” *Biophysical Journal*, vol. 59, no. 1, pp. 252–255, 1991.
- [99] J. Zasadzinski and M. Schneider, “Ripple wavelength, amplitude, and configuration in lyotropic liquid crystals as a function of effective head-group size,” *Journal de Physique*, vol. 48, no. 11, pp. 2001–2011, 1987.
- [100] J. Zasadzinski, J. Schneir, J. Gurley, V. Elings, and P. Hansma, “Scanning tunneling microscopy of freeze-fracture replicas of biomembranes,” *Science*, vol. 239, no. 4843, pp. 1013–1015, 1988.
- [101] R. Zhang, W. J. Sun, S. Tristram-Nagle, R. L. Headrick, R. M. Suter, and J. F. Nagle, “Critical fluctuations in membranes,” *Physical Review Letters*, vol. 74, no. 14, p. 2832, 1995.
- [102] J. L. Baker, L. H. Jimison, S. Mannsfeld, S. Volkman, S. Yin, V. Subramanian, A. Salleo, A. P. Alivisatos, and M. F. Toney, “Quantification of thin film crystallographic orientation using X-ray diffraction with an area detector,” *Langmuir*, vol. 26, no. 11, pp. 9146–9151, 2010.
- [103] A. B. Rodriguez-Navarro, “Registering pole figures using an X-ray single-crystal diffractometer equipped with an area detector,” *Journal of Applied Crystallography*, vol. 40, pp. 631–634, Jun 2007.