MAC 1147 TEST IA SPRING 2012

- A. Sign your scantron sheet in the white area on the back in ink.
- B. Write and code in the spaces indicated:
 - 1) Name (last name, first initial, middle initial)
 - 2) UF ID number

- 3) Discussion section number
- C. Under "special codes" code in the test ID numbers 1, 1.
 - 2 3 4 5 6 7 8 9 0
 - 2 3 4 5 6 7 8 9 0
- D. At the top right of your answer sheet, for "Test Form Code" encode A.
 - B C D E
- E. While taking the test, please keep your answer sheet covered or turned over at all times.
- F. This test consists of 10 five-point multiple choice questions, 4 two-point bonus multiple choice questions and four pages of partial credit questions worth 30 points. The time allowed is 90 minutes.
- G. WHEN YOU ARE FINISHED:
 - 1) Before turning in your test check for <u>transcribing errors</u>. Any mistakes you leave in are there to stay.
 - 2) You must turn in your scantron and tear off sheets to your discussion leader. Be prepared to show your picture I.D. with a legible signature.
 - 3) The answers will be posted within one day after the exam in Sakai.

NOTE: Be sure to bubble the answers to questions 1–14 on your scantron.

Part I: 5 points each

1. Given the following numbers:

$$0.15 \; , \; -\frac{\pi}{2} \; , \; \sqrt{-9} \; , \; 0 \; , \; \frac{\sqrt{2}}{3} \; , \; 0.\overline{2}$$

How many numbers above are rational?

- a. 1
- b. 2
- c. 3
- d. 4
- e. 5

2. Which of the following graphs has the symmetry with respect to the $\underline{x-axis}$?

P.
$$x = y^2$$

$$Q. \ y = x^2$$

R.
$$x + y = 0$$

a. P only

- b. Q only
- c. R only

- d. P, Q and R
- e. None

3. If x > 1, simplify $\frac{2|1-x|}{x-1} - |3x-3| = \underline{\hspace{1cm}}$.

a. 1 - x

- b. -1 3x
- c. -1 + 3x

- d. -5 3x
- e. 5 3x

4. Rationalize the denominators and simplify:

$$\frac{2-\sqrt{2}}{\sqrt{3}+1}+\frac{3}{\sqrt{6}}$$

a.
$$\frac{\sqrt{3}}{2} + \frac{1}{2} + \frac{\sqrt{6}}{4} - \frac{\sqrt{2}}{4}$$
 b. $\frac{\sqrt{3}}{2} + \frac{1}{2} - \frac{\sqrt{6}}{4} - \frac{\sqrt{2}}{4}$

b.
$$\frac{\sqrt{3}}{2} + \frac{1}{2} - \frac{\sqrt{6}}{4} - \frac{\sqrt{2}}{4}$$

c.
$$\sqrt{3} - 1 - \sqrt{6} + \frac{\sqrt{2}}{2}$$

d.
$$\sqrt{3} - 1 + \frac{\sqrt{2}}{2}$$

- e. None of the above
- 5. If $f(x) = \frac{1}{x-1}$, simplify the quotient $\frac{f(x) f(3)}{x-3}$.

a.
$$\frac{-1}{2(x-3)}$$

b.
$$\frac{1}{2(x-3)}$$

d.
$$\frac{-1}{2(x-1)}$$

e.
$$\frac{1}{2(x-1)}$$

6. Solve: $\frac{-6}{x^2-2x}-\frac{x}{2-x}=\frac{2}{x}$

a.
$$\{-1 \pm 2\sqrt{3}\}$$

b.
$$\{1 \pm 2\sqrt{3}\}$$

d.
$$\{-1 \pm \sqrt{3}\}$$

e.
$$\{1 \pm \sqrt{3}\}$$

7. Find the domain of the expression: $\sqrt{|3-2x|-5}$

a.
$$[-4, 1]$$

a.
$$[-4,1]$$
 b. $(-\infty, -1] \cup [4, \infty)$

c.
$$(-\infty, \infty)$$

d.
$$[-1, 4]$$

e.
$$(-\infty, -4] \cup [1, \infty)$$

8. Simplify:
$$\frac{5^{2+\sqrt{2}}}{(4+\sqrt{11})^{\sqrt{2}}\cdot(4-\sqrt{11})^{\sqrt{2}}}$$

a.
$$\frac{1}{25}$$
 b. $\frac{1}{5}$ c. 1

b.
$$\frac{1}{5}$$

9. Find the equation of the circle if the endpoints of the diameter are (-8,1)and (2,7).

a.
$$(x-3)^2 + (y+4)^2 = 10$$
 b. $(x-3)^2 + (y+4)^2 = 34$

b.
$$(x-3)^2 + (y+4)^2 = 34$$

c.
$$(x+3)^2 + (y-4)^2 = \sqrt{34}$$
 d. $(x+3)^2 + (y-4)^2 = 34$

d.
$$(x+3)^2 + (y-4)^2 = 34$$

e.
$$(x+3)^2 + (y-4)^2 = 10$$

10. Find the <u>sum</u> of all the zeros of the function

$$f(x) = \sqrt{x+3} - 3\sqrt{x} + 1.$$

a.
$$\frac{17}{16}$$
 b. $\frac{1}{16}$ c. 1 d. $\frac{1}{4}$ e. $\frac{5}{4}$

b.
$$\frac{1}{16}$$

d.
$$\frac{1}{4}$$

Continue on the next page!

Part II: Bonus section, 2 points each

Assume that $x \neq 0$ and $y \neq 0$.

11.
$$\frac{2}{x} + \frac{x}{5} =$$

- a. $\frac{x+2}{x+5}$ b. $\frac{2}{5}$ c. $\frac{x^2+10}{5x}$

12.
$$\sqrt{x^2} + \sqrt{5} =$$

- a. $\sqrt{x^2 + 5}$
- b. $|x| + \sqrt{5}$
- c. $x+\sqrt{5}$

13.
$$\frac{1}{x^{-1}+y} =$$

- a. $\frac{x}{y}$ b. $\frac{x}{1+xy}$ c. $\frac{x}{1+y}$

14.
$$(-3x^2)^3 =$$

- a. $-27x^6$
- b. $-27x^5$ c. $-3x^5$


This is the end of the multiple choice section.

Be sure to work on the free response problems on the next four pages.

MAC 1147 TEST IA PART II SPRING 2012

Sect #	Name
UF ID	Signature
SHOW ALL	WORK TO RECEIVE FULL CREDIT.
If the cost to make a	d doughnut store sells a dozen of doughnuts for \$5.50. a dozen of doughnuts is \$2.00 plus a daily fixed cost at the store sells x dozen of doughnuts daily.
a) Find the daily re	venue function.
	Revenue =
b) Find the daily to	tal cost.
	*
The state of the s	$\operatorname{Total} \operatorname{cost} =$
c) Find the daily pr	
	e a ver ez treje je
	$Profit = \underline{}$
	ns of doughnuts must the store sell in order to make of \$160 each day?
	m \

2. a) Find the equation of the line which is parallel to the x-axis and passes through the point (-1,3).



c) Find the slope of the line 2x - 3y = 5.

$$slope = \underline{\hspace{1cm}}$$

1, 5

d) Find the slope of the line that is perpendicular to the line 2x - 3y = 5.

Salah Albah Salah Basar Ba

e) Find the equation of the line that is perpendicular to 2x - 3y = 5 and passes though the point (-1,3).

y =_____

- 3. Consider the quadratic equation $2x^2 2x 3 = 0$.
 - a) Solve the equation using the completing the square.

$$x =$$

b) Solve the equation using the $\underline{\text{quadratic formula}}$.

$$x =$$

c) Find the product of the two zeros of the equation.

4. Simplify the expression and write your answer without negative exponents.

a)
$$\frac{x^2(x^2+4)^{-1/2}-(x^2+4)^{1/2}}{x^2+4}$$

b) $\frac{x^2y^{-1} + x^{-1}y^2}{y^{-1} + x^{-1}}$