

PMATH 965: Mirror Symmetry for GIT Quotients

Content by Elana Kalashnikov
Typeset by Kaleb Ruscitti
Last updated: November 13, 2023

1 Motivation

Mirror symmetry is an enormous area of research. Here we provide motivation from just one perspective, which is Fano classification. Let X be a smooth n -dimensional algebraic variety over \mathbb{C} . Suppose we also have a line bundle $L \rightarrow X$, with some global sections $s_1, s_2, \dots, s_m \in \Gamma(X, L)$. Then we can try and write down a map

$$\begin{aligned}\iota : X &\rightarrow \mathbb{P}^{m-1} \\ x &\rightarrow [s_1(x) : s_2(x) : \dots : s_m(x)].\end{aligned}$$

This is well defined as long as there is no $x \in X$ where all the sections vanish; $s_i(x) = 0$ for all $i = 1, \dots, m$.

Definition 1.1. A line bundle L over an algebraic variety X is called *very ample* if there exist some global sections s_1, \dots, s_m of L for which the map ι defined above is an embedding of X into \mathbb{P}^{m-1} .

If there exists a natural number k such that $L^{\otimes k}$ is very ample, then we say L is *ample*.

For example, the line bundles $\mathcal{O}(n) \rightarrow \mathbb{P}^{m-1}$ (not to be confused with orthogonal groups!) are very ample for all $n \geq 1$.

Definition 1.2. The variety X is *Fano* if $-K_X := \bigwedge^n TX$ is *ample*.

If X is Fano, then it is projective, since $\bigwedge^n TX$ is very ample, meaning it has some sections which define an embedding ι of X into projective space. Some examples of Fano varieties include \mathbb{P}^n , any degree d projective curve in \mathbb{P}^n with $d < n + 1$, and Grassmannians. Naturally then we can ask *Why study Fano varieties?*

- Fano varieties are often the ambient spaces in algebraic geometry. For example, Calabi-Yaus can be cut-out from Fano varieties.
- Fano varieties are special in that there are only finitely many of them in any given dimension.

Theorem 1.3 (Kollár-Miyaoka-Mori). *Up to deformation, there are finitely many Fano varieties in each dimension.*

Here X_1 and X_2 are considered equivalent up to deformation if there exists a flat family $\mathcal{X} \rightarrow B$ over an irreducible base B such that X_1 and X_2 are fibers over some points $b_1, b_2 \in B$.

This raises the big question: Can we classify the Fano varieties? The current progress is:

- In dimension 1, there is just one Fano; \mathbb{P}^1 .
- In dimension 2, there are 10, called the del Pezzo surfaces.
- In dimension 3, there are 105, which were classified throughout the 70s and 90s.
- All higher dimensions are yet to be classified.

In this course, we are also concerned with *mirror symmetries* for Fano varieties. A conjectured mirror symmetry is between n -dimensional Fano varieties and Laurent polynomials in n -dimensions up to an equivalence called *mutation*. Loosely, we can say

Definition 1.4. A variety X is *mirror* to a polynomial f , if you can determine *enumerative info* about X from f .

By enumerative info for X , we mean things like Gromov-Witten invariants, quantum cohomology and quantum periods. The mirror symmetry conjecture is that these can be computed in terms of corresponding quantities of f . For example:

Definition 1.5. Let $f \in \mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ be a Laurent polynomial. The *classical period* of f is the quantity

$$\begin{aligned}\pi_f(t) &= \int_{(S^1)^n} \frac{1}{1 - tf} \frac{dx_1}{x_1} \cdots \frac{dx_n}{x_n} \\ &= \sum_{k=0}^{\infty} \frac{c_1(f^k)}{k!} t^k.\end{aligned}$$

where $c_1(f^k)$ means the coefficient of the constant term of f^k .

The idea therefore, is that mirror symmetry can help us compute hard things in geometry by using easier polynomial data. Then what is the status of this conjecture?

- Established in dimensions 1 and 2 by checking all cases.
- In dimension 3, all Fanos X have a mirror polynomial f , but the classification of other maximally mutable polynomials f is unknown.
- In all dimensions, the symmetry is established for *toric varieties*.

Toric varieties are Fano varieties which have the form $V // T$ where V is a (complex) vector space and $T = (\mathbb{C}^*)^k$, which is called the *algebraic torus*. The double slash $//$ indicates a *geometric invariant theory* quotient, which will be discussed in the first part of the course. Recently, there is a lot of work on extending mirror symmetry to GIT quotients $V // G$ more generally, for G a reductive algebraic group.

As we will see, toric varieties are very nice to work with. This is because they are extremely computable. Essentially, there is a dictionary between the geometry of a toric variety X and the combinatorics of a polytope P corresponding to X . The basic question then, is can a similar correspondence be generalised to other Fano varieties? What should play the role of the polytope? Mirror symmetry answers this question: X corresponds to f , which has a *Newton polytope* with some additional coefficient data.

Exercise 1.6. 1. Show that a degree d hypersurface in \mathbb{P}^n is Fano for $d < n + 1$.

2. Find a closed formula for the classical period of $f(x, y) = x + y + \frac{1}{xy}$, and find a differential equation that it satisfies.

2 Quotients in Algebraic Geometry

Definition 2.1. An *algebraic* group is a group which is also an algebraic variety. An *action* of an algebraic group G on a variety X is a morphism

$$\begin{aligned} G \times X &\rightarrow X \\ (g, x) &\rightarrow g \cdot x \end{aligned}$$

such that for all $g, g' \in G$ and $x \in X$, we have $(gg') \cdot x = g \cdot (g' \cdot x)$ and $e \cdot x = x$.

For example: \mathbb{C}^* , $GL(n)$ and $SL(n)$.

Definition 2.2. Given an action of G on X and some $x \in X$, the *orbit* of x is

$$G \cdot x = \{g \cdot x, \mid g \in G\}. \quad (1)$$

The *stabiliser* of x is

$$G_x = \{g \in G \text{ s.t. } g \cdot x = x\}. \quad (2)$$

Note that G_x is a closed subgroup of G .

Example: Let $T = \mathbb{C}^*$ and $V = \mathbb{C}^2$. Define an action of T on V by $\lambda \cdot (z_1, z_2) = (\lambda z_1, \lambda z_2)$ for all $\lambda \in T$, $(z_1, z_2) \in V$. Then the orbit of $(z_1, z_2) \neq (0, 0)$ is the line through (z_1, z_2) , except the origin. The stabilizer is $\{1\}$. If $(z_1, z_2) = (0, 0)$ then the orbit is $\{(0, 0)\}$ and the stabilizer is all of T . Notice that $(0, 0)$ is in the closure of $G \cdot z$ for all $z \in \mathbb{C}^2$.

Proposition 2.3. For any G , X and $x \in X$,

- The orbit $G \cdot x$ is locally closed and a smooth subvariety of X .
- Each of its irreducible components has dimension $\dim(G) - \dim(G_x)$.
- The closure of $G \cdot x$ is a union of $G \cdot x$ and orbits of strictly smaller dimension.

The last point implies that minimal dimension orbits must be closed, and $\overline{G \cdot x}$ always contains a closed orbit.

Definition 2.4. The action of a group G is called *closed* if every orbit of G is closed.

Definition 2.5. A *linear algebraic group* is a closed subgroup of $GL(n)$.

Goal: If we have an action $G \curvearrowright X$, we want to build some quotient X/G in an algebraic-geometric way. As a naive attempt we can just take the quotient as topological spaces. Consider the action of \mathbb{C}^* on \mathbb{C}^2 from before. If we endow the set $\mathbb{C}^2/\mathbb{C}^*$ with the quotient topology, then since $[(0, 0)]$ is in every open neighbourhood of every other point (as we can always take a sequence of $\lambda_i \in \mathbb{C}^*$ approaching zero), this quotient is not even Hausdorff.

To solve this, we essentially want to delete the origin, and obtain $(\mathbb{C}^2 - \{(0, 0)\})/\mathbb{C}^* = \mathbb{P}^1$. The putative quotient Y we want to define could have the following desirable properties.

1. There exists a surjection $p : X \rightarrow Y$ which is G -invariant.
2. Y is separated.
3. Y satisfies the following universal property: if $f : X \rightarrow Z$ is G invariant, then it factors uniquely through p . That is:

$$\begin{array}{ccc} X & \xrightarrow{p} & Y \\ & \searrow f & \downarrow \\ & & Z \end{array}$$

4. For all U open, $\mathcal{O}_Y(U) \cong \mathcal{O}_X(p^{-1}(U))^G$, where the superscript denotes G -invariant functions.
5. If $Z \subset X$ is closed and G -invariant, then $p(Z)$ is closed. If Z_1, Z_2 are disjoint and closed then $p(Z_1)$ and $p(Z_2)$ are disjoint.

Definition 2.6. If a map p exists as in property 1, and it satisfies property 3, we say it is a *categorical quotient*. If p satisfies properties 4 and 5, then we say it is a *good quotient*.

We can also consider a geometric quotient, which is a good quotient whose points are orbits of $G \curvearrowright X$.

Remark 2.7. The properties of being good or geometric are local on the base, meaning that $p : X \rightarrow Y$ is good or geometric if and only if there exists an open cover of Y with the restrictions of p being good or geometric.

Lemma 2.8. *If $p : X \rightarrow Y$ is good, then it is categorical.*

Proof. Suppose $g : X \rightarrow Z$ is another H invariant morphism and $p : X \rightarrow Y$ is good. Then we want to define $h : Y \rightarrow Z$ such that $p \circ h = g$. Consider $g(p^{-1}(y))$ for some $y \in Y$, which we claim is a singleton set. Suppose for contradiction that there are $z_1 \neq z_2 \in g(p^{-1}(y))$. Then $g^{-1}(z_1) \cap g^{-1}(z_2) = \emptyset$, and these are closed, G -invariant sets because g is continuous and G -invariant. Hence, by the hypothesis that p is good we have:

$$p(g^{-1}(z_1)) \cap p(g^{-1}(z_2)) = \emptyset. \quad (3)$$

However, we must also have that $y \in p(g^{-1}(z_i)), i = 1, 2$ because $z_i \in g(p^{-1}(y))$; hence we have a contradiction and must have that $g(p^{-1}(y))$ is a singleton.

Therefore, we can define a map $h : Y \rightarrow Z$ by $y \rightarrow g(p^{-1}(y))$ and it is well-defined and clearly $p \circ h = g$. It remains to show that this is a morphism of schemes (namely we need it to be locally induced by ring morphisms $\mathcal{O}_X \rightarrow \mathcal{O}_Y$). Let $\{U_i\}$ be a finite open affine cover of Z . Let $W_i = X - g^{-1}(U_i) = g(U_i)^c$. Then U_i being open implies that $g^{-1}(U_i)$ is open and hence W_i is closed. Similarly, g being G -invariant implies W_i is also. Finally, since U_i is a cover, $\bigcap W_i = \emptyset$. Thus by goodness of p we have that $p(W_i)$ are all closed and

$$\bigcap p(W_i) = \emptyset. \quad (4)$$

Define $V_i = Y - p(W_i) = p(W_i)^c$. Then the V_i are an open cover of Y by equation 4. Note further that $p^{-1}(V_i) \subset g^{-1}(U_i)$. Thus we have a sequence of maps

$$\mathcal{O}_Z(U_i) \xrightarrow{\mathcal{O}}_X (g^{-1}(U_i))^G \xrightarrow{\text{res}|_{p^{-1}(V_i)}} \mathcal{O}_X(p^{-1}(V_i))^G \cong_{p \text{ good}} \mathcal{O}_Y(V_i). \quad (5)$$

The G -invariance on the second ring comes from the invariance of g . The last isomorphism is one of the hypothesis conditions of p being good. Thus since U_i and V_i are affine, this defines a local morphism $h_i : V_i \rightarrow U_i$, and it suffices to verify that $h|_{V_i} = h$. \square

Proposition 2.9. *Let $p : X \rightarrow Y$ be a good quotient. Then*

1. $\overline{G \cdot x_1} \cap \overline{G \cdot x_2} \neq \emptyset \iff p(x_1) = p(x_2)$.
2. For all $y \in Y$, there exists a unique closed orbit in $p^{-1}(y)$.

Proof. 1) Suppose $\overline{G \cdot x_1} \cap \overline{G \cdot x_2} \neq \emptyset$. Since p is continuous and constant on orbits, it is constant on orbit closures and hence $p(\overline{G \cdot x_1}) = p(\overline{G \cdot x_2})$ and in particular $p(x_1) = p(x_2)$. On the other hand if $\overline{G \cdot x_1} \cap \overline{G \cdot x_2} = \emptyset$, then since p is good, the image of each orbit is disjoint.

2) Suppose there exist two closed orbits in $p^{-1}(y)$. If they are not equal, then again since p is continuous and constant on orbits they must be disjoint and hence their image is disjoint by goodness. However they must each contain y so this is a contradiction. (Existence was not given in class, maybe it is obvious?) \square

Now, let us construct good quotients for affine varieties.

2.1 Affine GIT Quotient

Suppose we have the action of a group G on an affine variety X . Then $X = \text{Spec}(\mathcal{O}_X)$ by definition, and we want our quotient to be good; so we want it to have functions \mathcal{O}_X^G . Thus the idea is to let our quotient be $\text{Spec}(\mathcal{O}_X^G)$. However \mathcal{O}_X^G is not finitely generated in general, so we will restrict ourselves to *reductive* groups.

Definition 2.10. Let G be a linear algebraic group.

- We say G is *reductive* if every smooth connected unipotent normal subgroup of G is trivial.
- We say G is *linearly reductive* if, for every representation $G \rightarrow GL(V)$ and every non-zero fixed point $v \in V$, there exists a homogeneous G -invariant degree-1 polynomial f on V such that $f(v) \neq 0$.
- We say G is *geometrically reductive* if, for every representation $G \rightarrow GL(V)$ and every non-zero fixed point $v \in V$, there exists a homogeneous G -invariant polynomial f on V such that $f(v) \neq 0$.

Theorem 2.11. The three properties above are equivalent over \mathbb{C} .

For example, $GL(n, \mathbb{C})$, $SL(n, \mathbb{C})$ and $PGL(n, \mathbb{C})$ are reductive.

Theorem 2.12 (Nagata's Theorem). Let G be geometrically reductive acting on a finitely generated \mathbb{C} -algebra R . Then R^G is finitely generated.

Lemma 2.13. Let G be a geometrically reductive group acting on an affine variety X . Let Z_1 and Z_2 be two closed, G -invariant disjoint subsets of X . Then there exists a G -invariant function $\psi \in \mathcal{O}_X^G$ such that $\psi(Z_1) = 1$ and $\psi(Z_2) = 0$.

Proof. Firstly

$$\langle 1 \rangle = I(\emptyset) = I(Z_1 \cap Z_2) = I(Z_1) + I(Z_2), \quad (6)$$

therefore $1 = f_1 + f_2$ for some f_1, f_2 with $f_i(Z_j) = \delta_{ij}$. Claim: (c.f. Hoskins) The subspace spanned by $\{g \cdot f, \mid g \in G\} \subset \mathcal{O}_X$ is G -invariant and finite dimensional. Therefore we can pick a basis h_1, \dots, h_n , and because G acts on all of the h_i , we get an induced action of G on \mathbb{C}^n such that the map

$$\begin{aligned} \phi : X &\rightarrow \mathbb{C}^n \\ x &\rightarrow (h_i(x)) \end{aligned}$$

is G -equivariant, meaning $\phi(g \cdot x) = g\phi(x)$. Note then that $\phi(Z_1) = 0$ and $\phi(Z_2) \neq 0$, and define $v = \phi(Z_2) \in \mathbb{C}^n$.

Since ϕ is G -equivariant, v is fixed by the action of G . Then by the hypothesis of geometric reductivity, there exists some G -invariant homogenous f_0 such that $f_0(v) \neq 0$ and $f_0(0) = 0$. Finally, let

$$\psi = \frac{1}{f_0(v)} f_0 \circ \phi. \quad (7)$$

□

Definition 2.14. The *affine GIT quotient* of an affine variety X under reductive group G , denoted $X // G$ is $\text{Spec}(\mathcal{O}_X^G)$.

Theorem 2.15. Let X be an affine variety and G a reductive group acting on X . Then $p : X \rightarrow Y = \text{Spec}(\mathcal{O}_X^G)$ is a good quotient.

Proof. First we show p is G -invariant. Suppose for contradiction there exist $x \in X$, $g \in G$ such that $p(x) \neq p(g \cdot x)$. Since Y is affine, there exists an $x \in \mathcal{O}_Y$ such that $f(x) \neq f(g \cdot x)$. However $\mathcal{O}_Y = \mathcal{O}_X^G$ by definition, so f must be G invariant, giving a contradiction.

Next we show p is surjective. Let $y \in Y$ and let $\langle f_1, \dots, f_n \rangle$ be the ideal defining y . Let \mathfrak{m} be the maximal ideal containing $\langle f_1, \dots, f_n \rangle$. The point corresponding to \mathfrak{m} in X maps to y under p .

Now let $U \subset Y$ be open. We want to show $\mathcal{O}_Y(U) \cong \mathcal{O}_X(p^{-1}(U))^G$; it suffices to show this for $U = D_f^Y$ for any $f \in \mathcal{O}_Y$.

$$\begin{aligned}\mathcal{O}_Y(D_f^Y) &= (\mathcal{O}_Y)_f \\ &= [\mathcal{O}_X(X)^G]_f \\ &= [\mathcal{O}_X(X)_f]^G \\ &= [\mathcal{O}_X(D_f^X)]^G \\ &= \mathcal{O}_X(p^{-1}(D_f^Y))^G\end{aligned}$$

Let Z_1, Z_2 be G -invariant closed disjoint subsets. By the lemma, there exists $\psi \in \mathcal{O}_X^G$ with $\psi(Z_1) = 0$ and $\psi(Z_2) = 1$. Then $p(Z_1) \cap p(Z_2) = \emptyset$, because there is a G -invariant function which separates them. This turns out to be equivalent to the topological condition for goodness, that $p(Z_1) \cap p(Z_2) = \emptyset$. To prove this, it suffices to prove that if Z is closed and G -invariant then $p(Z)$ is closed.

Suppose Z is closed and G -invariant. For contradiction, suppose there exists $g \in \overline{p(Z)} - p(Z)$. Then Z and $p^{-1}(y)$ are both closed and G -invariant, so

$$\overline{p(Z)} \cap \overline{p^{-1}(y)} = \emptyset. \quad (8)$$

however y must be in this intersection, giving a contradiction. \square

Proposition 2.16. *If the action of G is closed then $X // G$ is a geometric quotient.*

This GIT construction separates orbits as much as possible while still being good.

Example: Consider $\mathbb{C}^* \curvearrowright \mathbb{C}^2$ by $t(x, y) = (tx, t^{-1}y)$. Then the affine GIT quotient is given by $\mathbb{C}^2 // \mathbb{C}^* = \text{Spec}(\mathbb{C}[x, y]^G)$, and $\mathbb{C}[x, y]^G = \mathbb{C}[xy] \cong \mathbb{C}[z]$. Therefore $\mathbb{C}^2 // \mathbb{C}^* = \mathbb{C}$. The quotient map is $(x, y) \rightarrow xy$ and the orbits come in three types:

1. The orbit of the origin is $G \cdot (0, 0) = (0, 0)$.
2. The orbits of $(x, 0)$ and $(0, y)$, for $x, y \neq 0$ are the x and y axes in \mathbb{C}^2 .
3. The remaining orbits have the form $G \cdot (x, y) = \{(z_1, z_2) \mid z_1 z_2 = \lambda\}$ for some λ , which are conics.

The GIT quotient sends the type 1 and 2 orbits to the same point, $0 \in \mathbb{C}$, so this is not a geometric quotient.

Example: Consider the additive complex group $G_a = (\mathbb{C}, +)$. Let it act on \mathbb{C}^4 by embedding it into $GL(4, \mathbb{C})$ by the map

$$s \rightarrow \begin{pmatrix} 1 & s & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & s \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (9)$$

G_a is not reductive, but in this case the ring of invariants is still finitely generated. Note however that in our proof that the GIT quotient is surjective, we again used that G is reductive. In this case, we will not have a good quotient. If f is invariant, it must send

$$\begin{aligned}x_1 &\rightarrow x_2 & x_2 &\rightarrow sx_1 + x_2 \\ x_3 &\rightarrow x_3 & x_4 &\rightarrow sx_3 + x_4\end{aligned}$$

So x_1, x_3 are invariant, and $x_1x_4 - x_2x_3$ is invariant. It turns out these three generate all the invariants, and so $\mathcal{O}_{\mathbb{C}^4}^{G_a} = \mathbb{C}[x_1, x_3, x_1x_4 - x_2x_3]$. Furthermore, $\text{Spec}(\mathbb{C}[x_1, x_3, x_1x_4 - x_2x_3]) = \mathbb{C}^3$. The quotient map is

$$(x_1, x_2, x_3, x_4) \rightarrow (x_1, x_3, x_1x_4 - x_2x_3), \quad (10)$$

which is not surjective as $(0, 0, \lambda)$ is not in its image for $\lambda \neq 0$.

2.2 Projective GIT Quotient

Recall the example of $\mathbb{C}^* \curvearrowright \mathbb{C}^2$ by scaling. We saw that $(0,0)$ is in the closure of every orbit. Hence $\mathbb{C}^2/\mathbb{C}^*$ is just one point. This is the same as saying that all the \mathbb{C}^* invariants in $\mathbb{C}[x_1, x_2]$ are just the constants. Projective GIT will allow us to loosen the definition of G -invariance and get that $\mathbb{C}^2 // \mathbb{C}^* = \mathbb{P}^1$. Recall, if f is homogeneous of degree k then $f(\lambda x, \lambda y) = \lambda^k f(x, y)$, so f is projectively invariant.

Consider $G \curvearrowright X$, with X a projective variety. We can think of X being projective in two ways, either there is an embedding $X \subset \mathbb{P}^k$, or X is equipped with an ample line bundle $L \rightarrow X$. We will swap between these two pictures as convenient. The idea of projective GIT is to replace Spec with Proj. If R is the graded ring with $X = \text{Proj}(R)$, then we want to define $X // G$ to be $\text{Proj}(R^G)$. To make sense of this in the (X, L) perspective, we need a G -action on the sections of the bundle L .

Definition 2.17. Let X be an algebraic variety and $\pi : L \rightarrow X$ a line bundle. Suppose $G \curvearrowright X$ via $\sigma : G \times X \rightarrow X$. Then a G -linearisation of L is a lift of σ to $\bar{\sigma} : G \times L \rightarrow L$ which commutes with σ under the projection π ; $\sigma(g, \pi(s)) = \pi(\bar{\sigma}(g, s))$ for all $s \in \Gamma(X, L)$, and such that the 0 section is invariant.

Remark 2.18. A linearisation defines a linear map between fibres of L , $\bar{\sigma} : L_x \rightarrow L_{g \cdot x}$.

Example: Let $X = \mathbb{C}^n$ and $L = \mathbb{C} \times \mathbb{C}^n$ be the trivial bundle. Then a linearisation of L is a character in $\chi(G)$. If we fix $\theta \in \chi(G)$, then the linearisation of L is

$$g \cdot (a, v) = (\theta(g)a, g \cdot v). \quad (11)$$

This defines an action on the sections of L ; for $U \subset X$ open and $s \in \Gamma(U, L)$, let $(g \cdot s)(x) = \theta(g)s(g^{-1}x)$.

In the other perspective, when $X \subset \mathbb{P}^k$ explicitly, then a linearisation is a way to think of $G \curvearrowright X$ via an embedding $G \hookrightarrow GL(k+1, \mathbb{C}) \curvearrowright \mathbb{P}^k$. In particular, if L is very ample, then $X \hookrightarrow \mathbb{P}(\Gamma(X, L)^\vee) = \mathbb{P}^k$. Then these two notions of linearisation agree. If $X = \text{Proj}(R)$, then a linearisation is an action $G \curvearrowright R$ which preserves the grading.

In any case, we can now define projective GIT.

Definition 2.19. The *projective GIT quotient* of (X, L) by G , with respect to a given linearisation, is

$$X // G = \text{Proj} \left(\bigoplus_{r \geq 0} \Gamma(X, L^r)^G \right) \quad (12)$$

with the quotient map induced by the injection $R^G \hookrightarrow R$.

Example: We construct \mathbb{P}^n as a GIT quotient of $X = \mathbb{C}^{n+1}$ by \mathbb{C}^* under scaling. A linearisation is given by a character of \mathbb{C}^* .

$$\begin{aligned} \chi(\mathbb{C}^*) &\cong \mathbb{Z} \\ (\lambda \rightarrow \lambda^a) &\leftrightarrow a \end{aligned}$$

Let $a \in \mathbb{Z}$ be a character, then \mathbb{C}^* acts on the trivial line bundle L over \mathbb{C}^{n+1} by $\lambda \cdot s(x) = \lambda^a s(x)$. We have that $\Gamma(\mathbb{C}^{n+1}, L^k) = \mathbb{C}[x_0, \dots, x_n]$. If we want an element f to be \mathbb{C}^* invariant, we need

$$t \cdot f(x_0, \dots, x_n) = t^a f(t^{-1}x_0, \dots, t^{-1}x_n) = f(x_0, \dots, x_n). \quad (13)$$

If $a = 1$ then equation 13 exactly means that f is a degree- k homogenous polynomial. Then

$$X // G = \text{Proj} \left(\bigoplus_{k \geq 0} \text{degree } k \text{ homogenous polynomials} \right) = \mathbb{P}^n.$$

If $a = 0$, then equation 13 is only solved by constants. In this case, $X // G$ has only one point and we recover the affine GIT quotient.

If $a < 0$ then equation 13 has no solutions and the quotient is the empty set. Finally, the case with $a > 1 \in \mathbb{N}$ is left as an exercise.

We can also think of \mathbb{C}^{n+1} as $\text{Proj}(\mathbb{C}[x_0, \dots, x_n, y])$, with the grading that lets x_i have degree 0 and y have degree 1. Then $\mathbb{C}^* \curvearrowright \mathbb{C}^{n+1}$ by $\lambda \cdot (x_0, \dots, x_n, y) = (\lambda x_0, \dots, \lambda x_n, \lambda^{-a} y)$ for $a \in \chi(\mathbb{C}^*)$. The quotient in each case works out exactly the same as above.

Let us try to get an intuitive sense for $X // G$. Suppose that L is very ample. Suppose further that some sections s_0, \dots, s_n generate the G -invariant sections in all degrees. Then the Proj construction is essentially doing

$$\begin{aligned} X &\rightarrow \mathbb{P}^n \\ x &\rightarrow [s_0(x) : \dots : s_n(x)]. \end{aligned}$$

This is defined where not all of the $s_i(x)$ vanish; the image is $X // G$, which contains all the points x that have some non-vanishing G -invariant section.

Definition 2.20. A point $x \in X$ is L -semistable for (X, L) if $\{y \in X \mid s(y) \neq 0\}$ is affine and there exists a G -invariant section s of L^r , for some r such that $s(x) \neq 0$.

A point which is not semistable is called unstable. The set of semistable points is denoted $X^{ss}(L)$, it is Zariski open and G -invariant.

If L is ample then $\{y \in X \mid s(y) \neq 0\}$ is always affine.

Definition 2.21. A semistable point $x \in X^{ss}$ is *stable* if there exists some $s \in \Gamma(X, L^k)^G$ such that $s(x) \neq 0$ and the action G on $Y = \{y \in X \mid s(y) \neq 0\}$ is closed, Y is affine and the stabiliser of x is finite. If the stabiliser is not finite, x is called *polystable*.

The set of polystable points is a disjoint union of open sets, each of which consists of polystable orbits of a fixed dimension.

Exercise 2.22. Suppose L is very ample and we have an embedding $X \subset \mathbb{P}^k$ for some k . Show that the following notions of semistable and stable agree with the definitions above.

- $x \in X$ is semi-stable if there exists a G -invariant homogeneous polynomial f with $f(x) \neq 0$.
- $x \in X$ is stable if $G \cdot x$ is finite and there exists a G -invariant homogeneous polynomial and the G -action on D_f is closed.

Theorem 2.23. *There is a G -invariant morphism*

$$p : X^{ss}(L) \rightarrow X // G$$

such that p is a good quotient and $X // G$ is quasi-projective. If L is ample, $X // G$ is projective.

Proof. We prove for L very ample. Write $X = V(I) \subset \mathbb{P}^K$, where $I \subset$ some homogeneous ideal. Then let $R = \mathbb{C}[x_0 : \dots : x_n]/I$ such that $X // G = \text{Proj}(R^G)$. The inclusion $R^G \hookrightarrow R$ induces a rational map $\text{Proj}(R) \rightarrow \text{Proj}(R^G)$, well-defined where points in $\text{Proj}(R)$ don't get mapped into points containing the irrelevant ideal. That is to say, well defined away from the *null cone*

$$N_{RG}(X) := \{x \in X \mid f(x) = 0, \forall f \in R^G\}. \quad (14)$$

Thus the map is well defined on

$$X^{ss} = X - N_{RG}(X) \rightarrow \text{Proj}(R^G).$$

Let $f \in R^G$, let Y_f be the affine open of f in $Y := X // G$. Then X_f is the affine open set in X^{ss} equal to $\text{Spec}((R_f)_0)$, and Y_f is the affine open equal to $\text{Spec}([(R^G)_f]_0)$ and the ring map

$$[(R^G)_f]_0 = [(R_f)_0]^G \hookrightarrow (R_f)_0$$

induces a map $X_f \rightarrow Y_f$. This map is exactly the affine GIT quotient which we proved has the required properties. Since being a good quotient is local on the base, being local on the distinguished affines implies that the quotient must be good everywhere. \square

The next question is to understand when $X // G$ will be a geometric quotient.

Definition 2.24. Let $G \cdot x_1$ and $G \cdot x_2$ be semistable orbits. Then we say that x_1 and x_2 are GIT equivalent if either of the following equivalent things happen:

- $\overline{G \cdot x_1} \cap \overline{G \cdot x_2} \cap X^{ss} = \emptyset$.
- x_1 and x_2 map to the same point in $X // G$.

Proposition 2.25 (c.f. Hoskins). *x is stable if and only if $G \cdot x$ is closed in X^{ss} and G_x is finite.*

Theorem 2.26. *The restriction of $p : X^{ss}(L) \rightarrow X // G$ to $p : X^s(L) \rightarrow X^s(L) // G$ is a geometric quotient.*

2.3 Stability Criteria

We've constructed good and geometric quotients of X^{ss} and X^s , but in general finding the semi-stable and stable points can be difficult. Therefore we will prove some criteria which help us compute these loci. We say f is HGI to mean $f : \mathbb{C}^{n+1} \rightarrow \mathbb{C}$ is a homogeneous, G -invariant polynomial.

Assume G is reductive, $X \subset \mathbb{P}^n$ is a projective variety, and G acts linearly on \mathbb{C}^{n+1} .

Proposition 2.27 (Topological criterion for stability). *Let \hat{x} be a lift of $x \in X$ to \mathbb{C}^{n+1} . Then*

1. x is semistable $\iff 0 \notin G \cdot \hat{x}$.
2. x is stable $\iff \dim G_{\hat{x}} = 0$ and $G \cdot \hat{x}$ is closed in \hat{X} .

Proof. 1) If x is semistable, then there exists f HGI such that $f(x) \neq 0$. Then $f(\hat{x}) \neq 0$, and by G -invariance $f(G \cdot \hat{x}) = f(\hat{x})$ is non-zero constant. Furthermore, by continuity this means $f(\overline{G \cdot \hat{x}}) \neq 0$. Thus $\overline{G \cdot \hat{x}} \cap \{0\} = \emptyset$ by the topological property of good quotients.

Conversely, these sets being disjoint means there exists an $f \in \mathbb{C}[x_0, \dots, x_n]^G$ such that $f(\overline{G \cdot \hat{x}}) = 1$ and $f(0) = 0$ by an earlier lemma. We can write $f = \sum h_i$ with the h_i all HIG, and since f does not vanish on $\overline{G \cdot \hat{x}}$ at least one h_i must not vanish there.

2) Suppose $\dim G_x = 0$ and there is a HIG f such that $x \in D_f$ and $G \cdot \hat{x}$ is closed in D_f . Note that $G_{\hat{x}} \subset G_x$ which implies $G_{\hat{x}}$ is finite. Let $\pi : \mathbb{C}^{n+1} \rightarrow \mathbb{P}^n$ be the quotient and define

$$Z = \{z \in \hat{X} \mid f(z) = f(\hat{x})\}. \quad (15)$$

Z is closed. Consider the map $\pi : Z \rightarrow D_f$, which is surjective and finite because things in $\pi^{-1}(a)$ must be in the stabiliser G_a which is finite. Then $\pi^{-1}(G \cdot x)$ is closed since $G\hat{x}$ is closed in D_f , and π is continuous. $\pi^{-1}(G \cdot x)$ is G -invariant, and so it is the union of a finite number of G orbits, all of dimension $\dim G_x$, hence they are closed, and so is $G \cdot \hat{x} \subset \pi^{-1}(G \cdot x)$.

Conversely, suppose $G \cdot \hat{x}$ is closed in \hat{X} . Then $0 \notin \overline{G \cdot \hat{x}}$ again and by 1), x is semi-stable. Hence there exists f HIG s.t. $f(x) \neq 0$.

Let Z, π be the same as before. Since $\pi(G \cdot \hat{x}) = G \cdot x$, then x has a finite stabiliser and $G \cdot x$ is closed in D_f . This argument works for every HGI f with $f(x) \neq 0$, hence $G \cdot x$ is closed in X^{ss} , and x is stable. \square

Now, we will work towards a numerical criterion, first by considering $G = \mathbb{C}^*$. Let $\mathbb{C}^* \curvearrowright X \subset \mathbb{P}^k$ linearly. Up to change of basis, we can assume \mathbb{C}^* acts diagonally on \mathbb{C}^{k+1} . To be precise, for $t \in \mathbb{C}^*$ we have

$$t \cdot (x_0, \dots, x_k) = (t^{w_0} x_0, \dots, t^{w_k} x_k), \quad w_i \in \mathbb{Z}. \quad (16)$$

For $x = [x_0 : \dots : x_n] \in X$, let $\hat{x} = (x_0, \dots, x_n)$, and let $\mu(x) = \max\{-w_i \mid i \text{ such that } x_i \neq 0\}$. Consider

$$\lim_{t \rightarrow 0} t^s(t \cdot \hat{x}) = \lim_{t \rightarrow 0} (t^{s+w_0} x_0, \dots, t^{s+w_k} x_k). \quad (17)$$

If $s > \mu(x)$, then the limit goes to 0. If $s < \mu(x)$ then the limit doesn't exist. Thus, $\mu(x)$ is the unique $s \in \mathbb{Z}$ such that this limit exists but is non-zero. Similarly, let $\mu^-(x) = \max\{w_i \mid i \text{ such that } x_i \neq 0\}$. Then

1. $\mu^-(x) < 0 \iff \lim_{t \rightarrow \infty} t \cdot \hat{x}$ does not exist.
2. $\mu^-(x) = 0 \iff \lim_{t \rightarrow \infty} t \cdot \hat{x}$ exists and is non-zero.

Using μ and μ^- we can find the following stability criterion.

Proposition 2.28. *x is (semi)-stable if and only if $\mu(x) > (\geq) 0$ and $\mu^-(x) > (\geq) 0$.*

Proof. The closure of $G \cdot \hat{x}$ is obtained by adding in the limits as $t \rightarrow 0$ and $t \rightarrow \infty$;

$$\overline{G \cdot \hat{x}} = G \cdot \hat{x} \cup \{\lim_{t \rightarrow 0} t \cdot \hat{x}, \lim_{t \rightarrow \infty} t \cdot \hat{x}\}. \quad (18)$$

From the topological criterion, we know semistability means $0 \notin \overline{G \cdot \hat{x}}$. This happens exactly when neither limit is zero, which happens if and only if $\mu(x) \geq 0$ and $\mu^-(x) \geq 0$, as discussed above.

Furthermore stability occurs $G \cdot \hat{x}$ is closed, namely $\overline{G \cdot \hat{x}} = G \cdot \hat{x}$. This happens when both limits do not exist, which is if and only if $\mu(x) > 0$ and $\mu^-(x) > 0$. \square

Now we will use this to build a criterion for general reductive G , called the *Hilbert-Mumford Numerical Criterion*. Let G be reductive acting on X projective via $\rho : G \hookrightarrow GL(n, \mathbb{C})$.

Definition 2.29. A *one-parameter subgroup* (1PS) of G is a non-trivial group homomorphism $\lambda : \mathbb{C}^* \rightarrow G$.

Let λ be a 1PS of G , $x \in X$ and \hat{x} a lift of x as before. Then there is an action $\mathbb{C}^* \curvearrowright X$ by $\mathbb{C}^* \xrightarrow{\lambda} G \xrightarrow{\rho} GL(n, \mathbb{C})$. If we write $x = \sum x_i e_i$ in a diagonal basis $\{e_i\}_{i=0}^k$ for this action, then as before $t \in \mathbb{C}^*$ acts by

$$t \cdot (x_0, \dots, x_k) = (t^{w_0} x_0, \dots, t^{w_k} x_k), \quad w_i \in \mathbb{Z}. \quad (19)$$

Define $\mu(x, \lambda) = -\min\{w_i \mid i \text{ such that } w_i \neq 0\}$.

Exercise 2.30. Prove that

1. $\mu(x, \lambda^n) = n\mu(x, \lambda)$,
2. $\mu(g \cdot x, g\lambda g^{-1}) = \mu(x, \lambda)$,
3. $\mu(x, \lambda) = \mu(x_0, \lambda)$, $x_0 := \lim_{t \rightarrow 0} \lambda(t) \cdot x$.

Note that we do not need a μ^- because $\lim_{t \rightarrow \infty} \lambda(t) \cdot \hat{x} = \lim_{t \rightarrow 0} \lambda^{-1}(t) \cdot x$.

Lemma 2.31. x is (semi)-stable with respect to $\lambda(\mathbb{C}^*)$ if and only if $\mu(x, \lambda) > (\geq) 0$ and $\mu(x, \lambda^{-1}) > (\geq) 0$.

Proof. Exactly as in the \mathbb{C}^* case. □

Theorem 2.32 (Hilbert-Mumford Numerical Criterion). *Let G be a reductive group, acting linearly on $X \subset \mathbb{P}^n$. Then x is (semi)-stable if and only if $\mu(x, \lambda) > (\geq) 0$ for all 1PS λ of G .*

We won't prove this, but the work we've done so far shows that this theorem is equivalent to:

Theorem 2.33 (Fundamental Theorem of GIT). *Let G be a reductive group acting linearly on \mathbb{C}^{n+1} , and let $x \in \mathbb{C}^{n+1}$. If $y \in G \cdot x$ then there is a 1PS λ of G such that $\lim_{t \rightarrow 0} \lambda(t)x = y$.*

Exercise 2.34. Let $G = \mathbb{C}^*$, $X = \mathbb{P}^n$. Consider the action given by weights $(-1, \dots, -1, 1, \dots, 1)$, k and then $n - k$ times. Determine, using the Hilbert-Mumford criterion, the GIT quotient $\mathbb{P}^n // \mathbb{C}^*$.

Next we will rephrase this criterion for GIT quotients defined in terms of an ample line bundle $L \rightarrow X$. Suppose L has a G linearisation and let λ be a 1PS of G . Then since X is projective,

$$x_0 := \lim_{t \rightarrow 0} \lambda(t)x \in X \quad (20)$$

is a fixed point for λ . Then λ acts on the fibre L_{x_0} by some character $t \rightarrow t^r$ and we define $\mu^L(x, \lambda) = r$. To compare this definition with the previous one, choose a basis such that λ acts diagonally with weights w_0, \dots, w_k , and write $\hat{x} = [a_0 : \dots : a_k]$. Then

$$\lim_{t \rightarrow 0} \lambda(t) = [b_0 : \dots : b_n], \quad b_i = \begin{cases} a_i & \text{if } w_i = -\mu(x, \lambda), \\ 0 & \text{else.} \end{cases} \quad (21)$$

On the fibres, the action of λ has weight $-\mu(x, \lambda)$. These fibres lift to $\mathcal{O}(-1)$ so the weight of λ on $\mathcal{O}(1)$ is $\mu(x, \lambda)$. Using the Hilbert-Mumford criterion, we obtain

Theorem 2.35. *Let G be reductive, $G \curvearrowright X$ projective, and $L \rightarrow X$ ample and with a linearisation. Then x is (semi)-stable if and only if $\mu^L(x, \lambda) > (\geq) 0$ for all 1PS λ of G .*

Proof. L^n is very ample, and $\mu^{L^n}(x, \lambda) = \mu^L(x, \lambda)$. Since we are only checking for μ non-zero, we can assume L is very ample without loss of generality, embed X into \mathbb{P}^n and then reduce to the Hilbert-Mumford criterion. □

There is one other important case. If $X = \mathbb{C}^n$ then the criterion does not apply as written as \mathbb{C}^n is not projective. However, projectivity was only used to look at the limit $\lim_{t \rightarrow 0} \lambda(t) \cdot x$. Now for \mathbb{C}^n this limit may not exist, but if it does not exist then it cannot add anything to the closure of $G \cdot x$, and that is exactly what we want for stability. Thus we have

Theorem 2.36. *A point $x \in \mathbb{C}^n$ is (semi)-stable if and only if $\mu(x, \lambda) > (\geq) 0$ for all 1PS λ of G for which $\lim_{t \rightarrow 0} \lambda(t) \cdot x$ exists.*

Example - Grassmannian: Let $0 < r < n \in \mathbb{N}$ and let $GL(r) \curvearrowright M_{r \times n}$ by left multiplication. Let $L = M_{r \times n} \times \mathbb{C}$ be the trivial line bundle and let it be linearized by $g \rightarrow \det(g)$. First we claim that :

$$A \in M_{r \times n} \text{ is stable} \iff A \text{ is stable} \iff \text{rk}(A) = r.$$

Suppose that $\text{rk}(A) < r$. Since stability is G -invariant, we can replace A with gA and thus row reduce to A in row-reduced echelon form; in particular since the rank is not full, it will have a bottom row of zeros. Let λ be the 1PS defined by

$$\lambda(t) = \text{diag}(t, t, \dots, t, t^{-r}),$$

so that the limit $\lim_{t \rightarrow 0} \lambda(t)A$ exists, and $\langle \theta, \lambda \rangle = r - 1 - r = -1 < 0$, implying A is unstable.

Now suppose that $\text{rk}(A) = r$. Since $\mu_L(gx, g\lambda g^{-1}) = \mu_L(x, \lambda)$, we can assume that any λ is diagonal. Let $\lambda(t)$ be the 1PS given by weights w_1, \dots, w_r . Since every row of A is non-zero, $\lim_{t \rightarrow 0} \lambda(t)A$ exists and hence $w_i \geq 0$ unless $\langle \theta, \lambda \rangle = \sum w_i > 0$. Hence A is stable.

Using this claim, we have that the GIT quotient $M_{r \times n} // GL(r)$ is classes of full-rank matrices up to change of basis, which is the Grassmannian $\text{Gr}(n, r)$.

2.4 Symplectic Reduction

In this section, our goal is to connect GIT quotients to another form of geometric quotient called symplectic reduction.

Definition 2.37. A *symplectic manifold* is a pair (X, ω) where X is a real manifold, and ω is a closed, non-degenerate smooth 2-form on X , called the symplectic form.

In detail, ω is a skew-symmetric bilinear form $\omega_x : T_x X \times T_x X \rightarrow \mathbb{R}$ such that

- (Smooth) ω_x varies smoothly in x .
- (Non-degenerate) For all $x \in X$, ω_x is an isomorphism of $T_x X$ with $T_x^* X$ via

$$\xi \rightarrow \omega_x(\xi, -)$$

- (Closed) $d\omega = 0$.

Example: Let $X = \mathbb{C}^n$ with co-ordinates $z_k = x_k + iy_k$. Then

$$\omega = \sum_{k=1}^n dy_k \wedge dx_k = \frac{1}{2i} \sum dz_k \wedge d\bar{z}_k$$

is a symplectic form.

Example: Let $X = \mathbb{P}^n$, which is given by the quotient $\mathbb{C}^{n+1}/\mathbb{C}^*$. The symplectic form on \mathbb{C}^{n+1} is not invariant under \mathbb{C}^* , so it does not pass to the quotient. However, if we think of $\mathbb{P}^n = S^{2n+1}/S^1$, and restrict the form on \mathbb{C}^{n+1} to S^{2n+1} , we get a short exact sequence:

$$T_p(S^1 \cdot p) \hookrightarrow T_p S^{2n+1} \twoheadrightarrow T_{[p]} \mathbb{P}^n. \quad (22)$$

From this we define the *Fubini-Study* form ω_{FS} , by $\pi^* \omega_{FS} = \omega_{S^{2n+1}}$, where $\pi : S^{2n+1} \rightarrow \mathbb{P}^n$ is the projection. For this to be well defined, one must check that it vanishes on $T_p(S^1 \cdot p)$.

For the remainder of this section, fix (X, ω) to be a symplectic manifold.

Definition 2.38. Let $H : X \rightarrow \mathbb{R}$ be smooth. Then $dH \in \Omega^1(X)$. The *Hamiltonian vector field* corresponding to H is the unique vector field ξ such that

$$\iota_\xi \omega = dH. \quad (23)$$

We say a vector field ξ is *Hamiltonian* if $\iota_\xi \omega$ is exact.

Definition 2.39. A vector field ξ is *symplectic* if $\iota_\xi \omega$ is closed

Note then that being Hamiltonian implies being symplectic.

Definition 2.40. Let K be a compact, connected Lie group acting on (X, ω) . If $(g \cdot -) = l_g : X \rightarrow X$ is a symplectomorphism for all $g \in K$, then we say K *acts symplectically*.

Suppose $K \curvearrowright (X, \omega)$ symplectically. Let $\text{Lie}(K) = \mathfrak{k}$. There exists the exponential map

$$\exp(-, A) : \mathbb{R} \rightarrow X, \quad t \rightarrow \exp(tA) \cdot x. \quad (24)$$

We can take the derivative and evaluate at zero to get

$$\frac{d}{dt} e^{tA} \cdot x|_{t=0} \in T_x X, \quad (25)$$

and letting x vary, this defines a vector field denoted X_A .

Exercise 2.41. Since K acts symplectically, its image is closed. When is X_A Hamiltonian? Can we pick a map $\mathfrak{k} \rightarrow C^\infty(X)$ which is a Lie algebra homomorphism?

Definition 2.42. A *moment map* for a Hamiltonian action is a map $\mu : X \rightarrow \mathfrak{k}^*$ such that

1. μ is \mathfrak{k} -equivariant
2. $d\langle \mu(\xi), A \rangle = \omega(X_A, \xi)$ for all $A \in \mathfrak{k}$

Example: Let $U(n)$ act on \mathbb{C}^n in the natural way. Then $\mathfrak{u}(n)$ is the skew-Hermitian matrices and

$$\omega(z, v) = \frac{1}{2i} \left(H(z, v) - \overline{H(z, v)} \right), \quad (26)$$

where $H(z, v) = zv^\dagger$, defines a symplectic form. We identify $\mathfrak{u}(n) \cong \mathfrak{u}(n)^*$ by the map $A \rightarrow (B \rightarrow \text{Tr}(AB))$, which lets us define

$$\begin{aligned} \mu : X &\rightarrow \mathfrak{u}(n) \\ z &\rightarrow -\frac{i}{2} zz^\dagger \end{aligned}$$

or equivalently,

$$\begin{aligned} \mu : X &\rightarrow \mathfrak{u}(n)^* \\ z &\rightarrow -\frac{i}{2} \text{Tr}(z^\dagger Az). \end{aligned}$$

We show this is equivariant and leave condition 2 as an exercise. Let $K \in U(n)$. Then

$$\begin{aligned} \mu(K \cdot z)A &= -\frac{i}{2} \text{Tr} \left((Kz)^\dagger A(Kz) \right) \\ &= -\frac{i}{2} \text{Tr}(z^\dagger K^\dagger A K z) \\ &= \mu(z)(K \cdot A). \end{aligned}$$

This example allows us to compute the moment map for any unitary linear action on \mathbb{C}^n . If $\rho : K \rightarrow U(n)$, then $\mu_K = \rho^* \mu_{U(n)}$.

Example: If $K = (S^1)^n$ acting on \mathbb{C}^n with weights (r_1, \dots, r_n) , then $\mu : \mathbb{C}^n \rightarrow \mathbb{R}^n$ is given by

$$\mu(z_1, \dots, z_n) = \left(\frac{r_i |z_i|^2}{2} \right)_{i=1}^n.$$

Exercise 2.43. Show that the moment map for $U(r) \curvearrowright M_{r \times n}$ is given by $A \rightarrow i(A^\dagger A - \mathbb{I})$.

Now we have all the pieces to define symplectic reduction. Suppose we have $K \curvearrowright (X, \omega)$ and we want to build a quotient which is also symplectic. Even if X/K is a smooth manifold, it might not have even dimension and so it certainly might not be symplectic. Suppose there is a moment map μ for the K action.

Lemma 2.44. *If η is a regular value of μ , then $\mu^{-1}(\eta)$ is a closed submanifold of X with codimension $\dim K$.*

Moreover, $T_x \mu^{-1}(\eta)$ and $T_x(K \cdot x)$ are orthogonal with respect to the ω for all $x \in \mu^{-1}(\eta)$.

Suppose η is a regular value of μ which is fixed by the coadjoint orbit action. Since μ is K -equivariant, $\mu^{-1}(\eta)$ is K -invariant. Thus we can define the *symplectic reduction* to be

$$X_\eta^{\text{red}} = \mu^{-1}(\eta)/K. \quad (27)$$

Theorem 2.45 (Marsden-Weinstein-Meyer). *Fix a Hamiltonian action of a compact connected Lie group K on a symplectic manifold (X, ω) with moment map μ , and a regular, co-adjoint fixed value $\eta \in \mathfrak{k}^*$. If the action of K on $\mu^{-1}(\eta)$ is free then*

1. X_η^{red} is a smooth manifold of dimension $\dim X - 2 \dim K$.
2. There is a unique symplectic form ω^{red} on X_η^{red} such that

$$\pi^*(\omega^{\text{red}}) = \iota^* \omega$$

where $\iota : \mu^{-1}(\eta) \rightarrow X$ and $\pi : \mu^{-1}(\eta) \rightarrow X_\eta^{\text{red}}$ are the inclusion and quotient maps.

Reductive groups are the complexification of real compact Lie groups: Fix any reductive group G . There is a compact Lie group $K < G$ such that $\mathfrak{k} \subset \mathfrak{g}$ and $\mathfrak{g} = \mathfrak{k} + i\mathfrak{k}$.

Examples: $S^1 < \mathbb{C}^*$, $SU(n) < SL(n, \mathbb{C})$.

Now suppose we have G reductive, $K < G$ compact, and G acts on a smooth projective variety $X \subset \mathbb{P}^n$ linearly. Then this induces an action of K on X which is symplectic on \mathbb{P}^n . The moment-map corresponding to the Fubini-Study form on \mathbb{P}^n gives a moment map for the K action.

Theorem 2.46 (Kempf-Ness). *Let K, G, X, μ be as above. Then*

1. $G \cdot \mu^{-1}(0) = X^{\text{polystable}}$.
2. *If $x \in X$ is polystable, then $G \cdot x \cap \mu^{-1}(0) = K \cdot y$ for some $y \in X$.*
3. $x \in X^{ss} \iff \overline{G \cdot x} \cap \mu^{-1}(0) \neq \emptyset$.

Proof: Hoskins

As a corollary we have

Proposition 2.47. *The inclusion $\mu^{-1}(0) \subset X^{ss}$ induces a homeomorphism $\mu^{-1}(0)/K \rightarrow X // G$*

Proof. $X // G$ is a good quotient, the set X^{ss} / \sim , where \sim denotes GIT equivalence. Then for $x \in X^{ss}$, \overline{Gx} contains a unique closed orbit, which is polystable. Hence as a set

$$X // G = \{\text{polystable orbits}\}$$

Then point 2 of Kempf-Ness tells us the sets are the same and we have a continuous bijection between a compact space and a Hausdorff space; hence a homeomorphism. \square

Example: Let $U(r) \curvearrowright M_{r \times n}$ by multiplication. There is a moment map $\mu : A \rightarrow i(AA^* - \mathbb{I})$, and

$$\mu^{-1}(0) = \{A \mid AA^* = \mathbb{I}\} = \{A \mid A \text{ isometrically embeds } \mathbb{C}^r \rightarrow \mathbb{C}^n\}. \quad (28)$$

Then there is a surjection $V^{ss} = \{A \mid \text{rank } A = r\} \rightarrow \mu^{-1}(0)$.

Proposition 2.48. *If K acts freely on $\mu^{-1}(0)$, then $X^s = X^{ss}$ and $X // G$ is non-singular and a geometric quotient.*

Proof. Acting freely means G_x is trivial for all $x \in \mu^{-1}(0)$. If $x \in X^{ss}$ but not X^s , then $\overline{G \cdot x}$ contains exactly one closed orbit of strictly smaller dimension than $G \cdot x$. By Kempf-Ness, this orbit is in $\mu^{-1}(0)$ and hence has trivial stabilizer. However, this implies that this closed orbit has full dimension, a contradiction. Thus $X^{ss} = X^s$. \square

3 More GIT Quotient Stuff

3.1 Semi-Invariants

Let $\chi(G)$ be the group of characters of G . Recall that $\chi([C^*]^r) = \mathbb{Z}^r$, and $\chi(GL(r)) = \mathbb{Z}$, generated by $g \rightarrow \det g$. Recall further that there exists a pairing $\langle \cdot, \cdot \rangle$ between one-parameter subgroups (1PS) of G and characters. This is a perfect pairing, so we sometimes call 1PS *co-characters*.

Let V be a vector space and G a reductive group acting linearly on V ($G \hookrightarrow GL(V)$). Choose co-ordinates (x_1, \dots, x_n) for V , which gives us a G action on the ring $R = \mathbb{C}[x_1, \dots, x_n]$. In affine GIT, we studied the ring of invariants R^G , and now we will see that in projective GIT we are studying "semi-invariants" of R .

Definition 3.1. An element $f \in \mathbb{C}[x_1, \dots, x_n]$ is called *semi-invariant of weight θ* , for $\theta \in \chi(G)$, if $g \cdot f(x) = \theta(g)f$.

Example: Let $(\mathbb{C}^*)^k \curvearrowright \mathbb{C}^n$ diagonally with weights $\theta_1, \dots, \theta_n \in \mathbb{Z}^k$. Then we have a "multigrading" on $\mathbb{C}[x_1, \dots, x_n]$, where x_i has weight θ_i . In this case, the semi-invariants are the polynomials that are homogeneous with respect to this multigrading.

Example: Let $G = GL(r) \curvearrowright M_{r \times n}$. Then $\chi(G) = \mathbb{Z}$ and we can ask 'what are the semi-invariants of weight 1?' Let

$$A = \begin{bmatrix} [A]_1 & \dots & [A]_n \end{bmatrix}, \quad g \cdot f := \det(g)f. \quad (29)$$

For any $I = i_1, \dots, i_r$, let $P_I = \det [[A]_{i_1}, \dots, [A]_{i_r}]$. Then P_I is a semi-invariant because

$$\begin{aligned} gP_I(A) &= P_I(g \cdot A) = \det [g[A]_{i_1}, \dots, g[A]_{i_r}] \\ &= \det(g)P_I(A). \end{aligned}$$

Theorem 3.2. *There is a correspondence of sets between semi-invariants of weight k , and homogeneous degree- k polynomials in the P_I .*

In this example, the P_I finitely generate the ring of semi-invariants, $R_\theta := \bigoplus_{k \geq 0} \mathbb{C}[x_1, \dots, x_n]^{k\theta}$. However it is not hard to come up with examples where R_θ has no known list of finite generators (despite such generators having to exist by reductivity of G).

3.2 GIT for $V // G$

By definition, we have $V // G = \text{Proj}(R_\theta)$. Thankfully, due to the Hilbert-Mumford criterion, you don't always need to compute generators of R_θ in order to find the GIT quotient. However, these semi-invariants do contain useful information if you *can* compute them.

We can go from a G -linearised vector bundle of V to a vector bundle on $V // G$ under some assumptions. Let E be a representation of G . Then, consider the G -linearized trivial vector bundle $V^{ss} \times E$, and take the quotient by G :

$$\begin{array}{ccc} V^{ss} \times E & \xrightarrow{\quad /G \quad} & (V^{ss} \times E)/G := E_G \\ \downarrow & & \downarrow \\ V^{ss} & & V // G \end{array}$$

Question: When is E_G a vector bundle?

Definition 3.3. A vector bundle $E \rightarrow X$ descends to $F \rightarrow X // G$ if

$$E|_{X^{ss}} = p^* F, \quad (30)$$

where $p : X \rightarrow X // G$ is the GIT quotient.

Theorem 3.4 (Kempf's Descent Lemma). E_G is a vector bundle if and only if $\forall x \in X^{ss}$ such that $G \cdot x$ is closed in X , G_x acts trivially on the fibre E_x .

In particular, if $X^{ss} = X^s$ and the G action is trivial, then all vector bundles descend.

These vector bundles E_G are called (by Elana) *representation-theoretic vector bundles*. (Editor note: Xuemiao called a very similar construction 'local systems'.)

As part of the construction, we have

$$\Gamma(X // G, E_G) = \Gamma(X^{ss}, E)^G, \quad (\text{semi-invariants}). \quad (31)$$

3.3 Generalized Euler Sequence

Suppose that $V^s(\theta) \neq \emptyset$ and $V^{ss} = V^s$. Furthermore, suppose $G \curvearrowright V^s$ freely. Then $V // G$ is a smooth geometric quotient and all vector bundles descend. Consider the exact sequences

$$\begin{array}{ccccc} V^{ss} \times G & \hookrightarrow & V^{ss} & \twoheadrightarrow & V^{ss}/G \\ & & \parallel & & \\ & & \text{Differentiate} & & \\ & & \downarrow & & \\ V^{ss} \times \mathfrak{g} & \hookrightarrow & V^{ss} \times TV & \twoheadrightarrow & p^*(T(V // G)) \end{array}$$

This sequence is G -equivariant, so it descends to the sequence

$$(V^{ss} \times \mathfrak{g})/G \hookrightarrow (V^{ss} \times V)/G \twoheadrightarrow T(V // G)$$

Exercise 3.5. Show that this gives the Euler sequence for projective space.

3.4 Moduli Spaces

Moduli spaces are the solutions to *moduli problems*. These moduli problems come up in classifications. Suppose we want to classify some type of object, e.g. "curves of genus g " or "lines in projective space" or " r -dimensional subspaces of \mathbb{C}^n ". The first thing to decide is what it means for two objects to be equivalent. Then we are trying to understand the set of objects up to equivalence. In particular, we hope to put a geometric structure on this set. Part of this structure is deciding when two objects are near each other; this leads to the notion of families of objects.

Definition 3.6. A *family over U* of objects is a flat map $d : X \rightarrow U$ such that each fibre $d^{-1}(u)$ is an object.

Then the data of a moduli problem is a functor $F : C^0 \rightarrow U$, where for us $C = \text{Scheme}$, and $F(U)$ is the set of all families over U .

Definition 3.7. A *fine moduli space* is a scheme \mathcal{M} such that there is a natural isomorphism between F and the functor $\text{Hom}(-, \mathcal{M})$.

Equivalently, a scheme \mathcal{M} is a fine moduli space if it carries a *universal family*, which is a family $U \rightarrow \mathcal{M}$ such that any over family $T \rightarrow B$ is obtained by pulling back along a map $B \rightarrow \mathcal{M}$.

For many moduli problems, a fine moduli space does not exist. Thus we must be content with a *coarse* moduli space.

Definition 3.8. A scheme \mathcal{M} is a *coarse moduli space* if:

- points in \mathcal{M} correspond to isomorphism classes of objects
- any family $T \rightarrow B$ gives a map $\phi : B \rightarrow \mathcal{M}$.

Example: Consider the moduli problem of classifying surjective maps $\mathbb{C}^n \rightarrow \mathbb{C}^r$ up to change of basis. Then a family is a vector bundle $E \rightarrow X$ of rank r with a surjective map $\mathbb{C}^r \rightarrow E$ (E is globally generated). Then $Q \rightarrow \text{Gr}(n, r)$ from before is a universal family and the grassmannian is a fine moduli space. Given any globally generated vector bundle $E \rightarrow X$ generated by s_1, \dots, s_r , we have $X \rightarrow \text{Gr}(n, r)$ by

$$x \rightarrow (e_i \rightarrow s_i(x)). \quad (32)$$

3.5 Quiver Moduli Spaces

Definition 3.9. A *quiver* is a directed multigraph Q , with vertices Q_0 and edges Q_1 . There are maps $s, t : Q_1 \rightarrow Q_0$, and for any $e \in Q_1$, $s(e)$ and $t(e)$ are called the *source* and *target* of e .

Definition 3.10. A representation of a quiver Q is a set in the form

$$\{(V_i, \phi_a), \mid i \in Q_0, a \in Q_1, V_i \text{ a vector space}, \phi_a : V_{s(a)} \rightarrow V_{t(a)}\}.$$

The *dimension* of a representation is $(\dim V_i)_{i \in Q_0} \in \mathbb{N}^{|Q_0|}$.

A *morphism* of representations $(V_i, \phi_a) \rightarrow (W_i, \psi_a)$ is a collection of intertwining linear maps: $f_i : V_i \rightarrow W_i$ such that $f_i \psi_a = \phi_a f_i$.

Therefore for any quiver Q we have a moduli problem, which is to classify representations of Q with fixed dimension up to isomorphism. We can always fix $V_i = \mathbb{C}^{r_i}$, and define a space of representations

$$\text{Rep}(Q, r) = \bigoplus_{a \in Q_1} \text{Hom}(\mathbb{C}^{r_{s(a)}}, \mathbb{C}^{r_{t(a)}}). \quad (33)$$

Our isomorphisms live in the group $G = \prod_{i \in Q_0} GL(r_i)$, acting by change of basis:

$$(g_i) \circ (\phi_a) = g_{t(a)} \phi_a g_{s(a)}^{-1}.$$

The character group is $\chi(G) \cong \mathbb{Z}^{|Q_0|}$. To answer this moduli problem, our goal is to study the GIT quotient $\text{Rep}(Q, r)/G$. Using the Hilbert-Mumford criterion, we can work out a characterization of stability for this set. Let λ be a 1PS. Consider the action of λ on (V_i, ϕ_a) ; we have $\lambda : \mathbb{C}^* \rightarrow GL(V_i)$, and so

$$V_i = \bigoplus_{n \in \mathbb{Z}} V_i^{(n)},$$

where $V_i^{(n)}$ denotes the subspace on which \mathbb{C}^* has weight n . For each map ϕ_a , there is a component $\phi_a^{mn} : V_{s(a)}^{(n)} \rightarrow V_{t(a)}^{(m)}$. By construction, our 1PS acts on ϕ_a^{mn} by

$$\lambda(t) \cdot \phi_a^{mn} = t^{m-n} \phi_a^{mn}.$$

Therefore the limit $\lim_{t \rightarrow 0} \lambda(t) \cdot \phi_a$ exists iff $\phi_a^{mn} = 0$ for $m < n$ iff $\phi_a(V_{s(a)}^{(n)}) \subset \bigoplus_{m \geq n} V_{t(a)}^{(m)}$. Let $M_n := \bigoplus_{m \geq n} V_i^{(m)}$. Remark that for $n \gg 0$, $M_n = 0$, and for $0 \gg n$, $M_n = V_i$. Therefore the image of the limit $\lim_{t \rightarrow 0} \lambda(t) \cdot \phi_a$, (if it exists) will be

$$\bigoplus_{n \in \mathbb{Z}} M_n / M_{n+1}.$$

In summary, a 1PS subgroup with limit is turned into a filtration of V_i .

Suppose instead we are given a filtration $\{M_n\}$ of V_i . We can build $\bigoplus_{n \in \mathbb{Z}} M_n/M_{n+1}$ and define a 1PS λ by setting the action of \mathbb{C}^* on M_n/M_{n+1} to have weight n . Now we can connect this with the Hilbert-Mumford criterion. Consider a linearisation $\theta \in \chi(G)$. Since we are working with a non-trivial stabilizer Δ , we require $\theta(\Delta) = 1$. Consider

$$\begin{aligned} \langle \theta, \lambda \rangle &= \sum_{i \in Q_0} \sum_{n \in \mathbb{Z}} \theta_i n \dim(V_i^{(n)}) \\ (\text{exercise}) &= \sum_{i \in Q_0} \sum_{n \in \mathbb{Z}} \theta_i n \dim(M_n/M_{n+1})_i \\ &= \sum_{i \in Q_0} \sum_{n \in \mathbb{Z}} \theta_i \dim(M_n). \end{aligned}$$

This motivates us to define, for any representation $M = (M_i)_{i \in Q_0}$ of Q , $\theta(M) = \sum_{i \in Q_0} \theta_i \dim(M_i)$.

Theorem 3.11 (King). *A representation $V = (V_i, \phi_a)$ is θ -semistable if and only if $\theta(V) = 0$, and for all subrepresentations $V' \subset V$, $\theta(V') \geq 0$.*

Proof. Given a subrepresentation V' of V , define a filtration by

$$M_i = \begin{cases} 0, & i > 0 \\ V'_i, & i = 0 \\ V_i, & i < 0. \end{cases}$$

Then this also defines a 1PS λ whose limit is this filtration, and $\langle \lambda, \theta \rangle = \theta(M)$. Thus the result follows from Hilbert-Mumford criterion. \square

Theorem 3.12 (King). *The GIT quotient $M(Q, r) = \text{Rep}(Q, r) // G$ is a coarse moduli space for θ semistable reps of Q with dimension r , up to GIT equivalence.*

Exercise 3.13. Consider the quiver with $p + 1$ vertices

$$\bullet_{v_0} \xrightarrow[n \text{ edges}]{\quad} \bullet_{v_1} \longrightarrow \bullet_{v_2} \longrightarrow \dots \longrightarrow \bullet_{v_{p+1}}$$

and dimension vector $r = (1, r_1, r_2, \dots, r_p)$. Let $\theta = (-\sum_{i=1}^p r_i, 1, 1, \dots, 1)$, so that $\theta(\Delta) = 1$. Find, using King's characterization, the semi-stable locus, and use it to conclude that

$$\text{Rep}(Q, r) // G$$

is the flag variety of quotients $\mathbb{C}^n \twoheadrightarrow V_1 \rightarrow \dots \leftarrow V_p$.

When is $M(Q, r)$ a fine moduli space? To get a universal family, consider the bundle $\mathbb{C}^{r_i} \times \text{Rep}(Q, r) := W_i$ over $\text{Rep}(Q, r)$. If this descends to define a vector bundle on $M(Q, r)$, then it will be a universal family. King shows that on the stable locus of $\text{Rep}(Q, r)$, the stabilizer of the G action is Δ . This does not act trivially on the W_i , so we cannot directly apply Kempf's descent lemma, but the idea is not doomed. We need to adjust the action slightly. Suppose we had a character χ such that

$$\langle \chi, \Delta \rangle = \sum \chi_i r_i = -1.$$

Then define a new action $G \circ W_i$ where $(g_i)_{i \in Q_0} \cdot (v, \phi_a) = (\chi(g)g_i v, \chi(g)g \cdot \phi_a)$. This does not change the action on the base $\text{Rep}(Q, r)$. With this new action, the G -linearised vector bundle W_i descends to the GIT quotient, meaning $M(Q, r)$ is fine.

Thus the remaining question is 'When does such a χ exist?'. When (r_i) is a *primitive lattice element*, meaning it is not an integer multiple of some other lattice element, then χ exists:

Theorem 3.14. *Suppose r is a primitive lattice element. Then $M^s(Q, r) = \text{Rep}^s(Q, r) // G$ is a fine moduli space for θ -stable representations of Q with dimension r .*

Exercise 3.15. For the quiver Q in the previous exercise, check that the universal bundle built as above is a topological quotient bundle, and $W_0 = \mathcal{O}$.

3.6 Quiver Flag Varieties

Quiver flag varieties, introduced by A. Craw, are quiver moduli spaces with some simplifying assumptions:

1. Q is acyclic,
2. Q has a unique source vertex.

These assumptions mean we can choose an ordering of the vertices, $Q_0 = \{0, 1, \dots, p\}$, such that the number of edges $n_{ij} = 0$ if $i \geq j$. If you draw the vertices in order from left-to-right, this means the edges all point to the right. Furthermore we assume

3. The dimension vector has the form $r = (1, r_1, \dots, r_p)$; in particular this means r is a primitive lattice element.
4. The stability condition is given by $\theta = (-\sum_{i=1}^p r_i, 1, \dots, 1)$.

Remember

$$\begin{aligned} \text{Rep}(Q, r) &= \bigoplus_{a \in Q_1} \text{Hom}(\mathbb{C}^{r_{s(a)}}, \mathbb{C}^{r_{t(a)}}) \\ &= \bigoplus_{i=1}^p \text{Hom}(\mathbb{C}^{s_i}, \mathbb{C}^{r_i}), \quad s_i = \sum_{a \in Q_1, t(a)=i} r_{s(a)}. \end{aligned}$$

Proposition 3.16. *The GIT quotient $M_\theta(Q, r)$ is a smooth projective variety of dimension $\sum r_i(s_i - r_i)$. It is a fine moduli space for representations of Q .*

Proof. Our previous discussion shows that since r is a primitive lattice element, $M_\theta(Q, r)$ is a fine moduli space. By construction, we know that the GIT quotient is projective over $\text{Spec}(\mathbb{C}[R(Q, r)]^G)$, and $\mathbb{C}[R(Q, r)]^G = \mathbb{C}$, hence $M_\theta(Q, r)$ is projective.

Given a θ -stable representation (W_i) , we can use the G -action to write it as a tuple of elements in row-reduced echelon form. Then each W_i has $r_i(s_i - r_i)$ entries, and summing over i gives the dimension. It also gives smoothness as this is a chart for the GIT quotient. \square

Fix a quiver Q . Let $Q(i)$ be the subquiver of Q obtained by removing all vertices $k > i$ and their arrows. Then let $Y_i = M_\theta(Q(i), (r_0, \dots, r_i))$. Remember that we have tautological bundles W_i for each vertex $i \in Q_0$ (??). Let

$$F_i = \bigoplus_{a \in Q_1, t(a)=i} W_{s(a)}.$$

Theorem 3.17 (Craw). *There is a tower of Grassmannian bundles,*

$$\dots \rightarrow Y_2 \rightarrow Y_1 \rightarrow Y_0 = \{p\}$$

with $Y_i = \text{Gr}(F_i, r_i)$.

4 Toric Varieties as GIT Quotients

References: Chapter in Dolgachev's notes on GIT, Audin Toric Actions on Symplectic Manifolds, Cox-Little-Schenk

Our goal now is to develop as much theory as possible about toric varieties using GIT quotients instead of the theory of fans.

4.1 Building Toric Varieties with GIT

For the remainder of this section, fix:

- $K = (\mathbb{C}^*)^r$,
- L the lattice of 1PS, $L \cong \mathbb{Z}^r$ in K ,
- $M = L^\vee$, the dual lattice of characters,
- $L_{\mathbb{R}} = L \otimes \mathbb{R}$, and $L_{\mathbb{Q}} = L \otimes \mathbb{Q}$.

Let $D_1, \dots, D_m \in L^\vee$. They define an action of K on $\mathbb{C}^m = V$, by $t \cdot (v_1, \dots, v_m) = (D_i(t)v_i)_{i=1}^m$. We will assume there is no generic stabilizer. A linearization is given by a choice of $\omega \in L^\vee$, which defines

$$L_\omega = \mathbb{C}^m \times \mathbb{C}_\omega$$

We construct the GIT quotient from $\bigoplus_{d \geq 0} \Gamma(\mathbb{C}^m, L^{\otimes d}) \cong \bigoplus_{d \geq 0} \mathbb{C}[x_1, \dots, x_m]^{d\omega}$. Here superscript $d\omega$ means semi-invariants of weight $d\omega$. A polynomial is a semi-invariant of weight $d\omega$ if and only if each monomial is of the form $x_1^{a_1} \dots x_m^{a_m}$ with $\sum_{i=1}^m a_i D_i = d\omega$. So semi-invariants are integer points in the set of solutions to a linear system of equations. The *real* solutions form a cone.

Definition 4.1 (Cone). A *cone* is a set of the form

$$\{x \in \mathbb{R}^n \mid c_1 \cdot x \geq 0 \dots c_s \cdot x \geq 0, c_i \in \mathbb{R}^n\}.$$

A *rational polyhedral cone* is one where all $c_i \in \mathbb{Q}^n$.

Note that $c_i \geq 0$ and $c_i \cdot x \leq 0$ is equivalent to $c_i \cdot x = 0$, so the latter type of inequalities also forms a cone.

Given a cone C defined by $\{c_i\}$, the dual cone is

$$C^\vee = \{y \in \mathbb{R}^n \mid x \cdot y \geq 0 \forall x \in C\} = \left\{ \sum a_i \cdot c_i \mid a_i \geq 0 \right\}.$$

Note that $(C^\vee)^\vee = C$.

We will show that ring of semi-invariants is finitely generated by proving Gordon's lemma about cones.

Theorem 4.2 (Gordon's Lemma). *Let C be a rational convex polyhedral cone in \mathbb{R}^n . Then $C \cap \mathbb{Z}^n$ is a finitely generated submonoid of \mathbb{Z}^n*

Proof. C is spanned by some vectors v_1, \dots, v_k . Consider the compact box $B = \{\sum a_i v_i \mid 0 \leq a_i \leq 1\}$. By compactness $|B \cap \mathbb{Z}^n| < \infty$, so we can enumerate the points $B \cap \mathbb{Z}^n = \{w_1, \dots, w_l\}$. Then we claim that $\{w_1, \dots, w_l\}$ generates $C \cap \mathbb{Z}^n$. Suppose that $w \in C \cap \mathbb{Z}^n$, and write

$$\begin{aligned} w &= \sum d_i v_i, \quad d_i \geq 0 \in \mathbb{Q} \\ &= \sum (d_i + a_i) v_i, \quad d_i \in \mathbb{Z}, a_i \in [0, 1) \\ &= \sum d'_i v_i + \sum a_i v_i \end{aligned}$$

However the d'_i are in $\text{span}\{w_i\}$, and the left side is in $B \cap \mathbb{Z}^n$, hence equal to w_k for some k . \square

Let M be a commutative monoid. Then we will let $\mathbb{C}[M]$ denote the associated *monoid algebra*. For example $\mathbb{C}[\mathbb{Z}] = \mathbb{C}[z_1^\pm, \dots, z_i^\pm]$. Gordon's lemma tells us that our GIT data defines a submonoid of \mathbb{Z}^n , which we call S . Then

$$\bigoplus_{d \geq 0} \Gamma(\mathbb{C}^m, L^{\otimes d}) = \mathbb{C}[S] = \bigoplus_{d \geq 0} \mathbb{C}[S_d].$$

Let m_1, \dots, m_s denote the monomials generating $\mathbb{C}[S]$. For each m_j , let $I_j = \{i \mid (m_j)_i \neq 0\} \subseteq [m]$, and for any $I \subseteq [m]$ let

$$(\mathbb{C}^*)^I \times \mathbb{C}^{\bar{I}} := \{(x_i) \mid x_i \neq 0 \text{ if } i \in I\}.$$

Proposition 4.3.

$$V_\omega^{ss} = \bigcup_{j=1}^s (\mathbb{C}^*)^{I_j} \times \mathbb{C}^{\bar{I}_j}.$$

Proof. Exercise: Roughly speaking, V^{ss} are those $x \in \mathbb{C}[S]$ for which there exists a HIG f with $f(x) \neq 0$. Then evaluate $m_i(x)$ and check that it vanishes exactly on $(\mathbb{C}^*)^{I_j} \times \mathbb{C}^{\bar{I}_j}$. \square

For example, if $D_1 = \dots = D_m = 1$ and $\omega = 1$, then the generating monomials are x_1, \dots, x_m and $I_k = \{k\}$. We can characterize the semistable locus another way.

Definition 4.4. Given $I \subset m$, let $\angle_I = \{\sum_{i \in I} a_i D_i, \mid a_i > 0 \in \mathbb{Q}\}$, called the *open cone* over $\{D_i\}$. Let

$$\mathcal{A}_\omega = \{I \subset [m] \mid \omega \in \angle_I\},$$

called the set of *anticones*.

Proposition 4.5.

$$V_\omega^{ss} = \bigcup_{I \in \mathcal{A}_\omega} (\mathbb{C}^*)^I \times \mathbb{C}^{\bar{I}}.$$

Proof. Using the previous proposition, it suffices to show that $I \in \mathcal{A}_\omega \iff I = I_j$ for a generating monomial m_j .

If $I = I_j$ for some m_j , then I_j records which co-ordinates have non-zero powers in m_j ; $m_j = \prod_{i \in I_j} x_i^{a_i}$. Furthermore, since $m_j \in S$,

$$\sum_{i \in I_j} a_i D_i = d\omega$$

and thus, $\omega = \sum \frac{a_i}{d} D_i \in \angle_{I_j}$, and thus $I_j \in \mathcal{A}_\omega$.

On the other hand, if $I \in \mathcal{A}_\omega$, then $\omega = \sum_{i \in I} a_i D_i$ for some $a_i > 0 \in \mathbb{Q}$. Hence $d\omega = \omega \sum_{i \in I} a'_i D_i$, with $a_i \in \mathbb{Z}_{\geq 0}$. The monomial $\prod_{i \in I} \omega_i^{da_i}$ is a semi-invariant of weight $d\omega$, and it remains to check that it is a generating monomial of $\mathbb{C}[S]$. \square

Note: each $I \in \mathcal{A}_\omega$ defines an open subset $D(\prod_{i \in I} x_i) \subset V_\omega^{ss}$, which gives an open affine chart on the quotient.

Definition 4.6. A *toric variety* is an algebraic variety X containing an algebraic torus T as a dense open subset, such that the action $T \curvearrowright T$ extends to an action on X .

Proposition 4.7. *The GIT quotient $V // K$ is a toric variety.*

Proof. Let $\tilde{T} = (\mathbb{C}^*)^m \subset V_\omega^{ss} \subset V$. Then $(\mathbb{C}^*)^n \cong (\tilde{T}/K) \subset V // K$, where $n = \dim V // K$, and the action of \tilde{T}/K on itself naturally extends to the whole space. \square

Example: Weighted Projective Spaces.

Let $K = \mathbb{C}^*$, $D_i = a_i \in \mathbb{Z}_{\geq 0}$ and $\omega = 1$. Then $\mathcal{A}_\omega = \{I \subset [m] \mid I \neq \emptyset\}$, and $V^{ss} = \mathbb{C}^m - \{0\}$. The toric variety is

$$V //_\omega K = \mathbb{P}(a_1, \dots, a_m).$$

Note that the co-ordinate points $[l_i] \in \mathbb{C}^m / \mathbb{C}^*$ have non-trivial but finite stabilizer.

Example: Let $K = (\mathbb{C}^*)^2$ and $A = [D_1, \dots, D_m]$ be the weight matrix

$$A = \begin{bmatrix} 1 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 5 & 6 \end{bmatrix}.$$

Let $\omega = [1, 1]$. Then

$$\mathcal{A}_\omega \{ \{5\}, \{6\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\} \},$$

and any unions of these elements. Notably, it does not contain $\{1\}$, $\{2\}$, $\{3\}$, etc. The point $p = (0, 0, 0, 0, 1, 0)$ is semi-stable because it doesn't vanish in the 5th co-ordinate; the 1PS $(t, t^{-1}) \subset K$ acts trivially on p , so it is not stable.

If we instead set $\omega = [2, 1]$, then \mathcal{A}_ω contains $\{1, *\}$ and $\{2, *\}$, where $*$ $\in \{3, 4, 5, 6\}$. There are no elements with non-trivial stabilizers.

Proposition 4.8. $V_\omega^{ss} = V_\omega^s$ if and only if $\{D_i \mid i \in I\}$ contains a \mathbb{Q} -basis (of S ?) for each anticone $I \in \mathcal{A}_\omega$.

Proof. Suppose $I \in \mathcal{A}_\omega$, but I does not contain a \mathbb{Q} -basis. Then there exists a lattice point $(b_1, \dots, b_r) \in \mathbb{Z}^r$ with $b \neq 0$ and $\langle b_i, D_i \rangle = 0$ for all $i \in I$. Let $z = \sum_{i \in I} e_i \in V_\omega^{ss}$. Then $\lambda_b(\mathbb{C}^*) \subset \text{Stab}_{\mathbb{C}}(z)$ so z has a 1-dimensional stabilizer and is hence not stable.

Conversely, suppose that $z \in \mathbb{C}^m$ is strictly semi-stable. Then $z \in (\mathbb{C}^*)^I \times \mathbb{C}^{\bar{I}}$ for some $I \in \mathcal{A}_\omega$, and there is a 1PS λ such that $\lim_{t \rightarrow 0} \lambda(t)z$ exists and $\langle \lambda, \omega \rangle = 0$. Also

$$\lambda(t)z = (t^{\langle \lambda, D_i \rangle} z_i)$$

and since the limit exists, $\langle \lambda, D_i \rangle \geq 0$ for all $i \in I$. Then since $I \in \mathcal{A}_\omega$ there exists a_i with $\omega = \sum a_i D_i$. So

$$0 = \langle \lambda, \omega \rangle = \sum a_i \langle \lambda, D_i \rangle$$

Hence, $\langle \lambda, D_i \rangle = 0$ for all $i \in I$, and hence $\lambda \notin \text{span}\{D_i, i \in I\}$. \square

Proposition 4.9. *Suppose that for each $I \in \mathcal{A}$, $\{D_i, i \in I\}$ contains a \mathbb{Z} -basis. Then X_ω is smooth.*

Proof. It suffices to show that K acts trivially on the semi-stable locus, which by the previous proposition is V_ω^s . Let $z \in V_\omega^s$. Then $z \in (\mathbb{C}^*)^I \times \mathbb{C}^{\bar{I}}$ for some $I \in \mathcal{A}_\omega$. Say, without loss of generality, that $I = \{1, \dots, r, \dots, l\}$, where D_1, \dots, D_r are the \mathbb{Z} -basis. As before, if $t \in K$ then $t \cdot (z_1, \dots, z_m) = (\chi_{D_i}(t)z_i)_{i=1}^m$. If $t \cdot z = z$ then $\chi_{D_i}(t) = 1$ for all $i \in I$. This happens iff $t = 1 \in K$ (exercise). \square

Theorem 4.10. *Assume $\omega \neq 0$. Then X_ω is projective iff 0 is not contained in the convex hull of the D_i .*

Proof. By definition $\text{Proj}(\mathbb{C}[S])$ is projective over $\text{Spec}(\mathbb{C}[S])$, so we need to show $\mathbb{C}[S] = \mathbb{C}$. This happens iff the only semi-invariants of weight 0 are cconstants, iff there exists no non-constant monomial, iff there does not exist $\{a_i\}$ greater than zero with $\sum a_i D_i = 0$.

Then suppose X_ω is not projective. There exists $\{a_i\}_{i=1}^m$ such that $\sum a_i D_i = 0$ with $a_i > 0$. Thus

$$\frac{1}{\sum a_i} \sum a_i D_i = 0$$

meaning 0 is in the convex hull. \square

Gale Duality and Fans

Let W be a vector space over \mathbb{R} , spanned by β_1, \dots, β_r . Consider the exact sequence given by defining $\gamma : \mathbb{R}^r \rightarrow W$ by $e_i \rightarrow \beta_i$, and $V = \ker \gamma$.

$$V \xhookrightarrow{\delta} \mathbb{R}^r \xrightarrow{\gamma} W$$

and the dual sequence

$$W^* \xhookrightarrow{\gamma^*} \mathbb{R}^r \xrightarrow{\delta^*} V^*.$$

Define $v_i = \delta^*(e_i)$. Then $v \in V$ has

$$\delta(v) = (\langle v, v_1 \rangle, \dots, \langle v, v_r \rangle)$$

, and for $w \in W^*$,

$$\gamma^*(w) = (\langle \beta_1, w \rangle, \dots, \langle \beta_r, w \rangle).$$

Lemma 4.11. *Let $I \subset [r]$. Then $\{\beta_i \mid i \in I\}$ is a basis of W if and only if $\{v_j \mid j \in \bar{I}\}$ is a basis of V^* .*

Proof. Suppose that the $\{v_j \mid j \in \bar{I}\}$ are a basis of V^* . Let \bar{v}_j be a dual basis (of V). Then

$$\delta(\bar{v}_j) = \sum_{i=1}^r \langle \bar{v}_j, v_i \rangle e_i = e_j + \sum_{i \in I} \langle \bar{v}_j, v_i \rangle e_i,$$

and

$$0 = \gamma \circ \delta(\bar{v}_j) = \beta_j + \sum_{i \in I} \langle \bar{v}_j, v_i \rangle \beta_i.$$

Therefore $\beta_j \in \text{span}\{\beta_i \mid i \in I\}$ for all $j \notin I$; the β_i form a basis. The other direction is essentially the same argument. \square

Remark: We can replace \mathbb{R} -basis with \mathbb{Z} -basis if the β_i and v_i are lattice points.

Remark: If the $\{\beta_i \mid i \in I\}$ contain a basis, then $\{v_j \mid j \in \bar{I}\}$ are contained in an basis.

Definition 4.12. A convex cone σ is *strongly convex* if $\sigma \cap (-\sigma) = \{0\}$.

Lemma 4.13. *Consider the cones*

$$C_\beta = \{\beta_i \mid i = 1, \dots, m\} \subset W,$$

$$C_v = \{v_i \mid i = 1, \dots, m\} \subset V^*.$$

Then if $C_\beta = W$, C_v is strongly convex in V^ . If $v_i \neq 0, \forall i$, and C_v is strongly convex, then $C_\beta = W$.*

Definition 4.14. A fan Σ is a collection of cones that is closed under taking intersections and faces.

Let $A = [D_1, \dots, D_m, K]$, ω be the GIT data for a toric variety. Then we have an exact sequence defined by $A : \mathbb{Z}^m \rightarrow \mathbb{Z}^r$ by $A(e_i) = D_i$; let $M = \ker A$ and write

$$M \longrightarrow \mathbb{Z}^m \xrightarrow{A} \mathbb{Z}^r$$

Let $M^* = N$ and $L = (\mathbb{Z}^r)^\vee$. N is a free abelian group of rank $n = m - r$. If we tensor the exact sequence with \mathbb{R} and take duals we have

$$L_{\mathbb{R}} \longrightarrow \mathbb{R}^m \xrightarrow{(e_i \mapsto \rho_i)} N_{\mathbb{R}}$$

Proposition 4.15. *For the sets I_j as defined above, let the anticone over I_j be*

$$\sigma_{\bar{I}_j} = \left\{ \sum a_i \rho_i \mid a_i > 0, i \in \bar{I}_j \right\}.$$

Then $\Sigma_\omega = \{\sigma_{\overline{I}_j} \mid I_j\} (?)$ is a fan

For proof see notes of Dolgachev.

Exercise: can we substitute \bar{I}_j with $I \in \mathcal{A}_\omega$?

Now we've defined a way to build a fan Σ_ω out of GIT data for a toric variety. Every fan can be used to build a toric variety directly. Loosely, this is done by gluing together $\text{Spec}(\mathbb{C}[[\sigma^\vee \cap M]]) =: X_\sigma$ along the intersections $\sigma_1 \cap \sigma_2$. If $\sigma = \sigma_1 \cap \sigma_2$, then X_σ is an affine open set in $\sigma_1 \cap \sigma_2$. The scheme X_Σ built this way is a more general construction, but if your fan $\Sigma = \Sigma_\omega$ for some GIT data, then the toric built this way agrees with the one built by GIT.

Definition 4.16. A fan Σ is called

- *complete* if its support is \mathbb{R}^n ,
- *simplicial* if each cone is simplicial, meaning the rays of the cone are contained in a \mathbb{Q} basis,
- *smooth* if each cone is smooth, meaning the rays of the cone are contained in a \mathbb{Z} basis.

Proposition 4.17. *Let X_Σ be the toric variety built from a fan Σ . Then*

- X_σ is projective iff Σ is complete,
- X_σ is an orbifold iff Σ is simplicial,
- X_σ is smooth iff Σ is smooth.

4.2 Example: \mathbb{P}^1

As an addendum to Elana's notes, we will go through the above theory for the case of the Riemann sphere \mathbb{P}^1 . First, we will construct \mathbb{P}^1 as a GIT quotient. Then we will compute the fan corresponding to the GIT data that we used, and verify that the toric variety of that fan is again \mathbb{P}^1 .

Our GIT data consists of $V = \mathbb{C}^2$, $K = \mathbb{C}^*$, with action $\mathbb{C}^* \curvearrowright \mathbb{C}^2$ defined by weights $A = [D_1, D_2] = [1, 1]$. Recall that $D_i \in \chi(\mathbb{C}^*) = \mathbb{Z}$. Then, we can let \mathcal{L} be the trivial line bundle $\mathcal{L} = \mathbb{C} \times \mathbb{C}^2$, and linearize our action by $\omega = 1 \in \chi(\mathbb{C}^*)$. This GIT data is equivalent to saying that our action is:

$$t \cdot ((z, w), l) = \left((t^{D_1} z, t^{D_2} w), t^\omega l \right) = ((tx, ty), tl),$$

for $t \in \mathbb{C}^*$, $((x, y), l) \in \mathcal{L}$. Then, we need to compute the ring of degree d semi-invariants, $\Gamma(\mathbb{C}^2, \mathcal{L})^{d\omega}$. Note that $\Gamma(\mathbb{C}^2, \mathcal{L}) = \mathbb{C}[x, y]$. The degree d semi-invariants are $f \in \mathbb{C}[x, y]$ with

$$f(t \cdot (x, y)) = t^{d\omega} f(x, y).$$

In this case we need those such that

$$f(tx, ty) = t^d f(x, y).$$

This is exactly the degree d homogeneous polynomials. Thus we have that $\bigoplus_{d \geq 0} \Gamma(\mathbb{C}^2, \mathcal{L})^{d\omega} \cong \mathbb{C}[x, y]$ with the usual grading, and

$$\mathbb{C}^2 // \mathbb{C}^* := \text{Proj} \left(\bigoplus_{d \geq 0} \Gamma(\mathbb{C}^2, \mathcal{L})^{d\omega} \right) = \text{Proj}(\mathbb{C}[x, y]) = \mathbb{P}^1. \quad (34)$$

Next, let us build the fan corresponding to this GIT data. Proposition 4.15 tells us it is built out of cones defined by some vectors $\rho_i \in N$, where $N := M^\vee = (\ker A)^\vee$. Let us find the ρ_i . Consider:

$$\begin{array}{ccc} \ker A & \hookrightarrow & \mathbb{Z}^2 \rightarrow \mathbb{Z} \\ & & e_i \xrightarrow{A} D_i \end{array}$$

Since $A = [1, 1]$, $\ker A = \{[l, -l] \mid l \in \mathbb{Z}\}$. Thus the map $\mathbb{Z} \cong \ker A \rightarrow \mathbb{Z}^2$ is given by $(l \rightarrow [l, -l])$ or as a matrix, $[1, -1]$. Then the dual map is given by $[1, -1]^T$;

$$\begin{array}{ccc} \mathbb{Z}^2 & \rightarrow & \mathbb{Z} \\ v & \mapsto & [1, -1]^T v. \end{array}$$


Thus $e_1 \rightarrow 1$ and $e_2 \rightarrow -1$, and hence $\rho_1 = 1, \rho_2 = -1$. Next, we need to find the set \mathcal{A}_ω , so we know which anticones we need to include. Recall that

$$\mathcal{A}_\omega = \left\{ I \subset [m] \mid \omega = \sum_{i \in I} a_i D_i \text{ for some } a_i > 0 \in \mathbb{Q} \right\}.$$

In this case, $[m] = [2] = \{1, 2\}$, so we have only four possible cones. Since $D_1 = D_2 = 1$ and $\omega = 1$, all of $\{1\}, \{2\}$ and $\{3\}$ are in \mathcal{A}_ω . $I = \emptyset$ is not. Now we are ready to apply proposition 4.15, which tells us our fan Σ consists of three cones:

$$\sigma_1 = \{a\rho_2 \mid a > 0\}, \quad \sigma_2 = \{a\rho_1 \mid a > 0\}, \quad \sigma_{12} = \{0\}.$$

Since $\rho_2 = -1$ and $\rho_1 = 1$, Σ looks like:



Although it is outside the scope of the course content, we will conclude by computing the toric variety corresponding to Σ to verify it is \mathbb{P}^1 . See Fulton pages 3-6 for details. To each cone in Σ , we associate an affine variety. Note that in this case, σ_1 and σ_2 are their own dual cones, and $\sigma_{12}^\vee = \mathbb{Z}$. Thus, σ_1^\vee is generated (over $\mathbb{Z}_{\geq 0}$) by $-1 = -e_1$, σ_2^\vee is generated by $1 = e_1$ and σ_{12}^\vee is generated by $\{1, -1\} = \{e_1, -e_1\}$. Thus we have

$$U_1 = \text{Spec}(\mathbb{C}[x^{-1}]), \quad U_2 = \text{Spec}(\mathbb{C}[x]), \quad U_{12} = \text{Spec}(\mathbb{C}[x, x^{-1}]).$$

By definition, we glue U_1 and U_2 along U_{12} . Thus we have the diagram

$$\begin{array}{c} \mathbb{C} \rightarrow \mathbb{C}^* \leftarrow \mathbb{C} \\ x^{-1} \rightarrow x \leftarrow x. \end{array}$$

Giving exactly \mathbb{P}^1 .

4.3 Delzant Polytopes

todo

5 Bundles and Divisors

5.1 Line Bundles

Let $V = \mathbb{C}^m$, $K = (\mathbb{C}^*)^k$, $A = [D_1, \dots, D_m]$ and ω be the GIT data for a toric variety X_ω . We will consider line bundles on X_ω built as quotients of line bundles on \mathbb{C}^m . Let U_ω denote the semi-stable locus of V , and consider the trivial line bundle

$$U_\omega \times \mathbb{C} \rightarrow U_\omega$$

with K -action given by A on U_ω and $d \in \chi(K)$ on \mathbb{C} . Kempf's descent lemma tells us that this linearized trivial line bundle descends to form $L_d \rightarrow X_\omega$ if the stabilizers of elements in U_ω act trivially on the fibres of $U_\omega \times \mathbb{C}$.

Example: Let $m = 3, k = 1$, $A = [1, 1, 2]$, $\omega = 1$, so that $X_\omega = \mathbb{P}(1, 1, 2)$. Then the anti-cones are all non-empty sets: $\mathcal{A}_\omega = \{I \subset [m] \mid I \neq \emptyset\}$. The point $z = [0, 0, 0, 1] \in \mathbb{C}^3$ has stabiliser

$$K_z = \{t \mid [0, 0, t^2] = [0, 0, 1]\} = \{\pm 1\} \cong \mathbb{Z}_2$$

If we let $U_\omega \times \mathbb{C}$ be linearized by weight $d = 1$, then the action of K_z on the fibre at z is non-trivial. However if we pick $d = 2$ then the action on the fibre is trivial. Thus the first does not descend to a line bundle, but the second does.

Example: Let $k = 2$, $m = 5$ and $\omega = [1, 1]^T$ and

$$A = \begin{bmatrix} 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 \end{bmatrix}.$$

Then $[0, 0, 0, 0, z_5, z_6]$ is a semistable point, with stabilizer $K_z = \{(\lambda, \lambda^{-1}) \mid \lambda \in \mathbb{C}^*\} \subset K$. If we take $d = r[1, 1]^T$, then the action of K_z on the fibre is $(\lambda, \lambda^{-1}) \cdot l \rightarrow \lambda^r \lambda^{-r} l = l$, which is trivial, but for any other linearization the action is non-trivial and the bundle will not descend to X_ω .

Can we find an easier criterion for when a weight will descend to define a line bundle? Observe the stabilizer at some semi-stable element z depends only on the set $I = \{i \mid z_i = 0\}$, which corresponds to some $\bar{I} \in \mathcal{A}_\omega$. This means we should be able to find a condition on anti-cones which tells us if a bundle descends. The stabilizer is the kernel of the map $f_{\bar{I}} : K \rightarrow (\mathbb{C}^*)^{\bar{I}}$ given by

$$\lambda \rightarrow (\chi_{D_i}(\lambda))_{i \in \bar{I}}.$$

Therefore $d \in \chi(X)$ defines a line bundle if $\forall J \in \mathcal{A}_\omega$, $d(\ker f_J) = \{e\}$.

We can rewrite $\ker(f_J) \cong \mathbb{Z}^k / \langle D_i : i \in J \rangle$. The natural surjection

$$\mathbb{Z}^k \rightarrow \mathbb{Z}^k / \langle D_i : i \in J \rangle$$

factors through

$$\mathbb{Z}^k / \langle d \rangle \mathbb{Z}^k / \langle D_i : i \in J \rangle$$

if and only if $d \in \langle D_i : i \in J \rangle$. Therefore

Proposition 5.1. *The bundle $U_\omega \times \mathbb{C}$ linearized by $d \in \chi(K)$ descends to a line bundle on X_ω if and only if, for all $J \in \mathcal{A}_\omega$, $d \in \langle D_i : i \in J \rangle$.*

Furthermore, we can convert this into a statement about the fan Σ_ω defining X_ω . Let $\rho_1, \dots, \rho_n \in M$ be the rays of the fan, and choose a lift of $d \in \chi(K) = \mathbb{Z}^k$ to \mathbb{Z}^m . Recall that the map $\mathbb{Z}^m \rightarrow \mathbb{Z}^k$ is given by the weight matrix A , so a lift is equivalent to a choice $(a_1, \dots, a_m) \in \mathbb{Z}^m$ such that

$$A(a_1, \dots, a_m) = \sum_{i=1}^m a_i D_i = d.$$

Associated to such a lift, we can define a map $M \rightarrow \mathbb{Z}^m$ by sending $\rho_i \rightarrow a_i$. Note that such a map can fail to be well defined. Consider the example of $\mathbb{P}(1, 1, 2)$ and linearization $d = 1$ from before. From the weight matrix A , we can find that the rays of the cone are

$$\rho_1 = [1, 1], \quad \rho_2 = [-1, 1], \quad \rho_3 = [0, -1]$$

Pick the lift $(1, 2, -1) \in \mathbb{Z}^3$. This is a valid lift because $1(1) + 2(1) - 1(2) = 1 = d$. Then the map $f : M_\mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$f(\rho_1) = 1, \quad f(\rho_2) = 2, \quad f(\rho_3) = 1.$$

This is not \mathbb{R} -linear. For example, if f was linear, then $\rho_3 = -\frac{1}{2}(\rho_1 + \rho_2)$, and hence

$$f\left(-\frac{1}{2}(\rho_1 + \rho_2)\right) = -\frac{1}{2}(f(\rho_1) + f(\rho_2)) = -\frac{3}{2}.$$

This contradicts that $f(\rho_3) = 1$.

Lemma 5.2. *Let (a_1, \dots, a_m) and (b_1, \dots, b_m) be two lifts of $d \in \mathbb{Z}^k$. Let $m_a, m_b : \mathbb{R}^n \rightarrow \mathbb{R}$ be the maps defined by $m_a(\rho_i) = a_i$ and $m_b(\rho_i) = b_i$. Then m_a is a linear map if and only if m_b is a linear map.*

Proof. By definition, $\sum a_i D_i = \sum b_i D_i = d$. Therefore $\sum (a_i - b_i) D_i = 0$, and $(a_1 - b_1, \dots, a_m - b_m) \in \ker(\mathbb{Z}^m \xrightarrow{A} \mathbb{Z}^k)$. By linearity of A this means $m_a \circ A = 0 \iff m_b \circ A = 0$. ??? i don't understand \square

Proposition 5.3. *Let $U_\omega \times \mathbb{C}$ be linearized by a weight $d \in \chi(K) = \mathbb{Z}^k$. This descends to a line bundle on X_ω if and only if, for all $\sigma_I \in \Sigma_\omega$ and any lift $(a_i)_{i=1}^m$ of d , the map $\rho_i \rightarrow a_i, i \in I$ defines a linear map.*

Equivalently, there exists $m_{\sigma_I} \in M_\mathbb{R}$ such that $\langle m_{\sigma_I}, \rho_i \rangle = a_i$.

Proof. From the previous proposition, we know the bundle descends if for all $I \in \mathcal{A}_\omega$, $d \in \langle D_i : i \in I \rangle$. The cones σ_I correspond 1-1 with the anticones $\bar{I} \in \mathcal{A}_\omega$, so we want to show that

$$d \in \langle D_i : i \in \bar{I} \rangle \iff (\rho_i \rightarrow a_i)_{i \in I} \text{ is a linear map.}$$

If $d \in \langle D_i : i \in \bar{I} \rangle$, then $d = \sum_{i \in \bar{I}} a_i D_i$ for some $a_i \in \mathbb{Z}$. Therefore letting $a_i = 0$ for $i \in I$, $(a_i)_{i=1}^m$ is a lift of d . The corresponding map is the zero map, which is linear.

On the other hand, if $d = \sum_{i=1}^1 a_i D_i$ is a lift defining a linear map, then let

$$B_I := \begin{bmatrix} \rho_{i_1} \\ \dots \\ \rho_{i_k} \end{bmatrix}_{i_j \in I}$$

The linearity means that

$$B_I^T \begin{bmatrix} b_1 \\ \dots \\ b_n \end{bmatrix} = 0 \implies [a_1, \dots, a_m] \cdot [b_1, \dots, b_m] = 0$$

I.e. $\ker(B_I^T) \subset \ker([a_1, \dots, a_m])$. Moreover

$$\ker(B_I^T) = \text{Im}(A^T) \cap \langle e_i \mid i \in I \rangle^T = [\ker(A) + \langle e_i \mid i \in I \rangle]^T.$$

Also, $[a_1, \dots, a_m]^T \in (\ker A + \langle e_i \mid i \in I \rangle)$. Thus $d = \sum a_i D_i = \sum_{k \in I} b_k D_k$, meaning $d \in \langle D_i \mid i \in I \rangle$. \square

Definition 5.4. An *integral support function* is a continuous map $g : N \rightarrow \mathbb{R}$ such that

1. g is linear on each cone,
2. g takes integral values on $|\Sigma_\omega| \cap \mathbb{Z}^n$.

With this definition, the previous proposition says that line bundles over X_ω are defined by integral support functions on Σ_ω . If one drops condition 2, then instead of line bundles one obtains \mathbb{Q} -Cartier divisors up to linear equivalence.

Given an integral support function g on a fan $\Sigma_\omega \subset \mathbb{R}^n$, then we can consider the graph of g , $\Sigma_\omega^g \subset \mathbb{R}^n \times \mathbb{R}$. This defines a new fan, with rays

$$\{\rho'_i = (\rho_i, g(\rho_i))\} \cup \{\rho'_{m+1} = (\vec{0}, 1)\}$$

and cones

$$\sigma_i \in \Sigma_\omega^g \iff \sigma_{I-m_1} \in \Sigma_\omega.$$

The projection map $\mathbb{Z}^n \times \mathbb{Z} \rightarrow \mathbb{Z}^n$ defines a map of fans $\Sigma_\omega^g \rightarrow \Sigma_\omega$, and hence induces a map of toric varieties. We claim that the toric variety corresponding to Σ_ω^g is the total space of the line bundle defined by g .

Proposition 5.5. Let g be an integral support function g on a fan Σ_ω . Let X_ω^g be the fan associated to the graph fan Σ_ω^g . Then the induced map of toric varieties is a line bundle $X_\omega^g \rightarrow X_\omega$.

5.2 Weil and Cartier Divisors

Let X be a normal irreducible variety. A prime divisor $D \subset X$ is an irreducible subvariety with codimension 1. Define the sheaf $\mathcal{O}_{X,D}$ by

$$\mathcal{O}_{X,D}(U) = \mathcal{O}_X(U \cap D).$$

Then $\mathcal{O}_{X,D}(X)$ is a discrete valuation ring, with valuation $v_D(f)$ equal to the order of vanishing along D . Let $\text{Div}(X)$ denote the free abelian group over the prime divisors.

Definition 5.6. A *Weil divisor* is an element of $\text{Div}(X)$. If $D = \sum a_i D_i$, where D_i are prime divisors and all $a_i > 0$, then we say D is *effective*.

Definition 5.7. Let $f \in \mathcal{O}_X$. Then $\text{Div}(f) := \sum_{\text{prime divisors } D} v_D(f) D$. Let $\text{Div}(\mathcal{O}_X) := \text{Div}_0(X)$, called the group of principal divisors.

We say D_1 is *linearly equivalent* to D_2 or write $D_1 \sim D_2$ if $D_1 - D_2 \in \text{Div}_0(X)$.

Definition 5.8. A Weil divisor is called *Cartier* if it is locally principal. That is, there exists an open cover $\{U_i\}$ of X with $D|_{U_i} = \text{Div}(f_i)$ for some $f_i \in \mathcal{O}_X(U_i)$. We denote the group of Cartier divisors by $\text{CDiv}(X) \subset \text{Div}(X)$.

Definition 5.9. The *class group* of X is

$$\text{Cl}(X) := \text{Div}(X) / \text{Div}_0(X).$$

The *Picard group* of X is

$$\text{Pic}(X) := \text{CDiv}(X) / \text{Div}_0(X).$$

Remark 5.10. To any Weil divisor D we can associate a coherent sheaf $\mathcal{O}_X(D)$ by

$$\mathcal{O}_X(D)(U) = \{f \in \mathcal{O}_X(U) \mid \text{Div}(f) + D|_U \geq 0\}.$$

If D_1 and D_2 are linearly equivalent, then $\mathcal{O}_X(D_1) \cong \mathcal{O}_X(D_2)$. If D is Cartier, then $\mathcal{O}_X(D)$ will also be locally free, hence defining a line bundle. This means that $\text{Pic}(X)$ can also be viewed as the group of isomorphism classes of line bundles on X .

Now we will restrict our focus back to toric varieties. Let $A = [D_1, \dots, D_m]$, K, ω be GIT data for a quotient of $V = \mathbb{C}^m$. Let $S = \{i \in [m] \mid \{i\}^c \notin \mathcal{A}_\omega\}$, and suppose $S = \emptyset$. Consider the set $\{x_i = 0\} \cap V^{ss}$. It is closed, K invariant and the image in the quotient will be a prime divisor \tilde{D}_i . Now let $T \cong (\mathbb{C}^*)^{m-k}$ be the torus inside X_ω , and let $\chi_m \in \chi(T)$. This defines a non-vanishing function $N \rightarrow \mathbb{R}$ by $\langle m, \cdot \rangle$.

Proposition 5.11. *The valuation of m along \tilde{D}_i is given by*

$$v_{\tilde{D}_i}(m) = \langle m, \rho_i \rangle.$$

Proof. We can pick a basis of N in which $\rho_i = e_i$. Our assumption that $S = \emptyset$ means that associated to ρ_i is an affine open set U_i corresponding to the anticone $\{i\}^c$;

$$U_i = [(\mathbb{C}^*)^{\{i\}^c} \times \mathbb{C}^{\{i\}}] / K \cong \mathbb{C} \times (\mathbb{C}^*)^{n-1}$$

WLOG we set $i = 1$. Then

$$U_i = \text{Spec}(x_1, x_2^\pm, \dots, x_n^\pm).$$

Let $f \in \mathbb{C}[x_1, \dots, x_n]$. Then $v_{\tilde{D}_i}(f) = l \in \mathbb{Z}$ and we can write $f = u^l \frac{g}{h}$, with $g/h \in \mathbb{C}[x_2^\pm, \dots, x_n^\pm]$. We can also write

$$\chi^m = \prod_{i=1}^k x_i^{\langle m, \rho_i \rangle} = x_1^{\langle m, \rho_1 \rangle} \prod_{i=2}^k x_i^{\langle m, \rho_i \rangle}.$$

Letting $f = \chi^m$ and comparing these equations, we see $\langle m, e_i \rangle = l = v_{\tilde{D}_i}(\chi^m)$. □

Proposition 5.12 (See CLS). *Let $\chi^m \in \chi(K)$. Then $\text{Div}(\chi^m) = \sum \langle m, \rho_i \rangle \tilde{D}_i$.*

Let $\text{Div}_T(X_\omega)$ denote the T -invariant divisors.

Exercise 5.13. Show that $\text{Div}(X_\omega) = \sum_{i=1}^m \mathbb{Z} \tilde{D}_i$.

Theorem 5.14. *There exists a short exact sequence*

$$\chi(T) \hookrightarrow \text{Div}_T(X_\omega) \rightarrow \text{Cl}(X_\omega).$$

Proof. There's an exact sequence (CLS 4.0.20)

$$\text{Div}_T(X_\omega) \rightarrow \text{Cl}(X_\omega) \xrightarrow{\pi} \text{Cl}(T) \rightarrow 0$$

Moreover, $\text{Cl}(T) = 0$. Hence for exactness it remains to show that $\chi(T) = \ker(\pi)$. Certainly, $\chi(T) \subset \ker(\pi)$, since $\text{Div}(\chi^m) \in \text{Div}_T(X_\omega) = \ker \pi$.

Suppose that $\pi(D) = 0$ for some $D \in \text{Cl}(X_\omega)$. By exactness, $D \in \text{Div}_T(X_\omega)$, so $D = \text{Div}(f)$ for some $f \in \mathbb{C}[X_\omega]^*$. This means the support of D does not intersect T , i.e. $\text{Div}(f)|_T = 0$. Hence, we can say $f \in \mathbb{C}[T]^*$, meaning f is a multiple of χ^m for some m ; $\ker \pi \subset \chi(T)$.

Finally, we need to show that $\iota : \chi(T) \rightarrow \text{Div}_T(X_\omega)$ is injective.

$$\iota(\chi_m) = \sum \langle m, \rho_i \rangle D_i = 0$$

Therefore $\langle m, \rho_i \rangle = 0$ for each ρ_i . The ρ_i are a spanning set of M , hence $m = 0$. □

Fix a divisor $D = \sum a_i \tilde{D}_i$, and define $P_D = \{m \in M_\mathbb{R} \mid \langle m, \rho_i \rangle \geq -a_i \forall i\}$. P_D might not be a polytope, however its integer points are still important.

Proposition 5.15.

$$\Gamma(X_\omega, \mathcal{O}_{X_\omega}(D)) = \bigoplus_{m \in P_D \cap M} \mathbb{C} \cdot \chi^m.$$

Proof. Suppose $f \in \Gamma(X_\omega, \mathcal{O}_{X_\omega}(D))$. Then by definition, $\text{Div}(f) + D \geq 0$. However

$$(\text{Div}(f) + D)_T = \text{Div}(f), \implies \text{Div}(f) \geq 0$$

Therefore, by the previous theorem? (TODO: ask Elana how this proof goes) □

Example: Consider the *Hirzebruch surface* H_2 , defined by the GIT weight matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 2 \end{bmatrix}, \quad B^T = \begin{bmatrix} -1 & 2 \\ 0 & 1 \\ 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

With stability condition $\omega = (\omega_1, \omega_2)$ with $2\omega_2 > \omega_1$. (TODO add picture of fan and secondary fan) The minimal anticones are

$$\mathcal{A}_\omega \supset \{(1\ 4), (3\ 4), (1\ 2), (2\ 3)\}.$$

Each anticone contains a \mathbb{Z} -basis of N , and hence all divisors are Cartier, and any choice of linearizing weight d will define a line bundle.

Consider the divisor $D_a := a\tilde{D}_1 + \tilde{D}_2$ for some $a \in \mathbb{Z}$. Define P_a from this divisor;

$$m \in P_a \iff (-m_1 + 2m_2 \geq -a) \wedge (m_2 \geq -1) \wedge (m_1 \geq 0) \wedge (-m_2 \geq 0).$$

When $a \geq 3$, P_a is a polytope whose corresponding toric variety recovers H_2 exactly. However if $a = 2$, then the polytope is a triangle, which defines \mathbb{P}^2 . If $a = 1$, then P_a is no longer a lattice polytope. These choices of a correspond to moving through the chambers of the secondary fan. In summary, P_D has a normal fan which recovers the original fan whenever $\sum a_i D_i \in \bigcap_{I \in \mathcal{A}_\omega} \angle_I$, or equivalently whenever $\sum a_i D_i$ defines an ample line bundle.

Theorem 5.16. 1. X_ω is smooth $\iff \text{Pic}(X_\omega) = \text{Cl}(X_\omega)$.
 2. X_ω is simplicial \iff every Weil divisor is \mathbb{Q} -Cartier.

5.3 Homogeneous Co-ordinate Ring

Consider GIT data $A = [D_1, \dots, D_m]$, $K = (\mathbb{C}^*)^k$ and ω . We know the GIT quotient is defined by

$$\text{Proj}\left(\bigoplus_{l \geq 0} \mathbb{C}[x_1, \dots, x_m]^{l\omega}\right)$$

Let $R = \bigoplus_{l \geq 0} \mathbb{C}[x_1, \dots, x_m]^{l\omega}$, and for any $\chi \in \chi(K)$, let R_χ denote the χ -graded part of R . For example, for $\mathbb{P}^2 \times \mathbb{P}^2$ whose weight matrix is $A = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{bmatrix}$ with linearization $\omega = (1, 1)$, we can consider things like $R_{\begin{bmatrix} 3 \\ 2 \end{bmatrix}}$. This ring R is called the *homogeneous co-ordinate ring* or the *Cox ring* of X_ω , and it behaves like the co-ordinate ring of \mathbb{P}^n .

The one may ask: When does $\{f = 0\} \subset X_\omega$ make sense for some $f \in \mathbb{C}[x_1, \dots, x_m]$? Well, this will make sense if $\{f = 0\} \subset \mathbb{C}^m$ passes to the GIT quotient, which requires that $\lambda \cdot f = \chi(\lambda)f$. Let $I \subset R$ be a homogeneous ideal. Let $\pi : (\mathbb{C}^m)^{ss} \rightarrow X_\omega$ be the quotient map, and let

$$V(I) = \{y \in X_\omega \mid \exists x \in \pi^{-1}(y), f(x) = 0 \forall f \in I\}$$

Then we have:

Proposition 5.17. • $V(I)$ is a closed subvariety of X_ω ,
 • All closed varieties are equal to $V(I)$ for some homogeneous ideal I .